

Extremal Multicenter Black Holes: Nilpotent Orbits and Tits Satake Universality Classes

Pietro Fré^{a1}, Alexander S. Sorin^b

^a *Dipartimento di Fisica, Università di Torino,
& INFN - Sezione di Torino
via P. Giuria 1, I-10125 Torino, Italy
fre@to.infn.it*

^b *Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research,
141980 Dubna, Moscow Region, Russia
sorin@theor.jinr.ru*

Abstract

Four dimensional supergravity theories whose scalar manifold is a symmetric coset manifold $U_{D=4}/H_c$ are arranged into a finite list of Tits Satake universality classes. Stationary solutions of these theories, spherically symmetric or not, are identified with those of an euclidian three-dimensional σ -model, whose target manifold is a Lorentzian coset $U_{D=3}/H^*$ and the extremal ones are associated with H^* nilpotent orbits in the K^* representation emerging from the orthogonal decomposition of the algebra $U_{D=3}$ with respect to H^* . It is shown that the classification of such orbits can always be reduced to the Tits-Satake projection and it is a class property of the Tits Satake universality classes. The construction procedure of Bossard et al of extremal multicenter solutions by means of a triangular hierarchy of integrable equations is completed and converted into a closed algorithm by means of a general formula that provides the transition from the symmetric to the solvable gauge. The question of the relation between H^* orbits and charge orbits \mathbf{W} of the corresponding black holes is addressed and also reduced to the corresponding question within the Tits Satake projection. It is conjectured that on the vanishing locus of the Taub-NUT current the relation between H^* -orbit and \mathbf{W} -orbit is rigid and one-to-one. All black holes emerging from multicenter solutions associated with a given H^* orbit have the same \mathbf{W} -type. For the S^3 model we provide a complete survey of its multicenter solutions associated with all of the previously classified nilpotent orbits of $\mathfrak{sl}(2) \times \mathfrak{sl}(2)$ within $\mathfrak{g}_{2,2}$. We find a new intrinsic classification of the W -orbits of this model that might provide a paradigm for the analogous classification in all the other Tits Satake universality classes.

¹Prof. Fré is presently fulfilling the duties of Scientific Counselor of the Italian Embassy in the Russian Federation, Denezhnij pereulok, 5, 121002 Moscow, Russia.

Contents

1	Introduction	3
2	The general framework	6
2.1	The σ -model approach to extremal black holes	8
2.1.1	General structure of the $\mathbb{U}_{D=3}$ Lie algebra	9
2.1.2	General form of the three-dimensional σ -model	10
2.2	Oxidation rules for extremal multicenter black holes	11
2.2.1	Reduction to the spherical case	12
2.3	A counter example: the extremal Kerr metric	13
3	Construction of multicenter solutions associated with nilpotent orbits	15
3.1	The coset representative in the symmetric gauge	15
3.2	Transformation to the solvable gauge	17
3.3	Extraction of the three dimensional scalar fields	18
3.4	General properties of the black hole solutions and structure of their poles	19
4	The example of the S^3 model: classification of the nilpotent orbits	22
5	Explicit construction of the multicenter Black Holes solutions of the S^3 model	27
5.1	The Very Small Black Holes of \mathcal{O}_{11}^4	27
5.2	The Small Black Holes of \mathcal{O}_{11}^2	32
5.3	The Large BPS Black Holes of \mathcal{O}_{11}^3	36
5.4	BPS Kerr-Newman solution	41
5.5	The Large non BPS Black Holes of \mathcal{O}_{22}^3	43
5.6	The Largest orbit \mathcal{O}_{11}^1	47
6	Classification of the sugra-relevant symmetric spaces and discussion of their general properties	50
6.1	Tits Satake projection	52
6.2	Classification of the sugra-relevant symmetric spaces	54
6.3	Dynkin diagram analysis of the principal models	55
7	Tits Satake decompositions of the \mathbf{W} representations	62
7.1	Universality class $\mathfrak{sp}(6, \mathbb{R}) \Rightarrow \mathfrak{f}_{4(4)}$	62
7.1.1	$\mathfrak{su}(3, 3)$ model	63
7.1.2	$\mathfrak{so}^*(12)$ model	63
7.1.3	$\mathfrak{e}_{7(-25)}$ model	64
7.2	Universality class $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 3) \Rightarrow \mathfrak{so}(4, 5)$	64
7.3	Universality class $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, 7) \Rightarrow \mathfrak{so}(8, 9)$	65
7.4	The universality classes $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, n) \Rightarrow \mathfrak{so}(8, n+2)$ with $n \leq 5$	65
7.5	\mathbf{W} -representations of the maximally split non exotic models	66
8	Tits Satake reduction of the \mathbb{H}^* subalgebra and of its representation \mathbb{K}^*	69

9	The general structure of the $\mathbb{H}^* \oplus \mathbb{K}^*$ decomposition in the maximally split models	71
10	\mathbb{K}^* orbits versus W-orbits	73
11	Conclusions and Perspectives	74
12	Acknowledgements	75
A	The complete form of the fields for the $\mathcal{O}_{3 11}$ orbit	76

1 Introduction

The classical solutions of supergravity models that, in the gravitational sector, include metrics of black hole type, have attracted a lot of interest in the course of the last fifteen-seventeen years, with various peaks of attention and research activity when some new conceptual input was introduced, which has occurred repeatedly.

The milestones in this new age of black hole physics have been:

- 1) The discovery of the attraction mechanism in the evolution of scalar fields [1].
- 2) The statistical interpretation of black hole entropy in terms of string microstates [2, 3].
- 3) The ample development of black hole study [4, 5, 6, 7, 8, 9, 10, 11] within the framework of special Kähler geometry, [12, 13, 14] using in particular the first order equations that follow from the preservation of supersymmetry. This meant an in depth study of BPS black holes and of the relation of symplectic invariants of the duality groups with the black hole entropy [15].
- 4) The construction of black hole microstates in suitable conformal field theories [16].
- 5) The discovery of *fake superpotentials* and of new attraction flows for non supersymmetric, non BPS black holes [17, 18, 19, 20, 21].
- 6) The introduction of a three-dimensional σ -model approach for the derivation of black hole solutions [22] and the association of these latter with nilpotent orbits of the U -duality group in the extremal case [23, 24, 25].
- 7) The complete integrability of the supergravity equations in the spherical symmetric case and their association with a Lax pair formulation, in the case where the scalar manifold is a symmetric homogenous space U/H [23, 24, 25, 26, 27, 28, 29].

What mainly concerns the present paper are the developments related with points 6) and 7) of the above list.

Concerning spherical symmetric supergravity black holes in the last couple of years there were significant parallel developments based on:

- A** The purely $D = 4$ approach centered on the geodesic potential for radial flows, its reexpression in terms of a *fake superpotential* and the analysis of critical points of the latter, corresponding to black hole horizons [5, 30, 31]. In this approach the classification of black holes is reduced to the classification of orbits of the symplectic \mathbf{W} -representation, namely the representation of the $D = 4$ duality group $U_{D=4}$, encoding the electromagnetic charges of the supergravity model.
- B** The $D = 3$ approach centered on the reduction of supergravity field equations to those of a $D = 3$ σ -model with a Lorentzian target manifold which, for scalar manifolds that are symmetric coset spaces $U_{D=4}/H_c$, is also a symmetric coset $U_{D=3}/H^*$, the isotropy group H^* being a non-compact real section of the complexification of the maximal compact subgroup of $U_{D=3}$. In this approach, which leads to a Lax pair formalism and to the development of explicit analytic formulae for the general integral [26, 23, 25], the classification of black holes is reduced to the classification of orbits of the non compact stability group H^* in the \mathbb{K}^* representation that corresponds to the orthogonal decomposition of the $D = 3$ Lie algebra: $U_{D=3} = \mathbb{H}^* \oplus \mathbb{K}^*$.

The main result of approach A) has been the classification of critical points in terms of certain special geometry invariants that are able to distinguish large and small black holes, BPS and non-BPS ones. Obviously the structure of these invariants depend on the electromagnetic charges which determine the geodesic potential and hence on the orbit of the \mathbf{W} representation yet such a link is not so clear in the $D = 4$ approach. On the other hand, also in the $D = 3$ approach the relation between the \mathbb{H}^* orbits and the \mathbf{W} orbits of the resulting black hole charges was not systematically investigated so far.

In view of these facts the basic goal of the present paper is precisely a systematic investigation of the relation between these two approaches to the classification of black hole solutions.

From the point of view of spherical symmetric solutions it was established that non-extremal black holes correspond to orbits of regular, diagonalizable Lax operators in the \mathbb{K}^* representation of \mathbb{H}^* , while extremal black holes are associated with nilpotent orbits in the same representation. For this reason a considerable amount of work was recently devoted to the study and classification of nilpotent orbits of the subgroup $\mathbb{H}^* \subset \mathbb{U}$ extending and completing work done by mathematicians for the nilpotent orbits of the full group \mathbb{U} . In the case of the so named S^3 -model, leading to $\mathbb{U}_{D=3} = \mathfrak{g}_{2,2}$, partial results were obtained in [35] confirmed and extended in [28].

In [29] the present authors, in collaboration with M. Trigiante, succeeded in elaborating a complete algorithm for the construction and classification of \mathbb{H}^* nilpotent orbits in the \mathbb{K}^* representation, applicable to all relevant supergravity models based on scalar manifolds that are symmetric coset spaces. The algorithm is based on the method of the standard triples $\{h, X, Y\}$, namely of the embedding of $\mathfrak{sl}(2, \mathbb{R})$ algebras where $X, Y \in \mathbb{K}^*$ and $h \in \mathbb{H}^*$, with the additional essential use of the Weyl group \mathcal{W} and of its subgroup \mathcal{W}_H which preserves the splitting $\mathbb{U}_{D=3} = \mathbb{H}^* \oplus \mathbb{K}^*$. The final labeling of the orbits is given by three set of eigenvalues named the α , β and γ labels (see [29], [34] for details).

In the same paper where it was introduced, the constructive orbit classification algorithm was applied to the $\mathfrak{g}_{2,2}$ case, reobtaining the 7 orbits found in previous calculations. It was also applied to the non maximally split case of the algebras $\mathfrak{so}(4, 4 + 2s)$ where, independently from the value of s , we were able to single out a fixed pattern of 37 nilpotent orbits. Indeed in [29] we showed that the structure of nilpotent orbits is a property of the Tits-Satake subalgebra of any algebra, giving a precise mathematical and physical relevance to the concept of *Tits Satake universality classes* introduced several years ago in [36].

In the present paper we address the systematic issue of organizing all supergravity models based on symmetric spaces into a finite list of *Tits Satake universality classes* (see tables 3,4) and by means of an in depth group-theoretical analysis we show that for all elements of a given class there are always enough and appropriate parameters in the adjoint representation of \mathbb{H}^* to rotate a generic element of \mathbb{K}^* into its subspace \mathbb{K}_{TS}^* obtained by means of the Tits Satake projection. We show that the same is true for the \mathbf{W} -representation of charges: in the adjoint representation of $\mathbb{U}_{D=4}$ there are always enough and appropriate parameters to rotate any element of \mathbf{W} into its Tits Satake projection \mathbf{W}_{TS} . Therefore we show constructively that both at the $D = 4$ and $D = 3$ level the classification of black hole solutions can be restricted to the classification of orbit structures for a finite list of maximally split Tits Satake algebras.

It is also appropriate to stress that the construction of nilpotent orbits for these universality classes has already consistently progressed, in view of the results of [29] and of [37] where the orbit pattern for the $\mathfrak{f}_{4,4}$ universality class was derived.

Having established this crucial point the comparison between the $D = 4$ and $D = 3$ approaches is reduced to investigate the relation between the corresponding \mathbf{W}_{TS} and \mathbb{K}_{TS}^* orbits. Although simplified to its essence, the problem remains, and in the spherical symmetric approach we obtain only some partial

answers. We can summarize the issue in the following question: *The electromagnetic charges that can be obtained from Lax operators belonging to the same H^* orbit do always fall in the same W^* orbit of $U_{D=4}$?* The answer is far from being positive. Indeed one immediately gets counterexamples by showing that Lax operators belonging to the same H^* orbit can yield both charges with vanishing and charges with non vanishing quartic invariant \mathfrak{J}_4 . However there is a caveat. Typically we are interested in asymptotically flat black holes so that we have to exclude a non vanishing Taub-NUT charge. The Taub-NUT charge is associated with the highest root of the algebra and the reduction of a dynamical system which kills the highest root corresponds to a consistent truncation. So we can consider the vanishing Taub-NUT locus in every H^* -orbit and it appears that all Lax operators belonging to such a locus produce charge vectors within the same W -orbit. Since the vanishing Taub-NUT locus is not a coset manifold rather an algebraic surface within a coset manifold, we were not able to provide a formal proof of this fact, yet we can put it in the form of a **conjecture** since no counter example has been found.

This is what we can say if we stick to spherically symmetric solutions, yet there are entirely new perspectives that open up if we take into account the brilliant strategy introduced in [38, 39, 40, 41, 42, 43] how to associate multicenter non spherically symmetric solutions to each H^* nilpotent orbit of \mathbb{K}^* .

The main catch of this strategy relies on the use of a *symmetric*, rather than *solvable gauge*, to represent the field equations of the three-dimensional σ -model (equivalent to supergravity for stationary solutions), that has the coset $U_{D=3}/H^*$ as target space. The next crucial ingredient in this menu is the use of the central element h of each standard triple $\{h, X, Y\}$ as a grading operator that determines a nilpotent subalgebra $\mathbb{N} \subset U_{D=3}$, spanned by its eigen-operators of positive grading. Denoting $\mathbb{N} \cap \mathbb{K}^*$ the intersection of such an algebra with the \mathbb{K}^* subspace and $\mathfrak{h}(x)$ a map $\mathcal{M}_3 \rightarrow \mathbb{N} \cap \mathbb{K}^*$ from the three-dimensional base space into such intersection, the symmetric coset representative is just $\mathcal{Y}(x) = \exp[\mathfrak{h}(x)]$ and obeys three-dimensional field equations forming a solvable system. The components of $\mathfrak{h}(x)$ associated with lower gradings are just harmonic functions, while those associated with higher gradings obey Laplace equation with a source term provided by a functional of the lower grading components. In this way one can construct non spherical symmetric solutions of the σ -model representing multicenter black holes and also, as we will show in the sequel, Kerr like solutions. A crucial technical problem that, in papers [38, 39, 40, 41, 42, 43] was only touched upon, or solved with ad hoc procedures for specific cases, concerns the final *oxidation procedure*. The correspondence between the fields of supergravity and the fields of the three-dimensional σ -model is precise and algorithmic in the solvable gauge realization of the coset representative, not in the symmetric gauge. Hence, in order to read off the supergravity solution one has to make such a gauge transformation which, at first sight, appears a matter of art and ingenuity, case by case. Fortunately this is not true, since the problem was already solved in complete generality in [26, 23, 24, 25], by means of a formula (see eq.(3.26) of the present paper) originally found in the context of spherical symmetric solutions, yet of much further reach. Indeed coupling eq.(3.26) with the strategy of papers [38, 39, 40, 41, 42, 43], the general construction of multicenter black hole solutions associated with nilpotent orbits becomes truly algorithmic and even implementable on a computer. We consider this one of the main results of the present paper.

We applied the algorithm to the case of the S^3 model, exploring the general form of multicenter solutions and we came to the following conclusion. The solutions associated with one nilpotent orbit have the property that, for each pole shared by all the harmonic functions springing from the corresponding nilpotent algebra, we have a black hole whose charges and other properties are those displayed by the spherical symmetric solution generated by the Lax operator in that orbit. However it may happen that a pole is located only in one subset of the harmonic building blocks, while other poles are located in different subsets. In that case the black holes springing from each pole have the charges and properties

of the spherical black hole pertaining to various smaller orbits associated with each subgroups. This mechanism unveils that larger orbit black holes can be viewed as composite objects built from the coalescence of smaller ones when their centers come together, namely when the corresponding poles overlap.

Our analysis also shows that the solutions associated with very large orbits, typically those for which the corresponding Lax operator has nilpotency degree larger than three, are generically singular not only in the spherical symmetric case but also in the multicenter case, since at each common pole of the involved harmonic functions we just retrieve the same conditions pertaining to the spherical symmetric solution. This explicit result is somewhat different from the claims made at various places of ref.s [39, 40, 38, 41, 42, 43] that multicenter solutions with generic positions of the poles can be regular for higher orbits. In any case we would like to stress that the solutions associate with higher orbits can be quite complex and deserve a special detailed study. Several surprises might still lie ahead.

In the course of our analysis we have found a number of other results that up to our knowledge were not known in the literature. We plan to summarize and high lite our findings in our conclusions, sect.11

The structure of our paper is displayed in the contents. We emphasize that it is organized in three parts. In the first part sect.s 2-3 we review the σ -model approach to stationary supergravity solutions, the construction of multicenter solutions attached to nilpotent orbits and we spell out the complete oxidation algorithm.

In the second part, sect.s 4-5 we apply the general scheme to the case of the S^3 model and we explore all the solutions attached to each of the classified nilpotent orbits.

Part three of the paper, sect.s 6-10 is devoted to organize all the available supergravity models based on symmetric scalar manifolds into Tits Satake universality classes, to show how within each class we can always reduce the analysis to the Tits Satake subalgebra both at the $D = 4$ and $D = 3$ level and finally to analyse the general structure of the \mathbb{H}^* subalgebra and the \mathbb{K}^* representation in the maximally split casse (Tits Satake projection).

2 The general framework

The addressed issue is that of stationary solutions of $D = 4$ ungauged supergravity models. For such field theories that include, the metric, several abelian gauge fields and several scalar fields, we have a general form of the bosonic lagrangian, which is the following one:

$$\begin{aligned} \mathcal{L}^{(4)} = & \sqrt{|\det g|} \left[\frac{R[g]}{2} - \frac{1}{4} \partial_\mu \phi^a \partial^\mu \phi^b h_{ab}(\phi) + \text{Im} \mathcal{N}_{\Lambda\Sigma} F_{\mu\nu}^\Lambda F^{\Sigma|\mu\nu} \right] \\ & + \frac{1}{2} \text{Re} \mathcal{N}_{\Lambda\Sigma} F_{\mu\nu}^\Lambda F_{\rho\sigma}^\Sigma \epsilon^{\mu\nu\rho\sigma}, \end{aligned} \quad (2.1)$$

where $F_{\mu\nu}^\Lambda \equiv (\partial_\mu A_\nu^\Lambda - \partial_\nu A_\mu^\Lambda)/2$. In eq.(2.1) ϕ^a denotes the whole set of n_s scalar fields parameterizing the scalar manifold $\mathcal{M}_{scalar}^{D=4}$ which, for supersymmetry $\mathcal{N} > 2$, is necessarily a coset manifold:

$$\mathcal{M}_{scalar}^{D=4} = \frac{\text{U}_{D=4}}{\text{H}_c} \quad (2.2)$$

For $\mathcal{N} = 2$ eq.(2.2) is not obligatory but it is possible: a well determined class of symmetric homogeneous manifolds that are special Kähler manifolds [14] falls into the set up of the present general discussion.

The theory includes also n_v vector fields A_μ^Λ for which

$$\mathcal{F}_{\mu\nu}^{\pm|\Lambda} \equiv \frac{1}{2} \left[F_{\mu\nu}^\Lambda \mp i \frac{\sqrt{|\det g|}}{2} \epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma} \right] \quad (2.3)$$

denote the self-dual (respectively antiself-dual) parts of the field-strengths. As displayed in eq.(2.1) they are non minimally coupled to the scalars via the symmetric complex matrix

$$\mathcal{N}_{\Lambda\Sigma}(\phi) = i \text{Im}\mathcal{N}_{\Lambda\Sigma} + \text{Re}\mathcal{N}_{\Lambda\Sigma} \quad (2.4)$$

which transforms projectively under $U_{D=4}$. Indeed the field strengths $\mathcal{F}_{\mu\nu}^{\pm|\Lambda}$ plus their magnetic duals:

$$G_{\Lambda|\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu}{}^{\rho\sigma} \frac{\delta \mathcal{L}^{(4)}}{\delta F_{\rho\sigma}^\Lambda} \quad (2.5)$$

fill up a $2n_v$ -dimensional symplectic representation of $U_{D=4}$ which we call by the name of \mathbf{W} .

We rephrase the above statements by asserting that there is always a symplectic embedding of the duality group $U_{D=4}$,

$$U_{D=4} \mapsto \text{Sp}(2n_v, \mathbb{R}) \quad ; \quad n_v \equiv \# \text{ of vector fields} \quad (2.6)$$

so that for each element $\xi \in U_{D=4}$ we have its representation by means of a suitable real symplectic matrix:

$$\xi \mapsto \Lambda_\xi \equiv \begin{pmatrix} A_\xi & B_\xi \\ C_\xi & D_\xi \end{pmatrix} \quad (2.7)$$

satisfying the defining relation:

$$\Lambda_\xi^T \begin{pmatrix} \mathbf{0}_{n \times n} & \mathbf{1}_{n \times n} \\ -\mathbf{1}_{n \times n} & \mathbf{0}_{n \times n} \end{pmatrix} \Lambda_\xi = \begin{pmatrix} \mathbf{0}_{n \times n} & \mathbf{1}_{n \times n} \\ -\mathbf{1}_{n \times n} & \mathbf{0}_{n \times n} \end{pmatrix} \quad (2.8)$$

Under an element of the duality group the field strengths transform as follows:

$$\begin{pmatrix} \mathcal{F}^+ \\ \mathcal{G}^+ \end{pmatrix}' = \begin{pmatrix} A_\xi & B_\xi \\ C_\xi & D_\xi \end{pmatrix} \begin{pmatrix} \mathcal{F}^+ \\ \mathcal{G}^+ \end{pmatrix} \quad ; \quad \begin{pmatrix} \mathcal{F}^- \\ \mathcal{G}^- \end{pmatrix}' = \begin{pmatrix} A_\xi & B_\xi \\ C_\xi & D_\xi \end{pmatrix} \begin{pmatrix} \mathcal{F}^- \\ \mathcal{G}^- \end{pmatrix} \quad (2.9)$$

where, by their own definitions:

$$\mathcal{G}^+ = \mathcal{N} \mathcal{F}^+ \quad ; \quad \mathcal{G}^- = \overline{\mathcal{N}} \mathcal{F}^- \quad (2.10)$$

and the complex symmetric matrix \mathcal{N} should transform as follows:

$$\mathcal{N}' = (C_\xi + D_\xi \mathcal{N}) (A_\xi + B_\xi \mathcal{N})^{-1} \quad (2.11)$$

Choose a parametrization of the coset $\mathbb{L}(\phi) \in U_{D=4}$, which assigns a definite group element to every coset point identified by the scalar fields. Through the symplectic embedding (2.7) this produces a definite ϕ -dependent symplectic matrix

$$\begin{pmatrix} A(\phi) & B(\phi) \\ C(\phi) & D(\phi) \end{pmatrix} \quad (2.12)$$

in the W -representation of $U_{D=4}$. In terms of its blocks the kinetic matrix $\mathcal{N}(\phi)$ is explicitly given by the Gaillard-Zumino formula:

$$\mathcal{N}(\phi) = [C(\phi) - iD(\phi)] [A(\phi) - iB(\phi)]^{-1} , \quad (2.13)$$

2.1 The σ -model approach to extremal black holes

A very powerful token in deriving stationary and in particular extremal black hole solutions of $D = 4$ supergravity that depend only on three space-like coordinates x_1, x_2, x_3 , is provided by the time-reduction of the four-dimensional field equations to those of an effective three-dimensional σ -model. Let us shortly review this procedure.

In all $\mathcal{N} = 2$ cases the number of vector fields in the theory is $n_v = n + 1$ where n is the complex dimension of the scalar manifold ($n_s = 2n$), while in the case of other theories the relation between n_v and n_s is different. Notwithstanding this difference, we can always introduce a $2n_v \times 2n_v$ field dependent matrix \mathcal{M}_4 defined as follows:

$$\mathcal{M}_4 = \left(\begin{array}{c|c} \text{Im}\mathcal{N}^{-1} & \text{Im}\mathcal{N}^{-1} \text{Re}\mathcal{N} \\ \hline \text{Re}\mathcal{N} \text{Im}\mathcal{N}^{-1} & \text{Im}\mathcal{N} + \text{Re}\mathcal{N} \text{Im}\mathcal{N}^{-1} \text{Re}\mathcal{N} \end{array} \right) \quad (2.14)$$

$$\mathcal{M}_4^{-1} = \left(\begin{array}{c|c} \text{Im}\mathcal{N} + \text{Re}\mathcal{N} \text{Im}\mathcal{N}^{-1} \text{Re}\mathcal{N} & -\text{Re}\mathcal{N} \text{Im}\mathcal{N}^{-1} \\ \hline -\text{Im}\mathcal{N}^{-1} \text{Re}\mathcal{N} & \text{Im}\mathcal{N}^{-1} \end{array} \right) \quad (2.15)$$

and we can introduce the following set of $2 + n_s + 2n_v$ fields depending on the three parameters x_i :

	Generic		$\mathcal{N} = 2$
warp factor	$U(x)$	1	1
Taub-NUT field	$a(x)$	1	1
D=4 scalars	$\phi^a(x)$	n_s	$2n$
Scalars from vectors	$Z^M(x) = (Z^\Lambda(x), Z_\Sigma(x))$	$2n_v$	$2n + 2$
Total	$2 + n_s + 2n_v$		$4n + 4$

the fields $\{U, a, \phi, Z\}$ are interpreted as the coordinates of a new $(2 + n_s + 2n_v)$ -dimensional manifold \mathcal{Q} , whose metric we declare to be the following:

$$ds_{\mathcal{Q}}^2 = \frac{1}{4} [dU^2 + h_{rs} d\phi^r d\phi^s + e^{-2U} (da + \mathbf{Z}^T \mathbb{C} d\mathbf{Z})^2 + 2e^{-U} d\mathbf{Z}^T \mathcal{M}_4 d\mathbf{Z}] \quad (2.16)$$

having denoted by \mathbb{C} the constant symplectic invariant metric in $2n_v$ dimensions that underlies the construction of the matrix $\mathcal{N}_{\Lambda\Sigma}$. The metric (2.16) has the following indefinite signature

$$\text{sign} [ds_{\mathcal{Q}}^2] = \left(\underbrace{+, \dots, +}_{2+n_s}, \underbrace{-, \dots, -}_{2n_v+2} \right) \quad (2.17)$$

since the matrix \mathcal{M}_4 is negative definite.

Moreover one very important point to be stressed is that the metric (2.16) admits a typically large group of isometries. Certainly it admits all the isometries of the original scalar manifold $\mathcal{M}_{\text{scalar}}$ enlarged with additional ones related to the new fields that have been introduced $\{U, a, Z^M\}$. In the case when the $D = 4$ scalar manifold is a homogeneous symmetric space:

$$\mathcal{M}_{\text{scalar}} = \frac{U_{D=4}}{H_c} \quad (2.18)$$

one can show [32, 33, 22, 44], that the manifold \mathcal{Q} with the metric (2.16) is a new homogeneous symmetric space

$$\mathcal{Q} = \frac{\mathbb{U}_{D=3}}{\mathbb{H}^\star} \quad (2.19)$$

whose structure is universal and can be described in general terms.

2.1.1 General structure of the $\mathbb{U}_{D=3}$ Lie algebra

The Lie algebra $\mathbb{U}_{D=3}$ of the numerator group always contains, as subalgebra, the duality algebra $\mathbb{U}_{D=4}$ of the parent supergravity theory in $D = 4$ and a universal $\mathfrak{sl}(2, \mathbb{R})_E$ algebra which is associated with the gravitational degrees of freedom $\{U, a\}$. Furthermore, with respect to this subalgebra, $\mathbb{U}_{D=3}$ admits the following universal decomposition, holding for all \mathcal{N} -extended supergravities:

$$\text{adj}(\mathbb{U}_{D=3}) = \text{adj}(\mathbb{U}_{D=4}) \oplus \text{adj}(\mathfrak{sl}(2, \mathbb{R})_E) \oplus W_{(2, \mathbf{W})} \quad (2.20)$$

where \mathbf{W} is the **symplectic** representation of $\mathbb{U}_{D=4}$ to which the electric and magnetic field strengths are assigned. Indeed the scalar fields associated with the generators of $W_{(2, \mathbf{W})}$ are just those coming from the vectors in $D = 4$. Denoting the generators of $\mathbb{U}_{D=4}$ by T^a , the generators of $\mathfrak{sl}(2, \mathbb{R})_E$ by L^x and denoting by W^{iM} the generators in $W_{(2, \mathbf{W})}$, the commutation relations that correspond to the decomposition (2.20) have the following general form:

$$\begin{aligned} [T^a, T^b] &= f^{ab}_c T^c \\ [L^x, L^y] &= f^{xy}_z L^z, \\ [T^a, W^{iM}] &= (\Lambda^a)^M_N W^{iN}, \\ [L^x, W^{iM}] &= (\lambda^x)^i_j W^{jM}, \\ [W^{iM}, W^{jN}] &= \epsilon^{ij} (K_a)^{MN} T^a + \mathbb{C}^{MN} k_x^{ij} L^x \end{aligned} \quad (2.21)$$

where the 2×2 matrices $(\lambda^x)^i_j$, are the canonical generators of $\mathfrak{sl}(2, \mathbb{R})$ in the fundamental, defining representation:

$$\lambda^3 = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix} \quad ; \quad \lambda^1 = \begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix} \quad ; \quad \lambda^2 = \begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix} \quad (2.22)$$

while Λ^a are the generators of $\mathbb{U}_{D=4}$ in the symplectic representation \mathbf{W} . By

$$\mathbb{C}^{MN} \equiv \left(\begin{array}{c|c} \mathbf{0}_{n_v \times n_v} & \mathbf{1}_{n_v \times n_v} \\ \hline -\mathbf{1}_{n_v \times n_v} & \mathbf{0}_{n_v \times n_v} \end{array} \right) \quad (2.23)$$

we denote the antisymmetric symplectic metric in $2n_v$ dimensions, n_v being the number of vector fields in $D = 4$ as we have already stressed. The symplectic character of the representation \mathbf{W} is asserted by the identity:

$$\Lambda^a \mathbb{C} + \mathbb{C} (\Lambda^a)^T = 0 \quad (2.24)$$

The fundamental doublet representation of $\mathfrak{sl}(2, \mathbb{R})$ is also symplectic and we have denoted by $\epsilon^{ij} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ the 2-dimensional symplectic metric, so that:

$$\lambda^x \epsilon + \epsilon (\lambda^x)^T = 0, \quad (2.25)$$

In eq. (2.21) we have used the standard convention according to which symplectic indices are raised and lowered with the appropriate symplectic metric, while adjoint representation indices are raised and lowered with the Cartan-Killing metric.

2.1.2 General form of the three-dimensional σ -model

Next we consider a gravity coupled three-dimensional euclidian σ -model, whose fields

$$\Phi^A(x) \equiv \{U(x), a(x), \phi(x), Z(x)\}$$

describe mappings:

$$\Phi : \mathcal{M}_3 \rightarrow \mathcal{Q} \quad (2.26)$$

from a three-dimensional manifold \mathcal{M}_3 , whose metric we denote by $\gamma_{ij}(x)$, to the target space \mathcal{Q} . The action of this σ -model is the following:

$$\mathcal{A}^{[3]} = \int \sqrt{\det \gamma} \mathfrak{R}[\gamma] d^3x + \int \sqrt{\det \gamma} \mathcal{L}^{(3)} d^3x \quad (2.27)$$

$$\begin{aligned} \mathcal{L}^{(3)} = & (\partial_i U \partial_j U + h_{rs} \partial_i \phi^r \partial_j \phi^s \\ & + e^{-2U} (\partial_i a + \mathbf{Z}^T \mathbb{C} \partial_i \mathbf{Z}) (\partial_j a + \mathbf{Z}^T \mathbb{C} \partial_j \mathbf{Z}) + 2 e^{-U} \partial_i \mathbf{Z}^T \mathcal{M}_4 \partial_j \mathbf{Z}) \gamma^{ij} \end{aligned} \quad (2.28)$$

where $\mathfrak{R}[\gamma]$ denotes the scalar curvature of the metric γ_{ij} .

The field equations of the σ -model are obtained by varying the action both in the metric γ_{ij} and in the fields $\Phi^A(x)$. The Einstein equation reads as usual:

$$\mathfrak{R}_{ij} - \frac{1}{2} \gamma_{ij} \mathfrak{R} = \mathfrak{T}_{ij} \quad (2.29)$$

where:

$$\mathfrak{T}_{ij} = \frac{\delta \mathcal{L}^{(3)}}{\delta \gamma^{ij}} - \gamma_{ij} \mathcal{L}^{(3)} \quad (2.30)$$

is the stress energy tensor, while the matter field equations assume the standard form:

$$\frac{1}{\sqrt{\det \gamma}} \gamma^{ij} \partial_i \left[\sqrt{\det \gamma} \frac{\delta \mathcal{L}^{(3)}}{\delta \partial^j \Phi^A} \right] - \frac{\delta \mathcal{L}^{(3)}}{\delta \Phi^A} = 0 \quad (2.31)$$

As it is well known, in $D = 3$ there is no propagating graviton and the Riemann tensor is completely determined by the Ricci tensor, namely, via Einstein equations, by the stress-energy tensor of the matter fields.

Extremal solutions of the σ -model are those for which the three-dimensional metric can be consistently chosen flat:

$$\gamma_{ij} = \delta_{ij} \quad (2.32)$$

corresponding to a vanishing stress-energy tensor:

$$(\partial_i U \partial_j U + h_{rs} \partial_i \phi^r \partial_j \phi^s + e^{-2U} (\partial_i a + \mathbf{Z}^T \mathbb{C} \partial_i \mathbf{Z}) (\partial_j a + \mathbf{Z}^T \mathbb{C} \partial_j \mathbf{Z}) + 2 e^{-U} \partial_i \mathbf{Z}^T \mathcal{M}_4 \partial_j \mathbf{Z}) = 0 \quad (2.33)$$

We will see in the sequel how the nilpotent orbits of the group H^* in the \mathbb{K}^* representation can be systematically associated with general extremal solutions of the field equations.

2.2 Oxidation rules for extremal multicenter black holes

Let us now describe the oxidation rules, namely the procedure by means of which to every configuration of the three-dimensional fields $\Phi(x) = \{U(x), a(x), \phi(x), Z(x)\}$, satisfying the field equations (2.31) and also the extremality condition (2.33), we can associate a well defined configuration of the four-dimensional fields satisfying the field equations of supergravity that follow from the lagrangian (2.1). We might write such oxidation rules for general solutions of the σ -model, also non extremal, yet given the goal of the present paper we confine ourselves to spell out such rule in the extremal case, which is somewhat simpler since it avoids the extra complications related with the three-dimensional metric γ_{ij} .

In order to write the $D = 4$ fields, the first necessary item we have to determine is the Kaluza-Klein vector field $\mathbf{A}^{[KK]} = A_i^{[KK]} dx^i$. This latter is worked out through the following dualization procedure:

$$\begin{aligned} \mathbf{F}^{[KK]} &= d\mathbf{A}^{[KK]} \\ \mathbf{F}^{[KK]} &= -\epsilon_{ijk} dx^i \wedge dx^j \left[\exp[-2U] \left(\partial^k a + Z \mathbb{C} \partial^k Z \right) \right] \end{aligned} \quad (2.34)$$

Given the Kaluza-Klein vector we can write the four-dimensional metric which is the following:

$$ds^2 = -\exp[U] \left(dt + \mathbf{A}^{[KK]} \right)^2 + \exp[-U] dx^i \otimes dx^j \delta_{ij} \quad (2.35)$$

The vielbein description of the same metric is immediate. We just write:

$$\begin{aligned} ds^2 &= -E^0 \otimes E^0 + E^i \otimes E^i \\ E^0 &= \exp\left[\frac{U}{2}\right] \left(dt + \mathbf{A}^{[KK]} \right) \\ E^i &= \exp\left[-\frac{U}{2}\right] dx^i \end{aligned} \quad (2.36)$$

Next we can present the form of the electromagnetic field strengths:

$$\begin{aligned} \mathbf{F}^\Lambda &= \mathbb{C}^{\Lambda M} \partial_i Z_M dx^i \wedge (dt + \mathbf{A}^{[KK]}) \\ &+ \epsilon_{ijk} dx^i \wedge dx^j \left[\exp[-U] (\text{Im} \mathcal{N}^{-1})^{\Lambda\Sigma} \left(\partial^k Z_\Sigma + \text{Re} \mathcal{N}_{\Sigma\Gamma} \partial^k Z^\Gamma \right) \right] \end{aligned} \quad (2.37)$$

Next we define the electromagnetic charges and the Taub-NUT charges for multicenter solutions. Considering the metric (2.35) the black hole centers are defined by the zeros of the warp-factor $\exp[U(\vec{x})]$. In a composite m -black hole solution there are m three-vectors \vec{r}_α ($\alpha = 1, \dots, m$), such that:

$$\lim_{\vec{x} \rightarrow \vec{r}_\alpha} \exp[U(\vec{x})] = 0 \quad (2.38)$$

Each of these zeros defines a non trivial homology two-cycle \mathbb{S}_α^2 of the 4-dimensional space-time which surrounds the singularity \vec{r}_α . The electromagnetic charges of the individual holes are obtained by integrating the field strengths and their duals on such homology cycles.

$$\left(\begin{array}{c} p^\Lambda \\ q_\Sigma \end{array} \right)_\alpha = \frac{1}{4\pi\sqrt{2}} \left(\begin{array}{c} \int_{\mathbb{S}_\alpha^2} \mathbf{F}^\Lambda \\ \int_{\mathbb{S}_\alpha^2} \mathbf{G}_\Sigma \end{array} \right) \equiv \frac{1}{4\pi} \int_{\mathbb{S}_\alpha^2} j^{EM} \quad (2.39)$$

Utilizing the form of the field strengths we obtain the explicit formula:

$$\mathcal{Q}_\alpha \equiv \left(\begin{array}{c} p^\Lambda \\ q_\Sigma \end{array} \right)_\alpha = \frac{1}{4\pi\sqrt{2}} \int_{\mathbb{S}_\alpha^2} \epsilon_{ijk} dx^i \wedge dx^j \left[\exp[-U] \mathcal{M}_4 \partial^k Z + \exp[-2U] \left(\partial^k a + Z \mathbb{C} \partial^k Z \right) \mathbb{C} Z \right] \quad (2.40)$$

which provides m -sets of electromagnetic charges associated with the solution. Similarly we have m Taub-NUT charges defined by:

$$\mathbf{n}_\alpha = -\frac{1}{4\pi} \int_{\mathbb{S}_\alpha^2} \epsilon_{ijk} dx^i \wedge dx^j \exp[-2U] \left(\partial^k a + Z \mathbb{C} \partial^k Z \right) \equiv \frac{1}{4\pi} \int_{\mathbb{S}_\alpha^2} j^{TN} \quad (2.41)$$

2.2.1 Reduction to the spherical case

The spherical symmetric one-center solutions are retrieved from the general case by assuming that all the three-dimensional fields depend only on one radial coordinate:

$$\tau = -\frac{1}{r} \quad ; \quad r = \sqrt{x_1^2 + x_2^2 + x_3^2} \quad (2.42)$$

On functions only of τ we have the identity:

$$\partial_i f(\tau) = -x^i \tau^3 \frac{d}{d\tau} f(\tau) \quad (2.43)$$

and introducing polar coordinates:

$$\begin{aligned} x_1 &= \frac{1}{\tau} \cos \theta \\ x_2 &= \frac{1}{\tau} \sin \theta \sin \varphi \\ x_3 &= \frac{1}{\tau} \sin \theta \cos \varphi \end{aligned} \quad (2.44)$$

we obtain:

$$\tau^3 \epsilon_{ijk} x^i dx^j \wedge dx^k = -2 \sin \theta d\theta \wedge d\varphi \quad (2.45)$$

By using these identities and restricting one's attention to the extremal case, the action of the σ -model (2.27) reduces to:

$$\begin{aligned} \mathcal{A} &= \int d\tau \mathcal{L} \\ \mathcal{L} &= \dot{U}^2 + h_{rs} \dot{\varphi}^r \dot{\varphi}^s + e^{-2U} (\dot{a} + \mathbf{Z}^T \mathbb{C} \dot{\mathbf{Z}})^2 + 2e^{-U} \dot{\mathbf{Z}}^T \mathcal{M}_4 \dot{\mathbf{Z}} \end{aligned} \quad (2.46)$$

where the dot denotes derivatives with respect to the τ variable. The σ -model field equations take the standard form of the Euler Lagrangian equations:

$$\frac{d}{d\tau} \frac{d\mathcal{L}}{d\dot{\Phi}} = \frac{d\mathcal{L}}{d\Phi} \quad (2.47)$$

and the extremality conditions (2.33) reduces to:

$$\mathcal{L} = \dot{U}^2 + h_{rs} \dot{\varphi}^r \dot{\varphi}^s + e^{-2U} (\dot{a} + \mathbf{Z}^T \mathbb{C} \dot{\mathbf{Z}})^2 + 2e^{-U} \dot{\mathbf{Z}}^T \mathcal{M}_4 \dot{\mathbf{Z}} = 0 \quad (2.48)$$

It appears from this that spherical extremal black holes are in one-to-one correspondence with light-like geodesics of the manifold \mathcal{Q} .

The reduced oxidation rules In the spherical case the above discussed oxidation rules reduce as follows. For the metric we have

$$ds_{(4)}^2 = -e^{U(\tau)} (dt + 2\mathbf{n} \cos \theta d\varphi)^2 + e^{-U(\tau)} \left[\frac{1}{\tau^4} d\tau^2 + \frac{1}{\tau^2} (d\theta^2 + \sin^2 \theta d\phi^2) \right] \quad (2.49)$$

where \mathbf{n} denotes the Taub-NUT charge obtained from the form of the Kaluza-Klein field strength:

$$\begin{aligned} \mathbf{F}^{KK} &= -2\mathbf{n} \sin \theta d\theta \wedge d\varphi \\ \mathbf{n} &= \left(\dot{a} + Z \mathbb{C} \dot{Z} \right) \end{aligned} \quad (2.50)$$

The electromagnetic field-strengths are instead the following ones:

$$F^\Lambda = 2p^\Lambda \sin \theta d\theta \wedge d\varphi + \dot{Z}_\Lambda d\tau \wedge (dt + 2\mathbf{n} \cos \theta d\varphi) \quad (2.51)$$

where the magnetic charges p^Λ are extracted from the reduction of the general formula (2.40), namely:

$$\mathcal{Q}^M = \begin{pmatrix} p^\Lambda \\ q_\Sigma \end{pmatrix} = \sqrt{2} \left[e^{-U} \mathcal{M}_4 \dot{Z} - \mathbf{n} \mathbb{C} Z \right]^M \quad (2.52)$$

2.3 A counter example: the extremal Kerr metric

In this section, in order to better clarify the notion of extremality provided by conditions (2.32-2.33) we consider the physically relevant counter-example of the extremal Kerr metric. Such static solution of Einstein equations is certainly encoded in the σ -model approach yet it is not extremal in the sense of eq.s (2.32-2.33) and therefore it is not related to any nilpotent orbit. Indeed the extremal Kerr metric is a solution of pure gravity and as such its σ -model representation lies in the euclidian submanifold:

$$\frac{\text{SL}(2, \mathbb{R})}{\text{O}(2)} \quad (2.53)$$

for which the coset tangent space \mathbb{K} contains no nilpotent elements.

Instead the so named BPS Kerr-Newman metric, which is not extremal in the sense of General Relativity and actually displays a naked singularity, is extremal in the sense of eq.s (2.32-2.33) and can be retrieved in one of the nilpotent orbits of the S^3 -model. We will show that explicitly in section 5.4.

As a preparation to such discussions let us recall the general form of the Kerr-Newmann metric which we represent in polar coordinates as it follows:

$$ds_{KN}^2 = -V^0 \otimes V^0 + \sum_{i=1}^3 V^i \otimes V^i \quad (2.54)$$

$$V^0 = \frac{\delta(r)}{\sigma(r, \theta)} (dt - \alpha \sin^2 \theta d\phi) \quad (2.55)$$

$$V^1 = \frac{\sigma(r, \theta)}{\delta(r)} dr \quad (2.56)$$

$$V^2 = \sigma(r, \theta) d\theta \quad (2.57)$$

$$V^3 = \frac{\sin(\theta)}{\sigma(r, \theta)} ((r^2 + \alpha^2) d\phi - \alpha dt) \quad (2.58)$$

$$\delta(r) = \sqrt{q^2 + r^2 + \alpha^2 - 2mr} \quad (2.59)$$

$$\sigma(r, \theta) = \sqrt{r^2 + \alpha^2 \cos^2(\theta)} \quad (2.60)$$

Parameters of the Kerr-Newman solution are the mass m , the electric charge q and the angular momentum $J = m\alpha$ of the Black Hole. The two particular cases we shall consider in this paper correspond to:

- a) The extremal Kerr solution: $q = 0$ and $m = \alpha$.
- b) The BPS Kerr-Newman solution $q = m$, arbitrary α .

Let us then focus now on the extremal Kerr solution. With the choice $m = \alpha$, $q = 0$, the metric (2.54) can be rewritten in the following form:

$$ds_{EK}^2 = -\exp[U] \left(dt + \mathbf{A}^{[KK]} \right)^2 + \exp[-U] \gamma_{ij} dy^i \otimes dy^j \quad (2.61)$$

where $y^i = \{r, \theta, \phi\}$ are the polar coordinates, the three dimensional metric γ_{ij} is the following one:

$$\gamma_{ij} = \begin{pmatrix} \frac{2r^2 - \alpha^2 + \alpha^2 \cos(2\theta)}{2r^2} & 0 & 0 \\ 0 & r^2 - \frac{\alpha^2}{2} + \frac{1}{2}\alpha^2 \cos(2\theta) & 0 \\ 0 & 0 & r^2 \sin^2(\theta) \end{pmatrix} \quad (2.62)$$

the warp factor is:

$$U = \log \left[\frac{r^2 - \alpha^2 \sin^2(\theta)}{(r + \alpha)^2 + \alpha^2 \cos^2(\theta)} \right] \quad (2.63)$$

and the Kaluza Klein vector has the following appearance:

$$\mathbf{A}^{[KK]} = \frac{2\alpha^2(r + \alpha) \sin^2(\theta)}{r^2 - \alpha^2 \sin^2(\theta)} d\phi \quad (2.64)$$

In presence of the metric γ_{ij} the duality relation between the Kaluza Klein vector field and the σ -model scalar field a reads as follows:

$$\mathbf{F}_{ij}^{[KK]} \equiv \partial_{[i} \mathbf{A}_{j]}^{[KK]} = \exp[-2U] \sqrt{\det \gamma} \epsilon_{ijk} \gamma^{k\ell} \partial_\ell a \quad (2.65)$$

and it is solved by:

$$a = -\frac{2\alpha^2 \cos(\theta)}{2r^2 + 4\alpha r + 3\alpha^2 + \alpha^2 \cos(2\theta)} \quad (2.66)$$

In this way, by means of inverse engineering we have showed how the extremal Kerr metric is retrieved in the σ -model approach. The crucial point is that the metric γ_{ij} is not flat and hence such a configuration of the U, a fields does not correspond to an extremal solution of the σ -model field equations. Indeed calculating the curvature two-form of the three-dimensional metric (2.62) we find

$$\mathfrak{R}^{12} = \frac{4\alpha^2 (2r^2 + \alpha^2 - \alpha^2 \cos(2\theta))}{(2r^2 - \alpha^2 + \alpha^2 \cos(2\theta))^3} e^1 \wedge e^2 \quad (2.67)$$

$$\mathfrak{R}^{13} = \frac{4\alpha^2}{(2r^2 - \alpha^2 + \alpha^2 \cos(2\theta))^2} e^1 \wedge e^3 \quad (2.68)$$

$$\mathfrak{R}^{23} = -\frac{4\alpha^2}{(2r^2 - \alpha^2 + \alpha^2 \cos(2\theta))^2} e^2 \wedge e^3 \quad (2.69)$$

where

$$e^1 = \frac{dr \sqrt{\frac{\cos(2\theta)\alpha^2}{r^2} - \frac{\alpha^2}{r^2} + 2}}{\sqrt{2}} \quad (2.70)$$

$$e^2 = d\theta \sqrt{r^2 - \frac{\alpha^2}{2} + \frac{1}{2}\alpha^2 \cos(2\theta)} \quad (2.71)$$

$$e^3 = d\phi r \sin(\theta) \quad (2.72)$$

is the *dreibein* corresponding to (2.62).

Hopefully this explicit calculation should have convinced the reader that the extremal Kerr solution and, by the same token, also the extremal Kerr-Newman solution are not extremal in the σ -model sense and are retrieved in regular rather than in nilpotent orbits of U/H^* .

3 Construction of multicenter solutions associated with nilpotent orbits

For spherically symmetric black holes the construction of solutions is associated with nilpotent orbits in the following way. A representative of the H^* orbit is a standard triple $\{h, X, Y\}$ and hence an embedding of an $\mathfrak{sl}(2, \mathbb{R})$ Lie algebra:

$$[h, X] = 2X \quad ; \quad [h, Y] = -2Y \quad ; \quad [X, Y] = 2h \quad (3.1)$$

into $U_{D=3}$ in such a way that $h \in \mathbb{H}^*$ and $X, Y \in \mathbb{K}^*$. The nilpotent operator X is identified with the Lax operator L_0 at euclidian time $\tau = 0$ and the corresponding solution depending on τ is constructed by using the algorithm described in [23, 25, 28]. In the multicenter approach of [38, 39, 40, 41, 42, 43] one utilizes the standard triple to single out a nilpotent subalgebra \mathbb{N} , as follows. One diagonalizes the adjoint action of the central element h of the triple on the Lie Algebra $U_{D=3}$:

$$[h, C_\mu] = \mu C_\mu \quad (3.2)$$

The set of all eigen-operators C_μ corresponding to positive gradings $\mu > 0$ spans a subalgebra $\mathbb{N} \subset U_{D=3}$ which is necessarily nilpotent

$$\mathbb{N} = \text{span}[C_2, C_3, \dots, C_{max}] \quad (3.3)$$

Such a nilpotent subalgebra has an intersection $\mathbb{N} \cap \mathbb{K}^*$ with the space \mathbb{K}^* which is not empty since at least the operator $C_2 = X$ is present by definition of a standard triple. The next steps of the construction are as follows.

3.1 The coset representative in the symmetric gauge

Given a basis A^i of the space $\mathbb{N}_{\mathbb{K}} \equiv \mathbb{N} \cap \mathbb{K}^*$, whose dimension we denote:

$$\ell \equiv \dim \mathbb{N}_{\mathbb{K}} \quad (3.4)$$

and a basis B^α of the subalgebra $\mathbb{N}_{\mathbb{H}} \equiv \mathbb{N} \cap \mathbb{H}^*$, whose dimension we denote

$$\mathfrak{m} \equiv \dim \mathbb{N}_{\mathbb{H}} \quad (3.5)$$

we can construct a map:

$$\mathfrak{H} : \mathbb{R}^3 \rightarrow \mathbb{N}_{\mathbb{K}} \quad (3.6)$$

by writing:

$$\mathbb{N}_{\mathbb{K}} \ni \mathfrak{H}(\vec{x}) = \sum_{i=1}^{\ell} \mathfrak{h}_i(\vec{x}) A^i \quad (3.7)$$

By construction, the point dependent Lie algebra element $\mathfrak{H}(\vec{x})$ is nilpotent of a certain maximal degree d_n , so that its exponential map to the nilpotent group $\mathbb{N} \subset \mathbb{U}_{D=3}$ truncates to a finite sum:

$$\mathcal{Y}(x) = \exp[\mathfrak{H}(\vec{x})] = \mathbf{1} + \sum_{a=1}^{d_n} \frac{1}{a!} \mathfrak{H}^a(\vec{x}) \quad (3.8)$$

The above constructed object realizes an explicit \vec{x} -dependent coset representative from which we can construct the Maurer Cartan left-invariant one form:

$$\Sigma = \mathcal{Y}^{-1} \partial_i \mathcal{Y} dx^i \quad (3.9)$$

Next let us decompose Σ along the \mathbb{K}^* subspace and the \mathbb{H}^* subalgebra, respectively. This is done by setting:

$$\mathbf{P} = \text{Tr}(\Sigma K^A) K_A \quad ; \quad \Omega = \text{Tr}(\Sigma H^m) H_m \quad (3.10)$$

where K_A and H_m denote a basis of generators for the two considered subspaces, K^A and H^m being their duals:

$$\text{Tr}(K^A K_B) = \delta_B^A \quad ; \quad \text{Tr}(H^m H_n) = \delta_n^m \quad ; \quad \text{Tr}(K^A H_n) = 0 \quad (3.11)$$

Denoting:

$$\star \mathbf{P} \equiv \frac{1}{2} \epsilon_{ijk} \delta^{im} \mathbf{P}_m dx^j \wedge dx^k \quad (3.12)$$

the Hodge-dual of the coset vielbein

$$\mathbf{P} = \mathbf{P}_m dx^m \quad (3.13)$$

the field equations of the three dimensional σ -model reduce to the following one:

$$d\star \mathbf{P} = \Omega \wedge \star \mathbf{P} - \star \mathbf{P} \wedge \Omega \quad (3.14)$$

Actually, since $\mathbb{N} \subset \mathbb{U}_{D=3}$ forms a nilpotent subalgebra the constructed object \mathcal{Y} realizes a map from the three-dimensional space to the much smaller coset manifold:

$$\mathcal{Y} : \mathbb{R}^3 \rightarrow \frac{\mathbb{N}}{\mathbb{N}_{\mathbb{H}}} \quad (3.15)$$

and due to the polynomial form of the coset representative the final equations of motion obtain a triangular solvable form that we describe here below. Since the algebra \mathbb{N} is nilpotent, its derivative series terminates, namely we have:

$$\mathbb{N} \supset \mathcal{D}\mathbb{N} \supset \dots \supset \mathcal{D}^n \mathbb{N} \supset \mathcal{D}^{n+1} \mathbb{N} = \mathbf{0} \quad (3.16)$$

where at each step $\mathcal{D}^i \mathbb{N}$ is a proper subspace of $\mathcal{D}^{i-1} \mathbb{N}$. Correspondingly let us define:

$$\mathcal{D}^i \mathbb{N}_{\mathbb{K}} = \mathcal{D}^i \mathbb{N} \bigcap \mathbb{K}^* \quad (3.17)$$

the intersections of the derivative subalgebras with the \mathbb{K}^\star subspace and let us introduce the complementary orthogonal subspaces:

$$\mathcal{D}^i \mathbb{N}_{\mathbb{K}} = \mathbb{N}_{\mathbb{K}}^{(i)} \oplus \mathcal{D}^{i+1} \mathbb{N}_{\mathbb{K}} \quad (3.18)$$

This yields an orthogonal graded decomposition of the space $\mathbb{N}_{\mathbb{K}}$ of the following form:

$$\mathbb{N}_{\mathbb{K}} = \bigoplus_{a=0}^n \mathbb{N}_{\mathbb{K}}^{(a)} \quad (3.19)$$

The space $\mathbb{N}_{\mathbb{K}}^{(0)}$ contains those generators that cannot be produced by any commutator within the algebra, $\mathbb{N}_{\mathbb{K}}^{(1)}$ contains those generators that are produced in simple commutators, $\mathbb{N}_{\mathbb{K}}^{(2)}$ contains those that are produced in double commutators and so on. Let us name

$$\ell_a = \dim \mathbb{N}_{\mathbb{K}}^{(a)} \quad ; \quad \sum_a \ell_a = \ell \quad (3.20)$$

Correspondingly we can arrange the ℓ functions $\mathfrak{h}_i(\vec{x})$ according to the graded decomposition (3.19), by writing:

$$\mathfrak{H}(\vec{x}) = \sum_{\alpha=0}^n \underbrace{\sum_{i=1}^{\ell_\alpha} \mathfrak{h}_i^{(\alpha)}(\vec{x}) A_\alpha^i}_{\in \mathbb{N}_{\mathbb{K}}^{(\alpha)}} \quad (3.21)$$

and equations (3.14) take the following triangular form:

$$\begin{aligned} \nabla^2 \mathfrak{h}_i^{(0)} &= 0 \\ \nabla^2 \mathfrak{h}_i^{(1)} &= \mathfrak{F}_i^{(1)} \left(\mathfrak{h}^{(0)}, \nabla \mathfrak{h}^{(0)} \right) \\ \nabla^2 \mathfrak{h}_i^{(2)} &= \mathfrak{F}_i^{(2)} \left(\mathfrak{h}^{(0)}, \nabla \mathfrak{h}^{(0)}, \mathfrak{h}^{(1)}, \nabla \mathfrak{h}^{(1)} \right) \\ \dots &= \dots \\ \nabla^2 \mathfrak{h}_i^{(n)} &= \mathfrak{F}_i^{(n)} \left(\mathfrak{h}^{(0)}, \nabla \mathfrak{h}^{(0)}, \mathfrak{h}^{(1)}, \nabla \mathfrak{h}^{(1)}, \dots, \mathfrak{h}^{(n-1)}, \nabla \mathfrak{h}^{(n-1)} \right), \end{aligned} \quad (3.22)$$

where ∇^2 denotes the three-dimensional Laplacian and at each level α , by $\mathfrak{F}_i^{(\alpha)}(\dots)$ we denote an $\mathfrak{so}(3)$ invariant polynomial of all the functions h^β up to level $\alpha - 1$ and of their derivatives.

Therefore the first ℓ_0 functions $\mathfrak{h}_i^{(0)}$ are just harmonic functions, while the higher ones satisfy Laplace equation with a source that is provided by the previously determined functions.

3.2 Transformation to the solvable gauge

Given the symmetric coset representative $\mathcal{Y}(\vec{x})$, parameterized by functions $\mathfrak{h}_i^{(\alpha)}(\vec{x})$ which satisfy the field equations (3.22), in order to retrieve the corresponding supergravity fields satisfying supergravity field equations, we need to solve a technical, yet quite crucial problem. We need to construct a new

upper triangular coset representative:

$$\mathbb{L}(\mathcal{Y}) = \begin{pmatrix} L_{1,1}(\mathcal{Y}) & L_{1,2}(\mathcal{Y}) & \cdots & L_{1,n-1}(\mathcal{Y}) & L_{1,n}(\mathcal{Y}) \\ 0 & L_{2,2}(\mathcal{Y}) & \cdots & L_{2,n-1}(\mathcal{Y}) & L_{2,n}(\mathcal{Y}) \\ 0 & 0 & L_{3,3}(\mathcal{Y}) & \cdots & L_{3,n}(\mathcal{Y}) \\ \vdots & \cdots & 0 & \cdots & \vdots \\ 0 & 0 & \cdots & 0 & L_{3,n}(\mathcal{Y}) \end{pmatrix} \quad (3.23)$$

which depends algebraically on the matrix entries of \mathcal{Y} and satisfies the following equivalence condition

$$\mathbb{L}(\mathcal{Y}) \mathcal{Q}(\mathcal{Y}) = \mathcal{Y} \quad ; \quad \mathcal{Q}(\mathcal{Y}) \in \mathbf{H}^* \quad (3.24)$$

where, as specified above, $\mathcal{Q}(\mathcal{Y})$ is a suitable element of the subgroup \mathbf{H}^* . It should be stressed that in the existing literature, this transition from the symmetric to the solvable gauge, which is compulsory in order to make the construction of the black hole solutions explicit, has been advocated, yet it has been to ad hoc procedures to be invented case by case.

Actually a universal and very elegant solution of such a problem exists and was found, from a different perspective, by the authors of the present paper in [23],[26],[24, 25],[28]. Indeed defining the following determinants:

$$\mathfrak{D}_i(\mathcal{Y}) := \text{Det} \begin{pmatrix} \mathcal{Y}_{1,1} & \cdots & \mathcal{Y}_{1,i} \\ \vdots & \vdots & \vdots \\ \mathcal{Y}_{i,1} & \cdots & \mathcal{Y}_{i,i} \end{pmatrix}, \quad \mathfrak{D}_0(\mathcal{Y}) := 1 \quad (3.25)$$

the matrix elements of the inverse of the upper triangular coset representative satisfying both equations (3.23) and (3.24) are given by the following expressions:

$$(\mathbb{L}(\mathcal{Y})^{-1})_{ij} \equiv \frac{1}{\sqrt{\mathfrak{D}_i(\mathcal{Y})\mathfrak{D}_{i-1}(\mathcal{Y})}} \text{Det} \begin{pmatrix} \mathcal{Y}_{1,1} & \cdots & \mathcal{Y}_{1,i-1} & \mathcal{Y}_{1,j} \\ \vdots & \vdots & \vdots & \vdots \\ \mathcal{Y}_{i,1} & \cdots & \mathcal{Y}_{i,i-1} & \mathcal{Y}_{i,j} \end{pmatrix} \quad (3.26)$$

Equation (3.26) provides a universal non-trivial and very elegant solution to the gauge-change problem and makes the entire construction based on harmonic functions truly algorithmic from the start to the very end.

3.3 Extraction of the three dimensional scalar fields

The result of the procedure described in the previous section is a triangular coset representative $\mathbb{L}(\mathfrak{h}_i^{(\alpha)})$ whose entries are polynomial and square root of polynomials in the functions $\mathfrak{h}_i^{(\alpha)}(x)$. The extraction of the scalar fields $\{U(x), a(x), Z(x), \phi(x)\}$ can now be performed according to the rules already presented in [28], which we recall here for completeness.

The general form of the solvable coset representative in terms of the fields is the following one:

$$\mathbb{L}(\Phi) = \exp[-a L_+^E] \exp[\sqrt{2} Z^M \mathcal{W}_M] \mathbb{L}_4(\phi) \exp[U L_0^E] \quad (3.27)$$

where L_0^E, L_\pm^E are the generators of the Ehlers group and $\mathcal{W}^M \equiv W^{1M}$ are the generators in the W -representation, according to the general structure (2.21) of the $\mathbb{U}_{D=3}$ Lie algebra; furthermore $\mathbb{L}_4(\phi)$ is the coset representative of the $D = 4$ scalar coset manifold immersed in the $\mathbb{U}_{D=3}$ group. From this structure, identifying $\mathbb{L}(\Phi) = \mathbb{L}(\mathfrak{h}_i^{(\alpha)})$ we deduce the following iterative procedure for the extraction of the relevant fields:

First of all we can determine the warp factor U by means of the following simple formula:

$$U(\mathfrak{h}) = \log \left[\frac{1}{2} \text{Tr} \left(\mathbb{L}(\mathfrak{h}) L_+^E \mathbb{L}^{-1}(\mathfrak{h}) L_-^E \right) \right] \quad (3.28)$$

Secondly we obtain the fields ϕ_i as follows. Defining the functionals

$$\Xi_i(\mathfrak{h}) = \text{Tr} \left(\mathbb{L}^{-1}(\mathfrak{h}) T_i \mathbb{L}(\tau) \right) \quad (3.29)$$

from the form of the coset representative (3.27) it follows that Ξ_i depend only on the $D = 4$ scalar fields and, according to the explicit form of the $D = 4$ coset, one can work out the scalar fields ϕ_i .

The knowledge of U, ϕ_i allows to define:

$$\Omega(\mathfrak{h}) = \mathbb{L}(\mathfrak{h}) \exp \left[-U L_0^E \right] \mathbb{L}_4(\phi)^{-1} \quad (3.30)$$

from which we extract the Z^M fields by means of the following formula:

$$Z^M(\mathfrak{h}) = \frac{1}{2\sqrt{2}} \text{Tr} \left[\Omega(\mathfrak{h}) \mathcal{W}_M^T \right] \quad (3.31)$$

where T means transposed. Finally the knowledge of $Z^M(\mathfrak{h})$ allows to extract the a field by means of the following trace:

$$a(\mathfrak{h}) = -\frac{1}{2} \text{Tr} \left[\Omega(\mathfrak{h}) \exp \left[-\sqrt{2} Z^M(\mathfrak{h}) \mathcal{W}_M \right] L_+^E \right] \quad (3.32)$$

3.4 General properties of the black hole solutions and structure of their poles

Having discussed the structure of supergravity solutions in terms of black-boxes that are a set of harmonic functions and of their descendants generated through the solution of the hierarchical equations (3.22), it is appropriate to study the general form of the geometries one obtains in this way and the properties of the available harmonic functions.

First of all, naming:

$$\mathfrak{W} = \exp[U(x)] \quad (3.33)$$

the warp factor that defines the 4-dimensional metric (2.35), we would like to investigate the general properties of the corresponding geometries. For the case where the Kaluza-Klein monopole is zero $\mathbf{A}^{[KK]} = 0$ we can write the general form of the curvature two-form of such spaces and therefore the intrinsic form of the Riemann tensor. Using the vielbein formalism introduced in eq.(2.36) we obtain:

$$\begin{aligned} \mathfrak{R}^{0i} &= -\mathfrak{W} \nabla^i \nabla_k \mathfrak{W} E^0 \wedge E^k - 2 \nabla^i \mathfrak{W} \nabla_k \mathfrak{W} E^0 \wedge E^k \\ \mathfrak{R}^{ij} &= -2 \mathfrak{W} \nabla^{[i} \nabla_k \mathfrak{W} E^{j]} \wedge E^k + (\nabla \mathfrak{W} \cdot \nabla \mathfrak{W}) \nabla_k \mathfrak{W} E^i \wedge E^j \end{aligned} \quad (3.34)$$

where the derivatives used in the above equations are defined as follows. Let the flat metric in three dimension be described by a euclidian *dreibein* e^i such that:

$$\begin{aligned} ds_{flat}^2 &= \sum_{i=1}^3 e^i \otimes e^i \\ E^i &= \frac{1}{\mathfrak{W}} e^i \end{aligned} \quad (3.35)$$

then the total differential of the warp factor expanded along e^i yields the derivatives $\nabla_k \mathfrak{W}$, namely:

$$d\mathfrak{W} = \nabla_k \mathfrak{W} e^k \quad (3.36)$$

Next let us consider the general form of harmonic functions. These latter form a linear space since any linear combination of harmonic functions is still harmonic. There are three types of building blocks that we can use:

a Real center pole:

$$\mathcal{H}_\alpha(\vec{x}) = \frac{1}{|\vec{x} - \vec{x}_\alpha|} \quad (3.37)$$

b Real part of an imaginary center pole:

$$\mathcal{R}_\alpha(\vec{x}) = \text{Re} \left[\frac{1}{|\vec{x} - i\vec{x}_\alpha|} \right] \quad (3.38)$$

c Imaginary part of an imaginary center pole:

$$\mathcal{J}_\alpha(\vec{x}) = \text{Im} \left[\frac{1}{|\vec{x} - i\vec{x}_\alpha|} \right] \quad (3.39)$$

Hence the most general harmonic function can be written as the following sum:

$$\text{Harm}(\vec{x}) = h_\infty + \sum_\alpha \frac{p_\alpha}{|\vec{x} - \vec{x}_\alpha|} + \sum_\beta q_\beta \text{Re} \left[\frac{1}{|\vec{x} - i\vec{x}_\beta|} \right] + \sum_\gamma k_\gamma \text{Im} \left[\frac{1}{|\vec{x} - i\vec{x}_\gamma|} \right] \quad (3.40)$$

where the constant h_∞ is the boundary value of the harmonic function at infinity far from all the poles. In order to study the behavior of $\text{Harm}(\vec{x})$ in the vicinity of a real pole ($|\vec{x} - \vec{x}_\alpha| \ll 1$) it is convenient to adopt local polar coordinates:

$$\begin{aligned} x^1 - x_\alpha^1 &= r \cos \theta \\ x^2 - x_\alpha^2 &= r \sin \theta \sin \phi \\ x^3 - x_\alpha^3 &= r \sin \theta \cos \phi \end{aligned} \quad (3.41)$$

In this coordinates the harmonic function is approximated by:

$$\text{Harm}(\vec{x}) \simeq h_\alpha + \frac{p_\alpha}{r} \quad (3.42)$$

where the effective constant h_α encodes the finite part of the function contributed by all the other poles. In polar coordinates the Laplacian operator on functions of r becomes:

$$\Delta = \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} \quad (3.43)$$

The general outcome of the construction procedure outlined in the previous section is that the warp factor is the square root of a rational function of n harmonic functions, where $n = \dim \mathbb{N}_\mathbb{K}$

$$\mathfrak{W}(\vec{x}) = \sqrt{\frac{\mathbb{P}(\widehat{\text{Harm}}_1(\vec{x}), \dots, \widehat{\text{Harm}}_n(\vec{x}))}{\mathbb{Q}(\widehat{\text{Harm}}_1(\vec{x}), \dots, \widehat{\text{Harm}}_n(\vec{x}))}} \quad (3.44)$$

where \mathbb{P} and \mathbb{Q} are two polynomials. By $\widehat{\text{Harm}}_1(\vec{x})$ we denote both harmonic functions and their descendants generated by the hierarchical system (3.22). For a given multicenter solution it is convenient to enumerate all the poles displayed by one or the other of the harmonic functions and in the vicinity of each of those poles we will have:

$$\widehat{\text{Harm}}_i(\vec{x}) \simeq \frac{p_i}{r^{m_i}} \quad (3.45)$$

where $p_i \neq 0$ if the considered pole belongs to the considered function and it is zero otherwise. Furthermore if $\text{Harm}_i(\vec{x})$ is one of the level one harmonic function the exponent $m_i = 1$. Otherwise it is bigger, but in any case $m_i \geq 1$. Taking this into account the effective behavior of the warp factor will always be of the following form:

$$\mathfrak{W}(\vec{x}) \simeq r^{\ell_\alpha} \sqrt{c_\alpha} \quad (3.46)$$

where ℓ is some integer or half integer power (positive or negative) and c_α is a constant. In order for the pole to be a regular point of the solution, two conditions have to be satisfied:

1. The constant $c_\alpha > 0$ must be positive so that the warp factor is real.
2. The power $\ell_\alpha \geq 1$ so that the Riemann tensor does not diverge at the pole.

The second condition follows from the form (3.34) of the Riemann tensor which implies that all of its components behave as:

$$\mathfrak{R}^{ab}_{cd} \simeq r^{2\ell_\alpha-2} \times \text{const} \quad (3.47)$$

Near the pole the metric behaves as follows:

$$ds^2 \simeq -\sqrt{c_\alpha} r^{\ell_\alpha} dt^2 + \frac{1}{\sqrt{c_\alpha}} \frac{1}{r^{\ell_\alpha}} [dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)] \quad (3.48)$$

In order for the pole to be an event horizon of finite or of vanishing area, we must have $2 - \ell_\alpha > 0$, so that the volume of the two-sphere described by $(d\theta^2 + \sin^2 \theta d\phi^2)$ does not diverge. Hence for regular black holes we have only three possibilities:

$$\underbrace{\ell_\alpha = 2}_{\text{Large Black Holes}} \quad ; \quad \underbrace{\ell_\alpha = \frac{3}{2}}_{\text{Small Black Holes}} \quad ; \quad \underbrace{\ell_\alpha = 1}_{\text{Very Small Black Holes}} \quad (3.49)$$

When we are in the case of Large Black Holes, the near horizon geometry is approximated by that:

$$\text{AdS}_2 \times \mathbb{S}^2 \quad (3.50)$$

The case of the harmonic functions with an imaginary center requires a different treatment. Their near singularity behavior is best analyzed by using spheroidal coordinates.

These are easily introduced by setting:

$$\begin{aligned} x^1 &= \sqrt{r^2 + \alpha^2} \sin \theta \sin \phi \\ x^2 &= \sqrt{r^2 + \alpha^2} \sin \theta \cos \phi \\ x^3 &= r \cos \theta \end{aligned} \quad (3.51)$$

where r, θ, ϕ are the new coordinates and α is a deformation parameter which represents the position of the center in the complex plane. In terms of these coordinates the flat euclidian three-dimensional metric takes the following form:

$$ds_{\mathbb{E}^3}^2 = d\Omega_{spheroidal}^2 \equiv \frac{(r^2 + \alpha^2 \cos^2 \theta) dr^2}{r^2 + \alpha^2} + (r^2 + \alpha^2) \sin^2 \theta d\phi^2 + (r^2 + \alpha^2 \cos^2 \theta) d\theta^2 \quad (3.52)$$

and the two harmonic functions that correspond to the real and imaginary part of a complex harmonic function with center on the imaginary z -axis at α -distance from zero are:

$$\mathcal{P}_\alpha(r, \theta) = \frac{r}{r^2 + \alpha^2 \cos^2 \theta} \quad (3.53)$$

$$\mathcal{R}_\alpha(r, \theta) = \frac{\alpha \cos \theta}{r^2 + \alpha^2 \cos^2 \theta} \quad (3.54)$$

and the Hodge duals of their gradients, in spheroidal coordinates have the following form:

$$\star \nabla \mathcal{P}_\alpha = \frac{\sin \theta}{(r^2 + \alpha^2 \cos^2 \theta)^2} [2\alpha^2 r \cos \theta \sin \theta dr \wedge d\phi + (r^2 + \alpha^2) (r^2 - \alpha^2 \cos^2 \theta) d\theta \wedge d\phi] \quad (3.55)$$

$$\star \nabla \mathcal{R}_\alpha = \frac{\alpha \sin \theta}{(r^2 + \alpha^2 \cos^2 \theta)^2} [(\alpha^2 \cos^2 \theta - r^2) \sin \theta dr \wedge d\phi + 2r (r^2 + \alpha^2) \cos \theta d\theta \wedge d\phi] \quad (3.56)$$

These are the building blocks we can use to construct Kerr-Newman like solutions and we shall outline a pair of examples in the sequel.

4 The example of the S^3 model: classification of the nilpotent orbits

As an illustration of the general procedure we explore the case of the S^3 model, leading to the $G_{2,2}$ group in $D = 3$. The spherical symmetric black hole solutions of this model were already discussed in a full-fledged way in [28], and in [29] the detailed classification of the corresponding nilpotent orbits was derived. Here we reconsider the same case from the point of view of the multicenter construction outlined in the previous section, relying on the results of [29]. According to that paper, for the case of the coset manifold²:

$$\frac{U_{D=3}}{H^\star} = \frac{G_{(2,2)}}{\widehat{SL(2, \mathbb{R})} \times SL(2, \mathbb{R})_{h^\star}} \quad (4.1)$$

we have just seven distinct nilpotent orbits of the $H^\star = \widehat{SL(2, \mathbb{R})} \times SL(2, \mathbb{R})_{h^\star}$ subgroup in the \mathbb{K}^\star representation $(2, \frac{3}{2})$, which are enumerated by the three set of labels $\alpha\beta\gamma$ and are denoted $\mathcal{O}_{\beta\gamma}^\alpha$ as described in table 1³. An explicit choice of a representative for each of the seven orbits is provided

²For the rationale of our notation we refer the reader to the later section 9.

³For the notation and organization of the $\alpha\beta\gamma$ labels we refer the reader to [29].

below.

$$\mathcal{O}_{11}^1 = \begin{pmatrix} \frac{\sqrt{3}}{2} & \frac{\sqrt{5}}{2} & \sqrt{\frac{3}{2}} & \frac{\sqrt{5}}{2} & 0 & \frac{\sqrt{5}}{2} & 0 \\ \frac{\sqrt{5}}{2} & \sqrt{6} & -\frac{\sqrt{5}}{2} & -\sqrt{3} & -\frac{\sqrt{5}}{2} & 0 & \frac{\sqrt{5}}{2} \\ -\sqrt{\frac{3}{2}} & \frac{\sqrt{5}}{2} & -\sqrt{\frac{3}{2}} & \frac{\sqrt{5}}{2} & 0 & \frac{\sqrt{5}}{2} & 0 \\ -\frac{\sqrt{5}}{2} & \sqrt{3} & \frac{\sqrt{5}}{2} & 0 & \frac{\sqrt{5}}{2} & -\sqrt{3} & -\frac{\sqrt{5}}{2} \\ 0 & \frac{\sqrt{5}}{2} & 0 & \frac{\sqrt{5}}{2} & \sqrt{\frac{3}{2}} & \frac{\sqrt{5}}{2} & \sqrt{\frac{3}{2}} \\ \frac{\sqrt{5}}{2} & 0 & -\frac{\sqrt{5}}{2} & \sqrt{3} & -\frac{\sqrt{5}}{2} & -\sqrt{6} & \frac{\sqrt{5}}{2} \\ 0 & \frac{\sqrt{5}}{2} & 0 & \frac{\sqrt{5}}{2} & -\sqrt{\frac{3}{2}} & \frac{\sqrt{5}}{2} & -\sqrt{\frac{3}{2}} \end{pmatrix} \quad (4.2)$$

$$\mathcal{O}_{11}^4 = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{2} & 0 & 0 & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & 0 & 0 & -\frac{1}{2} \end{pmatrix} \quad (4.3)$$

$$\mathcal{O}_{11}^2 = \begin{pmatrix} \frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ \frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{2} & 0 & -\frac{1}{2} \\ 0 & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{2} & 0 & -\frac{1}{2} \end{pmatrix} \quad (4.4)$$

$$\mathcal{O}_{11}^3 = \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 1 & -\frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & -\frac{1}{2} & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & -\frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{2} & 0 & -\frac{1}{2} & -1 & 0 \\ 0 & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & -1 \end{pmatrix} \quad (4.5)$$

N	d_n	α – label	$\gamma\beta$ – labels	Orbits			\mathcal{W}_H – classes
1	7	[j=3]	$\gamma\beta_1 = \{8_1 4_1 0_1\}$	\mathcal{O}_1^1			$(\times, \gamma_1, \times)$
2	3	[j=1]×2[j=1/2]	$\gamma\beta_1 = \{3_1 1_1 0_1\}$	\mathcal{O}_1^2			$(\gamma_1, \gamma_1, \times)$
7	3	2[j=1]×[j=0]	$\gamma\beta_1 = \{4_1 0_2\}$ $\gamma\beta_2 = \{2_2 0_1\}$		β_1	β_2	$(\gamma_1, \gamma_2, \gamma_2)$
				γ_1	$\mathcal{O}_{1,1}^3$	$\mathcal{O}_{1,2}^3$	
				γ_2	$\mathcal{O}_{2,1}^3$	$\mathcal{O}_{2,2}^3$	
4	2	2[j=1/2]×3 [j=0]	$\gamma\beta_1 = \{1_2 0_1\}$	\mathcal{O}_1^4			$(0, \gamma_1, \gamma_1)$

Table 1: Classification of the nilpotent orbits of $\frac{G_{(2,2)}}{SU(1,1) \times SU(1,1)}$.

$$\mathcal{O}_{22}^3 = \begin{pmatrix} 1 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 1 & -\frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 \\ -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & -\frac{1}{\sqrt{2}} \\ 0 & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & -\frac{1}{2} & -1 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & -1 \end{pmatrix} \quad (4.6)$$

$$\mathcal{O}_{21}^3 = \begin{pmatrix} -1 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & -1 & 0 & 0 & \frac{1}{2} & 0 \\ -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & -\frac{1}{\sqrt{2}} \\ 0 & \frac{1}{2} & 0 & 0 & 1 & \frac{1}{2} & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 1 \end{pmatrix} \quad (4.7)$$

$$\mathcal{O}_{12}^3 = \begin{pmatrix} -1 & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & -1 & 0 & 0 & -\frac{1}{2} & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & -\frac{1}{2} & 0 & 0 & 1 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{2} & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & 1 \end{pmatrix} \quad (4.8)$$

Note that, in some instances, these representatives do not coincide with the representatives shown in our previous papers [28], [29]. The reason is that in the approach pursued in the present paper, it is no

longer relevant to consider representatives possessing vanishing Taub-NUT charges. The vanishing of the Taub-NUT current will be anyhow implemented on the parametrization of the symmetric coset representative associated with the nilpotent orbit. So we rather prefer to choose the simplest representatives of each nilpotent orbit postponing the issue of the Taub-NUT charge at a later stage.

Each orbit representative $\mathcal{O}_{\beta\gamma}^\alpha$ identifies a standard triple $\{h, X, Y\}$ and hence an embedding of an $\mathfrak{sl}(2, \mathbb{R})$ Lie algebra:

$$[h, X] = 2X \quad ; \quad [h, Y] = -2Y \quad ; \quad [X, Y] = 2h \quad (4.9)$$

into $\mathfrak{g}_{(2,2)}$ in such a way that $h \in \mathbb{H}^*$ and $X, Y \in \mathbb{K}^*$. The triple is obtained by setting:

$$X_{\alpha|\beta\gamma} \equiv \mathcal{O}_{\beta\gamma}^\alpha \quad ; \quad Y_{\alpha|\beta\gamma} \equiv X_{\alpha|\beta\gamma}^T \quad ; \quad h_{\alpha|\beta\gamma} \equiv [X_{\alpha|\beta\gamma}, Y_{\alpha|\beta\gamma}] \quad (4.10)$$

The relevant item in the construction of solutions based on the integration of equations in the symmetric gauge is provided by the central element of the triple $h_{\alpha|\beta\gamma}$ which defines the gradings. In the present example of the S^3 model, it turns out the orbits having the same α and γ labels but different β -labels have the same central element, namely:

$$h_{\alpha|\beta\gamma} = h_{\alpha|\beta'\gamma} \quad (4.11)$$

so that the solutions pertaining both to orbit $\mathcal{O}_{\beta\gamma}^\alpha$ and to orbit $\mathcal{O}_{\beta'\gamma}^\alpha$ are obtained from the same construction and are distinguished only by different choices in the space of the available harmonic functions parameterizing the general solution.

The explicit form of the central elements are the following ones:

Large Orbit \mathcal{O}_{11}^1 : central element

$$h_{1|11} = \begin{pmatrix} 0 & 0 & -1 & 0 & 5 & 0 & 0 \\ 0 & 0 & 0 & \sqrt{2} & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 5 \\ 0 & \sqrt{2} & 0 & 0 & 0 & \sqrt{2} & 0 \\ 5 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & \sqrt{2} & 0 & 0 & 0 \\ 0 & 0 & 5 & 0 & -1 & 0 & 0 \end{pmatrix}$$

$$\text{Eigenvalues } \left[\frac{1}{2} h_{1|11} \right] = \{-3, 3, -2, 2, -1, 1, 0\} \quad (4.12)$$

Very Small Orbit \mathcal{O}_{11}^4 : central element

$$h_{4|11} = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\text{Eigenvalues } \left[\frac{1}{2} h_{4|11} \right] = \left\{ -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0 \right\} \quad (4.13)$$

Small Orbit \mathcal{O}_{11}^2 : central element

$$h_{2|11} = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\sqrt{2} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\sqrt{2} & 0 & 0 & 0 & -\sqrt{2} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -\sqrt{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\text{Eigenvalues } \left[\frac{1}{2} h_{2|11} \right] = \left\{ -1, 1, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0 \right\} \quad (4.14)$$

Large BPS Orbit \mathcal{O}_{11}^3 : central element

$$h_{3|11} = h_{3|21} = \begin{pmatrix} 0 & 0 & 0 & \sqrt{2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ \sqrt{2} & 0 & 0 & 0 & 0 & 0 & -\sqrt{2} \\ 0 & -1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -\sqrt{2} & 0 & 0 & 0 \end{pmatrix}$$

$$\text{Eigenvalues } \left[\frac{1}{2} h_{3|11} \right] = \{-1, -1, 1, 1, 0, 0, 0\} \quad (4.15)$$

Large non BPS Orbit \mathcal{O}_{22}^3 : central element

$$h_{3|12} = h_{3|22} = \begin{pmatrix} 0 & 0 & 0 & -\sqrt{2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & -1 & 0 \\ -\sqrt{2} & 0 & 0 & 0 & 0 & 0 & \sqrt{2} \\ 0 & 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & \sqrt{2} & 0 & 0 & 0 \end{pmatrix}$$

$$\text{Eigenvalues } \left[\frac{1}{2} h_{3|22} \right] = \{-1, -1, 1, 1, 0, 0, 0\} \quad (4.16)$$

5 Explicit construction of the multicenter Black Holes solutions of the S^3 model

Having enumerated the central elements for the independent orbits we proceed to the construction and discussion of the corresponding black hole solutions, whose properties are summarized in table (2).

5.1 The Very Small Black Holes of \mathcal{O}_{11}^4

We begin with the smallest orbits which, in a sense that will become clear further on, represent the elementary blocks in terms of which bigger black holes are constructed.

Focusing on any orbit $\mathcal{O}_{\beta\gamma}^\alpha$ and considering the nilpotent element of the corresponding triple $X_{\alpha|\beta\gamma} \in \mathbb{K}^\star$ as a Lax operator L_0 , we easily workout the electromagnetic charges by calculating the traces displayed below (see section 10, for more explanations)

$$\mathcal{Q}^\mathbf{w} = \text{Tr}(X_{\alpha|\beta\gamma} \mathcal{T}^\mathbf{w}) \quad (5.1)$$

W-representation In the case of the orbit \mathcal{O}_{11}^4 we obtain:

$$\mathcal{Q}_{4|11}^\mathbf{w} = (0, 0, 0, 1) \quad (5.2)$$

Substituting such a result in the expression for the quartic symplectic invariant (see [28]):

$$\mathfrak{I}_4 = \frac{1}{4} \left(4\sqrt{3}Q_4Q_1^3 + 3Q_3^2Q_1^2 - 18Q_2Q_3Q_4Q_1 - Q_2 \left(4\sqrt{3}Q_3^3 + 9Q_2Q_4^2 \right) \right) \quad (5.3)$$

of the **W** representation which happens to be the spin $\frac{3}{2}$ of $\mathfrak{sl}(2, \mathbb{R})$ we find:

$$\mathfrak{I}_4 = 0 \quad (5.4)$$

The result is meaningful since, by calculating the trace $\text{Tr}(X_{4|11}L_+^E) = 0$, we can also check that the Taub-NUT charge vanishes. We can also address the question whether there are subgroups of the original duality group in four-dimensions $\text{SL}(2, \mathbb{R})$ that leave the charge vector (5.2) invariant. Using the explicit form of the $j = \frac{3}{2}$ representation displayed in eq.(3.13) of [28], we realize that indeed such group exists and it is the parabolic subgroup described below:

$$\forall c \in \mathbb{R} \quad : \quad \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} \in \mathcal{S}_{4|11} \subset \text{SL}(2, \mathbb{R}) \quad (5.5)$$

This stability subgroup together with the vanishing of the quartic invariant are the intrinsic definition of the **W**-orbit pertaining to very small black holes.

H^{*}-stability subgroup In a parallel way we can pose the question what is the stability subgroup of the nilpotent element $X_{4|11}$ in $\text{H}^\star = \widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{\text{h}^\star}$ (For further explanations on H^\star and its structure see section 9). The answer is the following:

$$\mathfrak{S}_{4|11} = \text{ISO}(1, 1) \quad (5.6)$$

Name of orbit	pq charges	Quart. Inv. \mathcal{I}_4	\mathbf{W} – stab. group $\mathcal{S}_{\mathbf{W}} \subset \mathfrak{sl}(2, \mathbb{R})$	\mathbf{H}^* – stab. group $\mathfrak{S}_{\mathbf{H}^*} \subset \widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{\mathbf{h}^*}$	dim \mathbb{N}	dim $\mathbb{N} \cap \mathbb{K}^*$
\mathcal{O}_{11}^4	$\begin{pmatrix} 0 \\ 0 \\ 0 \\ q \end{pmatrix}$	0	$\begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$	$\underbrace{\text{ISO}(1, 1)}_{3 \text{ gen.}}$	3	3
\mathcal{O}_{11}^2	$\begin{pmatrix} \sqrt{3}p \\ 0 \\ 0 \\ 0 \end{pmatrix}$	0	$\mathbf{1}$	$\underbrace{\text{SO}(1, 1) \triangleright \mathbb{R}}_{2 \text{ gen.}}$	4	3
\mathcal{O}_{11}^3	$\begin{pmatrix} 0 \\ p \\ -\sqrt{3}q \\ 0 \end{pmatrix}$	$9pq^3 > 0$	\mathbb{Z}_3	$\underbrace{\mathbb{R}}_{1 \text{ gen. } A^2 = 0}$	5	4
\mathcal{O}_{22}^3	$\begin{pmatrix} 0 \\ p \\ \sqrt{3}q \\ 0 \end{pmatrix}$	$-9pq^3 < 0$	$\mathbf{1}$	$\underbrace{\mathbb{R}}_{1 \text{ gen. } A^3 = 0}$	3	3
\mathcal{O}_{11}^1	$\begin{pmatrix} \frac{1}{2}\sqrt{\frac{3}{2}}p \\ 0 \\ \frac{7}{6}p \\ \sqrt{2}q \end{pmatrix}$	$\frac{1}{128}p^3 \times$ $(49p + 72q)$	$\mathbf{1}$	$\mathbf{1}$	6	4

Table 2: Properties of the $\mathfrak{g}_{(2,2)}$ orbits in the S^3 model. The structure of the electromagnetic charge vector is that obtained for solutions with vanishing Taub-NUT current. The symbol \triangleright is meant to denote semidirect product. $\mathcal{S}_{\mathbf{W}}$ denotes the subgroup of the $D = 4$ duality group which leaves the charge vector invariant, while $\mathfrak{S}_{\mathbf{H}^*}$ denotes the subgroup of the \mathbf{H}^* isotropy group of the $D = 3$ sigma-model which leaves invariant the X element of the standard triple. This latter is the Lax operator in the one-dimensional spherical symmetric approach.

A generic element of the corresponding Lie algebra is a linear combination of three generators J, T_1, T_2 , satisfying the commutation relations:

$$[J, T_1] = \frac{1}{\sqrt{2}} T_1 + \frac{3}{2\sqrt{6}} T_2$$

$$[J, T_2] = \frac{3}{2\sqrt{2}} T_1 \quad ; \quad [T_1, T_2] = 0 \quad (5.7)$$

It is explicitly given by the following matrix:

$$\omega J + x T_1 + y T_2 = \begin{pmatrix} 0 & -\frac{x}{2\sqrt{2}} & \frac{\omega}{2\sqrt{2}} & -\frac{x}{2} & 0 & -\frac{1}{2}\sqrt{\frac{3}{2}}y & 0 \\ \frac{x}{2\sqrt{2}} & 0 & -\frac{1}{2}\sqrt{\frac{3}{2}}y & -\frac{\omega}{2} & \frac{x}{2\sqrt{2}} & 0 & -\frac{1}{2}\sqrt{\frac{3}{2}}y \\ \frac{\omega}{2\sqrt{2}} & -\frac{1}{2}\sqrt{\frac{3}{2}}y & 0 & -\frac{x}{2} & 0 & -\frac{x}{2\sqrt{2}} & 0 \\ -\frac{x}{2} & -\frac{\omega}{2} & \frac{x}{2} & 0 & -\frac{x}{2} & -\frac{\omega}{2} & \frac{x}{2} \\ 0 & \frac{x}{2\sqrt{2}} & 0 & \frac{x}{2} & 0 & \frac{1}{2}\sqrt{\frac{3}{2}}y & \frac{\omega}{2\sqrt{2}} \\ \frac{1}{2}\sqrt{\frac{3}{2}}y & 0 & -\frac{x}{2\sqrt{2}} & -\frac{\omega}{2} & \frac{1}{2}\sqrt{\frac{3}{2}}y & 0 & -\frac{x}{2\sqrt{2}} \\ 0 & \frac{1}{2}\sqrt{\frac{3}{2}}y & 0 & \frac{x}{2} & \frac{\omega}{2\sqrt{2}} & \frac{x}{2\sqrt{2}} & 0 \end{pmatrix} \quad (5.8)$$

Nilpotent algebra $\mathbb{N}_{4|11}$ Considering next the adjoint action of the central element $h_{4|11}$ on the subspace \mathbb{K}^* we find that its eigenvalues are the following ones:

$$\text{Eigenvalues}_{4|11}^{\mathbb{K}^*} = \{-2, 2, -1, -1, 1, 1, 0, 0\} \quad (5.9)$$

Therefore the three eigenoperators A_1, A_2, A_3 corresponding to the positive eigenvalues 2, 1, 1, respectively, form the restriction to \mathbb{K}^* of a nilpotent algebra $\mathbb{N}_{4|11}$. In this case A_i commute among themselves so that $\mathbb{N}_{4|11} = \mathbb{N}_{4|11} \cap \mathbb{K}^*$ and it is abelian. This structure of the nilpotent algebra implies that for the orbit \mathcal{O}_{11}^4 we have only three functions \mathfrak{h}_i^0 which will be harmonic and independent.

Explicitly we set:

$$\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3) = \sum_{i=1}^3 \mathfrak{h}_i A_i = \begin{pmatrix} -\mathfrak{h}_1 & \mathfrak{h}_3 & 0 & -\sqrt{2}\mathfrak{h}_3 & -\mathfrak{h}_1 & -\mathfrak{h}_2 & 0 \\ \mathfrak{h}_3 & 0 & -\mathfrak{h}_2 & 0 & \mathfrak{h}_3 & 0 & -\mathfrak{h}_2 \\ 0 & \mathfrak{h}_2 & -\mathfrak{h}_1 & \sqrt{2}\mathfrak{h}_3 & 0 & -\mathfrak{h}_3 & -\mathfrak{h}_1 \\ \sqrt{2}\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_3 \\ \mathfrak{h}_1 & -\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_3 & \mathfrak{h}_1 & \mathfrak{h}_2 & 0 \\ -\mathfrak{h}_2 & 0 & \mathfrak{h}_3 & 0 & -\mathfrak{h}_2 & 0 & \mathfrak{h}_3 \\ 0 & -\mathfrak{h}_2 & \mathfrak{h}_1 & -\sqrt{2}\mathfrak{h}_3 & 0 & \mathfrak{h}_3 & \mathfrak{h}_1 \end{pmatrix} \quad (5.10)$$

Considering $\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3)$ as a Lax operator and calculating its Taub-NUT charge and electromagnetic charges we find:

$$\mathbf{n}_{TN} = -2\mathfrak{h}_2 \quad ; \quad \mathcal{Q} = (0, 2\mathfrak{h}_2, -2\sqrt{3}\mathfrak{h}_3, -2\mathfrak{h}_1) \quad (5.11)$$

This implies that constructing the multi-centre solution with harmonic functions the condition $\mathfrak{h}_2 = 0$ should be sufficient to annihilate the Taub-NUT current.

For later convenience let us change the normalization in the basis of harmonic functions as follows:

$$\mathfrak{h}_1^{(0)} = \frac{1}{\sqrt{2}}\mathcal{H}_1 \quad ; \quad \mathfrak{h}_2^{(0)} = \frac{1}{2}(1 - \mathcal{H}_2) \quad ; \quad \mathfrak{h}_3^{(0)} = \frac{1}{\sqrt{2}}\mathcal{H}_3 \quad (5.12)$$

Implementing the symmetric coset construction with:

$$\mathcal{Y}(\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3) \equiv \exp \left[\mathfrak{H} \left(\frac{1}{\sqrt{2}} \mathcal{H}_1, \frac{1}{2} (1 - \mathcal{H}_2), \frac{1}{\sqrt{2}} \mathcal{H}_3 \right) \right] \quad (5.13)$$

and calculating the upper triangular coset representative $\mathbb{L}(\mathcal{Y})$ according to equations (3.26) we find a relatively simple expression which, however, is still too large to be displayed. Yet the extraction of the σ -model scalar fields produces a quite compact answer which we list below:

$$\exp[-U] = \sqrt{\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1} \quad (5.14)$$

$$\text{Im } z = \frac{\sqrt{\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1}}{\mathcal{H}_2^2 - \mathcal{H}_3^2 + \mathcal{H}_1} \quad (5.15)$$

$$\text{Re } z = -\frac{\sqrt{2}\mathcal{H}_3}{\mathcal{H}_2^2 - \mathcal{H}_3^2 + \mathcal{H}_1} \quad (5.16)$$

$$Z^M = \begin{pmatrix} \frac{\sqrt{6}\mathcal{H}_3^2}{\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1} \\ \frac{(\mathcal{H}_2 - 2\mathcal{H}_3)(\mathcal{H}_2 + \mathcal{H}_3)^2 + \mathcal{H}_1\mathcal{H}_2}{\sqrt{(\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1)^2}} \\ -\frac{\sqrt{3}\mathcal{H}_3}{\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1} \\ \frac{\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1 - 1}{\sqrt{2}(\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1)} \end{pmatrix} \quad (5.17)$$

$$a = \frac{\mathcal{H}_2^3 + (-3\mathcal{H}_3^2 + \mathcal{H}_1 + 1)\mathcal{H}_2 - 2\mathcal{H}_3^3}{\sqrt{2}(\mathcal{H}_2^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1)} \quad (5.18)$$

The Taub-NUT current Given this explicit result we can turn to the explicit oxidation formulae described in section 2.2 and calculate the Taub-NUT current which is the integrand of eq.(2.41). We find:

$$j^{TN} = \sqrt{2} \star \nabla \mathcal{H}_2 \quad (5.19)$$

Hence the vanishing of the Taub-NUT current is guaranteed by the very simple condition:

$$\mathcal{H}_2 = \alpha \quad ; \quad \nabla \mathcal{H}_2 = 0 \quad (5.20)$$

where α is just a constant. This confirms the preliminary analysis obtained from the Lax operator which requires a vanishing component of the Lax along the second generator A_2 of the nilpotent algebra.

General form of the solution Imposing this condition we arrive at the following form of the solution depending on two harmonic functions $\mathcal{H}_1, \mathcal{H}_3$:

$$\exp[-U] = \sqrt{\alpha^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1} \quad (5.21)$$

$$z = i \frac{1}{\sqrt{\alpha^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1}} - \frac{\sqrt{2}\mathcal{H}_3}{\alpha^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1} \quad (5.22)$$

$$j^{TN} = 0 \quad (5.23)$$

$$j^{EM} = \star \nabla \begin{pmatrix} 0 \\ 0 \\ \sqrt{3} \mathcal{H}_3 \\ -\frac{1}{\sqrt{2}} \mathcal{H}_1 \end{pmatrix} \quad (5.24)$$

Obviously the physical range of the solution is determined by the condition $(\alpha^2 - 3\mathcal{H}_3^2 + \mathcal{H}_1) > 0$ which can always be arranged, by tuning the parameters contained in the harmonic functions.

To this effect let us discuss the nature of the black holes encompassed by this solution, that, by definition, are located at the poles of the harmonic functions $\mathcal{H}_1, \mathcal{H}_3$.

According to the argument developed in section 3.4, in the vicinity of each pole $|\vec{x} - \vec{x}_I| = r < \epsilon$ we can choose polar coordinates centered at \vec{x}_α and the behavior of the harmonic functions, for $\epsilon \rightarrow 0$ is the following one:

$$\mathcal{H}_1 \sim a_1 + \frac{b_1}{r} \quad (5.25)$$

$$\mathcal{H}_3 \sim a_3 + \frac{b_3}{r} \quad (5.26)$$

which corresponds to the following behavior of the warp factor:

$$\exp[-U] \sim \sqrt{\alpha^2 - 3a_3^2 - \frac{3b_3^2}{r^2} + a_1 + \frac{b_1}{r} - \frac{6a_3b_3}{r}} \quad (5.27)$$

In order for the warp factor to be real for all values of $r \rightarrow 0$ we necessarily find

$$\begin{aligned} b_3 &= 0 \\ b_1 &> 0 \\ \alpha^2 - 3a_3^2 + a_1 &> 0 \end{aligned} \quad (5.28)$$

$$(5.29)$$

Since conditions (5.28) hold true for each available pole, it means the harmonic function \mathcal{H}_3 has actually no pole and is therefore equal to some constant. The boundary condition of asymptotic flatness fixes the value of such a constant:

$$\lim_{r \rightarrow \infty} \exp[-U] = 1 \quad \Leftrightarrow \quad \mathcal{H}_3 = \frac{\sqrt{\alpha^2 + \mathcal{H}_1(\infty) - 1}}{\sqrt{3}} \quad (5.30)$$

Under such conditions in the vicinity of each pole \vec{x}_α , the warp factor has the following behavior:

$$|\vec{x} - \vec{x}_\alpha|^2 \exp[-U] \stackrel{\vec{x} \rightarrow \vec{x}_\alpha}{\sim} \sqrt{b_1} |\vec{x} - \vec{x}_\alpha|^{3/2} + \mathcal{O}(|\vec{x} - \vec{x}_\alpha|^{5/2}) \quad (5.31)$$

leading to a vanishing horizon area:

$$\text{Area}_{H_\alpha} = \lim_{\vec{x} \rightarrow \vec{x}_\alpha} |\vec{x} - \vec{x}_\alpha|^2 \exp[-U] = 0 \quad (5.32)$$

At the same time using the form of the electromagnetic current in eq.(5.24) and the behavior of the harmonic function in the vicinity of the poles we obtain the charge vector of each black hole encompassed by the solution:

$$\mathcal{Q}_\alpha = \int_{\mathbb{S}_\alpha^2} j^{EM} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ -\frac{1}{\sqrt{2}} q_\alpha \end{pmatrix} ; \quad \text{where } q_\alpha = b_1 \text{ for pole } \vec{x}_\alpha \quad (5.33)$$

Summarizing For the regular multicenter solutions associated with the orbit 4|11 all black holes localized at each pole are of the same type, namely they are very small black holes with vanishing horizon area and a charge vector \mathcal{Q} belonging to \mathbf{W} -orbit which is characterized by both a vanishing quartic invariant and the existence of a continuous parabolic stability subgroup of $\text{SL}(2, \mathbb{R})$. Every black hole is a repetition in a different place of the spherical symmetric black hole which gives its name to the orbit.

5.2 The Small Black Holes of \mathcal{O}_{11}^2

Next let us consider the orbit \mathcal{O}_{11}^2 .

W-representation Applying the same strategy as in the previous case, from the general formula we obtain

$$\mathcal{Q}_{2|11}^{\mathbf{W}} = \text{Tr}(X_{2|11} \mathcal{T}^{\mathbf{W}}) = (\sqrt{3}, 0, 0, 0) \quad (5.34)$$

Substituting such a result in the expression for the quartic symplectic invariant (see eq.(5.3) we find:

$$\mathfrak{I}_4 = 0 \quad (5.35)$$

Just as before we stress that this result is meaningful since, by calculating the trace $\text{Tr}(X_{2|11} L_+^E) = 0$, we can also check that the Taub-NUT charge vanishes. Addressing the question whether there are subgroups of the original duality group in four-dimensions $\text{SL}(2, \mathbb{R})$ that leave the charge vector (5.34) invariant we realize that such a group contains only the identity

$$\text{SL}(2, \mathbb{R}) \supset \mathcal{S}_{2|11} = \mathbf{1} \quad (5.36)$$

Hence we clearly establish the intrinsic difference between the two type of small black holes at the level of the \mathbf{W} -representation. Both have vanishing quartic invariant, yet only the orbit 4|11 has a residual symmetry.

H*-stability subgroup Considering next the stability subgroup of the nilpotent element $X_{2|11}$ in $\mathbf{H}^* = \widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{\mathbf{h}^*}$ we obtain:

$$\mathfrak{S}_{2|11} = \text{SO}(1, 1) \triangleright \mathbb{R} \quad (5.37)$$

A generic element of the corresponding Lie algebra is a linear combination of two generators J, T , satisfying the commutation relations:

$$[J, T] = \frac{3}{2\sqrt{6}} T \quad (5.38)$$

We do not give its explicit form which we do not use in the sequel.

Nilpotent algebra $\mathbb{N}_{4|11}$ Considering next the adjoint action of the central element $h_{2|11}$ on the subspace \mathbb{K}^* we find that its eigenvalues are the following ones:

$$\text{Eigenvalues}_{4|11}^{\mathbb{K}^*} = \{-3, 3, -2, 2, -1, 1, 0, 0\} \quad (5.39)$$

Therefore the three eigenoperators A_3, A_2, A_1 corresponding to the positive eigenvalues 3, 2, 1, respectively, form the restriction to \mathbb{K}^* of a nilpotent algebra $\mathbb{N}_{2|11}$. In this case A_i do not all commute among themselves so that, differently from the previous case we have $\mathbb{N}_{4|11} \neq \mathbb{N}_{4|11} \cap \mathbb{K}^*$. In particular we find a new generator:

$$B \in \mathbb{H}^* \quad (5.40)$$

which completes a four-dimensional algebra with the following commutation relations:

$$0 = [A_3, A_2] = [A_1, A_3] \quad (5.41)$$

$$B = [A_2, A_1]$$

$$0 = [B, A_1]$$

$$0 = [B, A_2]$$

$$0 = [B, A_3] \quad (5.42)$$

As in the previous case, the structure of the nilpotent algebra implies that for the orbit \mathcal{O}_{11}^2 we have only three functions \mathfrak{h}_i^0 which will be harmonic and independent. This is so because $\mathcal{D}^2 \mathbb{N}_{2|11} = 0$ and $\mathcal{D} \mathbb{N}_{2|11} \cap \mathbb{K}^* = 0$.

Explicitly we set:

$$\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3) = \sum_{i=1}^3 \mathfrak{h}_i A_i =$$

$$\begin{pmatrix} -\mathfrak{h}_2 & \mathfrak{h}_1 - \mathfrak{h}_3 & \mathfrak{h}_2 & -\sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_3 & 0 & -3\mathfrak{h}_1 - \mathfrak{h}_3 & 0 \\ \mathfrak{h}_1 - \mathfrak{h}_3 & -2\mathfrak{h}_2 & \mathfrak{h}_3 - 3\mathfrak{h}_1 & -\sqrt{2}\mathfrak{h}_2 & \mathfrak{h}_1 + \mathfrak{h}_3 & 0 & -3\mathfrak{h}_1 - \mathfrak{h}_3 \\ -\mathfrak{h}_2 & 3\mathfrak{h}_1 - \mathfrak{h}_3 & \mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_3 & 0 & -\mathfrak{h}_1 - \mathfrak{h}_3 & 0 \\ \sqrt{2}\mathfrak{h}_1 + \sqrt{2}\mathfrak{h}_3 & \sqrt{2}\mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_3 & -\sqrt{2}\mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_1 + \sqrt{2}\mathfrak{h}_3 \\ 0 & -\mathfrak{h}_1 - \mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_3 & -\mathfrak{h}_2 & 3\mathfrak{h}_1 - \mathfrak{h}_3 & \mathfrak{h}_2 \\ -3\mathfrak{h}_1 - \mathfrak{h}_3 & 0 & \mathfrak{h}_1 + \mathfrak{h}_3 & \sqrt{2}\mathfrak{h}_2 & \mathfrak{h}_3 - 3\mathfrak{h}_1 & 2\mathfrak{h}_2 & \mathfrak{h}_1 - \mathfrak{h}_3 \\ 0 & -3\mathfrak{h}_1 - \mathfrak{h}_3 & 0 & -\sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_3 & -\mathfrak{h}_2 & \mathfrak{h}_1 - \mathfrak{h}_3 & \mathfrak{h}_2 \end{pmatrix} \quad (5.43)$$

Considering $\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3)$ as a Lax operator and calculating its Taub-NUT charge and electromagnetic charges we find:

$$\mathbf{n}_{TN} = -2(3\mathfrak{h}_1 + \mathfrak{h}_3) \quad ; \quad \mathcal{Q} = \left\{ -2\sqrt{3}\mathfrak{h}_2, 6\mathfrak{h}_1 - 2\mathfrak{h}_3, -2\sqrt{3}(\mathfrak{h}_1 + \mathfrak{h}_3), 0 \right\} \quad (5.44)$$

This implies that constructing the multi-centre solution with harmonic functions the condition $\mathfrak{h}_3 = -3\mathfrak{h}_1$ might be sufficient to annihilate the Taub-NUT current. We shall demonstrate that in this case the condition is slightly more complicated.

For later convenience let us change the normalization in the basis of harmonic functions as follows:

$$\mathfrak{h}_1^{(0)} = \frac{1}{4}\mathcal{H}_3 \quad ; \quad \mathfrak{h}_2^{(0)} = \frac{1}{2}(1 - \mathcal{H}_2) \quad ; \quad \mathfrak{h}_3^{(0)} = \frac{1}{4}\mathcal{H}_1 \quad (5.45)$$

Implementing the symmetric coset construction with:

$$\mathcal{Y}(\mathcal{H}_3, \mathcal{H}_2, \mathcal{H}_1) \equiv \exp \left[\mathfrak{H} \left(\frac{1}{4}\mathcal{H}_3, \frac{1}{2}(1 - \mathcal{H}_2), \frac{1}{4}\mathcal{H}_1 \right) \right] \quad (5.46)$$

calculating the upper triangular coset representative $\mathbb{L}(\mathcal{Y})$ according to equations (3.26) and extracting the σ -model scalar fields we obtain the answer which we list below:

$$\exp[-U] = \frac{1}{2} \sqrt{-\mathcal{H}_3^2 + (4\mathcal{H}_1^3 + 6\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 + \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2)} \quad (5.47)$$

$$\text{Im } z = \frac{\sqrt{-\mathcal{H}_3^2 + (4\mathcal{H}_1^3 + 6\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 + \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2)}}{2(\mathcal{H}_1^2 + \mathcal{H}_2)} \quad (5.48)$$

$$\text{Re } z = \frac{\mathcal{H}_3 - \mathcal{H}_2\mathcal{H}_1}{2(\mathcal{H}_1^2 + \mathcal{H}_2)} \quad (5.49)$$

$$Z^M = \begin{pmatrix} \frac{\sqrt{\frac{3}{2}}(\mathcal{H}_3^2 - 2\mathcal{H}_1(2\mathcal{H}_1^2 + 3\mathcal{H}_2 - 1)\mathcal{H}_3 + \mathcal{H}_2(-4\mathcal{H}_2^2 + (4 - 3\mathcal{H}_1^2)\mathcal{H}_2 + 2\mathcal{H}_1^2))}{\mathcal{H}_3^2 - 2(2\mathcal{H}_1^3 + 3\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 - \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2)} \\ \frac{\sqrt{2}(2\mathcal{H}_1^3 + 3\mathcal{H}_2\mathcal{H}_1 - \mathcal{H}_3)}{\sqrt{2}(\mathcal{H}_1\mathcal{H}_2^2 + \mathcal{H}_3(2\mathcal{H}_1^2 + \mathcal{H}_2))} \\ \frac{-\mathcal{H}_3^2 + (4\mathcal{H}_1^3 + 6\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 + \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2)}{\sqrt{6}(\mathcal{H}_1\mathcal{H}_2^2 + \mathcal{H}_3(2\mathcal{H}_1^2 + \mathcal{H}_2))} \\ \frac{\mathcal{H}_3^2 - 2(2\mathcal{H}_1^3 + 3\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 - \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2)}{4\mathcal{H}_3\mathcal{H}_1^3 + 3\mathcal{H}_2^2\mathcal{H}_1^2 + \mathcal{H}_3^2} \\ \frac{\sqrt{2}(-\mathcal{H}_3^2 + (4\mathcal{H}_1^3 + 6\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 + \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2))}{\sqrt{2}(-\mathcal{H}_3^2 + (4\mathcal{H}_1^3 + 6\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 + \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2))} \end{pmatrix} \quad (5.50)$$

$$a = \frac{\mathcal{H}_3(-6\mathcal{H}_1^2 - 3\mathcal{H}_2 + 1) - \mathcal{H}_1(3\mathcal{H}_2^2 + 3\mathcal{H}_2 + 2\mathcal{H}_1^2)}{\mathcal{H}_3^2 - 2(2\mathcal{H}_1^3 + 3\mathcal{H}_2\mathcal{H}_1)\mathcal{H}_3 - \mathcal{H}_2^2(3\mathcal{H}_1^2 + 4\mathcal{H}_2)} \quad (5.51)$$

The Taub-NUT current Given this explicit result we can turn to the explicit oxidation formulae described in section 2.2 and calculate the Taub-NUT current which is the integrand of eq.(2.41). We find:

$$j^{TN} = \frac{1}{2}(*\nabla\mathcal{H}_3 + 3(\mathcal{H}_2*\nabla\mathcal{H}_1 - \mathcal{H}_1*\nabla\mathcal{H}_2)) \quad (5.52)$$

Analyzing eq.(5.52) we see that there are just two possible solutions to the condition $j^{TN} = 0$:

case a) $\mathcal{H}_3 = \beta = \text{const}$; $\mathcal{H}_1 = 0$. With this condition we obtain:

$$\exp[-U] = \frac{1}{2} \sqrt{4\mathcal{H}_2^3 - \beta^2} \quad (5.53)$$

$$z = \frac{\beta + i\sqrt{4\mathcal{H}_2^3 - \beta^2}}{2\mathcal{H}_2} \quad (5.54)$$

$$j^{EM} = \star \nabla \begin{pmatrix} -\sqrt{\frac{3}{2}}\mathcal{H}_2 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (5.55)$$

case b) $\mathcal{H}_3 = \beta = \text{const}$; $\mathcal{H}_2 = 0$

$$\exp[-U] = \frac{1}{2} \sqrt{\beta (4\mathcal{H}_1^3 - \beta)} \quad (5.56)$$

$$z = \frac{\beta + i \sqrt{\beta (4\mathcal{H}_1^3 - \beta)}}{2\mathcal{H}_3^2} \quad (5.57)$$

$$j^{EM} = \begin{pmatrix} 0 \\ 0 \\ -\sqrt{\frac{3}{2}} \mathcal{H}_1 \\ 0 \end{pmatrix} \quad (5.58)$$

It might seem that these two solutions correspond to different types of black holes but this is not the case, as we now show. From the asymptotic flatness boundary condition we find that the value of β is fixed in terms of the value at infinity of the corresponding harmonic function $\mathcal{H}_{1,2}$, which of course must satisfy the necessary condition for reality of the solution $\mathcal{H}_{1,2}(\infty) \geq 1$:

$$\begin{cases} \beta = 2\sqrt{[\mathcal{H}_2(\infty)]^3 - 1} & \text{case a} \\ \beta = 2\left([\mathcal{H}_1(\infty)]^3 + \sqrt{[\mathcal{H}_1(\infty)]^6 - 1}\right) & \text{case b} \end{cases} \quad (5.59)$$

In the vicinity of a pole by means of the usual argument we obtain the following behavior of the warp factor:

$$|\vec{x} - \vec{x}_\alpha|^2 \exp[-U] \underset{\vec{x} \rightarrow \vec{x}_\alpha}{\sim} \begin{cases} \sqrt{b_2^3} \sqrt{|\vec{x} - \vec{x}_\alpha|} + \mathcal{O}(|\vec{x} - \vec{x}_\alpha|^{3/2}) & : \text{ case a} \\ \sqrt{\beta b_1^3} \sqrt{|\vec{x} - \vec{x}_\alpha|} + \mathcal{O}(|\vec{x} - \vec{x}_\alpha|^{3/2}) & : \text{ case b} \end{cases} \quad (5.60)$$

Hence in both cases the horizon area vanishes at all poles \vec{x}_α and the reality conditions are satisfied choosing the appropriate sign of $b_{1,2}$. The charge vector has the same structure for all black holes encompassed in the first or in the second solution, namely:

$$\mathcal{Q}_\alpha = \begin{cases} \left\{ -\sqrt{\frac{3}{2}} p_\alpha, 0, 0, 0 \right\} & : p_\alpha = b_2 \quad \text{for pole } \alpha \\ \left\{ 0, 0, -\sqrt{\frac{3}{2}} q_\alpha, 0 \right\} & : q_\alpha = b_1 \quad \text{for pole } \alpha \end{cases} \quad (5.61)$$

In both cases the quartic invariant \mathfrak{I}_4 is zero for all black holes in the solutions, yet one might still doubt whether the **W**-orbit for the two cases might be different. It is not so, since a direct calculation shows that the image in the $j = \frac{3}{2}$ representation $\Lambda[\mathfrak{A}]^4$ of the following $\text{SL}(2, \mathbb{R})$ element:

$$\mathfrak{A} = \begin{pmatrix} 0 & \frac{p}{q} \\ -\frac{q}{p} & 0 \end{pmatrix} \quad (5.62)$$

maps the charge vector $\mathcal{Q}_{[q]} = \{0, 0, -q, 0\}$, into the charge vector $\mathcal{Q}_{[p]} = \{p, 0, 0, 0\}$, namely we have $\Lambda[\mathfrak{A}] \mathcal{Q}_{[q]} = \mathcal{Q}_{[p]}$. Hence the two solutions we have here discussed simply give different representatives of the same **W**-orbit.

⁴See [28] for details, in particular eq.(3.13) of that reference for the explicit form of the spin $\frac{3}{2}$ matrices.

Summary Just as in the previous case for a multicenter solution associated with the \mathcal{O}_{11}^2 orbit all the black holes included in one solution are of the same type, namely small black holes with the same identical properties.

5.3 The Large BPS Black Holes of \mathcal{O}_{11}^3

Next let us consider the orbit \mathcal{O}_{11}^3 , which in the spherical symmetric case leads to BPS Black holes with a finite horizon area.

W-representation In order to better appreciate the structure of these solutions, let us slightly generalize our orbit representative, writing the following nilpotent matrix that depends on two parameters (p, q) to be interpreted later as the magnetic and the electric charge of the hole:

$$X_{3|11}(p, q) = \begin{pmatrix} q & 0 & 0 & -\frac{q}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & \frac{p+q}{2} & -\frac{p}{2} & 0 & \frac{q}{2} & 0 & 0 \\ 0 & \frac{p}{2} & \frac{q-p}{2} & 0 & 0 & -\frac{q}{2} & 0 \\ \frac{q}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & \frac{q}{\sqrt{2}} \\ 0 & -\frac{q}{2} & 0 & 0 & \frac{p-q}{2} & \frac{p}{2} & 0 \\ 0 & 0 & \frac{q}{2} & 0 & -\frac{p}{2} & \frac{1}{2}(-p-q) & 0 \\ 0 & 0 & 0 & -\frac{q}{\sqrt{2}} & 0 & 0 & -q \end{pmatrix} \quad (5.63)$$

The standard triple representative mentioned in eq.(4.5) is just the particular case $X_{3|11}(1, 1)$. Applying the same strategy as in the previous case, from the general formula we obtain

$$\mathcal{Q}_{3|11}^{\mathbf{w}} = \text{Tr}(X_{3|11}(p, q)\mathcal{T}^{\mathbf{w}}) = (0, p, -\sqrt{3}q, 0) \quad (5.64)$$

Substituting such a result in the expression for the quartic symplectic invariant (see eq.(5.3)) we find:

$$\mathfrak{I}_4 = 9pq^3 > 0 \quad \text{if } p \text{ and } q \text{ have the same sign} \quad (5.65)$$

Just as before we stress that this result is meaningful since, by calculating the trace $\text{Tr}(X_{3|11}L_+^E) = 0$, we can also check that the Taub-NUT charge vanishes. Furthermore we note that the condition that p and q have the same sign was singled out in [28] as the defining condition of the orbit \mathcal{O}_{11}^3 which, in the spherical symmetry approach leads to regular BPS solutions. The choice of opposite signs was proved in [28] to correspond to a different \mathbf{H}^* orbit, the non diagonal \mathcal{O}_{21}^3 which instead contains only singular solutions. Here we will show another important and intrinsically four dimensional reason to separate the two cases.

Addressing the question whether there are subgroups of the original duality group in four-dimensions $\text{SL}(2, \mathbb{R})$ that leave the charge vector (5.64) invariant we realize that such a subgroup exists and is the finite cyclic group of order three ⁵:

$$\text{SL}(2, \mathbb{R}) \supset \mathcal{S}_{3|11} = \mathbb{Z}_3 \quad (5.66)$$

⁵The authors express their gratitude to Alessio Marrani who attracted their attention, after the first appearance of the present paper in the arXive to paper [46] where the existence of a discrete stability subgroup for the charges of the regular BPS orbit in the S^3 model had already been found.

$\mathcal{S}_{3|11}$ is made by the following three elements:

$$\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (5.67)$$

$$\mathfrak{B} = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \sqrt{\frac{p}{q}} \\ \frac{\sqrt{3}}{2} \sqrt{\frac{q}{p}} & -\frac{1}{2} \end{pmatrix} \quad (5.68)$$

$$\mathfrak{B}^2 = \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \sqrt{\frac{p}{q}} \\ -\frac{\sqrt{3}}{2} \sqrt{\frac{q}{p}} & -\frac{1}{2} \end{pmatrix} ; \quad \mathfrak{B}^3 = \mathbf{1} \quad (5.69)$$

It is evident that such a \mathbb{Z}_3 subgroup exists if and only if the two charges p, q have the same sign. Otherwise the corresponding matrices develop imaginary elements and migrate to $\text{SL}(2, \mathbb{C})$. The existence of this isotropy group \mathbb{Z}_3 can be considered the very definition of the \mathbf{W} -orbit corresponding to BPS black holes. Indeed let us name $\lambda = \sqrt{\frac{p}{q}}$ and consider the algebraic condition imposed on a generic charge vector: $\mathcal{Q} = \{Q_1, Q_2, Q_3, Q_4\}$ by the request that it should admit the above described \mathbb{Z}_3 stability group:

$$\Lambda[\mathfrak{B}]\mathcal{Q} = \mathcal{Q} \Leftrightarrow \mathcal{Q} = \left(\sqrt{3}\lambda^2 Q_4, -\frac{\lambda^2 Q_3}{\sqrt{3}}, Q_3, Q_4 \right) \quad (5.70)$$

It is evident from the above explicit result that the charge vectors having this symmetry depend only on three parameters (λ^2, Q_3, Q_4) . The very relevant fact is that substituting this restricted charge vector in the general formula (5.3) for the quartic invariant we obtain:

$$\mathfrak{I}_4 = \lambda^2 (Q_3^2 + 3\lambda^2 Q_4^2)^2 > 0 \quad (5.71)$$

Hence the \mathbb{Z}_3 guarantees that the quartic invariant is a perfect square and hence positive. It is an intrinsic restriction characterizing the \mathbb{W} -orbit.

H*-stability subgroup Considering next the stability subgroup of the nilpotent element $X_{3|11}(1, 1)$ in $\mathcal{H}^* = \widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{\mathcal{H}^*}$ we obtain:

$$\mathfrak{S}_{3|11} = \mathbb{R} \quad (5.72)$$

the group being generated by a matrix $\mathbb{A}_{3|11}$ of nilpotency degree 2:

$$\mathbb{A}_{3|11}^2 = \mathbf{0} \quad (5.73)$$

We do not give its explicit form which we do not use in the sequel.

Nilpotent algebra $\mathbb{N}_{3|11}$ Considering next the adjoint action of the central element $h_{3|11}$ on the subspace \mathbb{K}^* we find that its eigenvalues are the following ones:

$$\text{Eigenvalues}_{3|11}^{\mathbb{K}^*} = \{-2, -2, -2, -2, 2, 2, 2, 2\} \quad (5.74)$$

Therefore the four eigenoperators A_1, A_2, A_3, A_4 corresponding to the four positive eigenvalues 2, respectively, form the restriction to \mathbb{K}^* of a nilpotent algebra $\mathbb{N}_{3|11}$. Also in this case the A_i do not all commute among themselves so that, we have $\mathbb{N}_{3|11} \neq \mathbb{N}_{3|11} \cap \mathbb{K}^*$. In particular we find a new generator:

$$B \in \mathbb{H}^* \quad (5.75)$$

which completes a five-dimensional algebra with the following commutation relations:

$$\begin{aligned}
[A_i, A_j] &= \Omega_{ij} B \\
[B, A_i] &= 0 \\
B &= \begin{pmatrix} 0 & 0 & -1 & 1 \\ 0 & 0 & -1 & -1 \\ 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \end{pmatrix}
\end{aligned} \tag{5.76}$$

The structure of the nilpotent algebra implies that for the orbit \mathcal{O}_{11}^3 we have only four functions \mathfrak{h}_i^0 which will be harmonic and independent. This is so because $\mathcal{D}^2 \mathbb{N}_{3|11} = 0$ and $\mathcal{DN}_{3|11} \cap \mathbb{K}^\star = 0$.

Explicitly we set:

$$\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3, \mathfrak{h}_4) = \sum_{i=1}^4 \mathfrak{h}_i A_i =$$

$$\begin{pmatrix} 2\mathfrak{h}_3 & \mathfrak{h}_1 - 2\mathfrak{h}_2 & 2\mathfrak{h}_1 - \mathfrak{h}_2 & -\sqrt{2}\mathfrak{h}_3 & -3\mathfrak{h}_2 & -3\mathfrak{h}_1 & 0 \\ \mathfrak{h}_1 - 2\mathfrak{h}_2 & \mathfrak{h}_3 - \mathfrak{h}_4 & \mathfrak{h}_4 & \sqrt{2}\mathfrak{h}_2 - 2\sqrt{2}\mathfrak{h}_1 & \mathfrak{h}_3 & 0 & -3\mathfrak{h}_1 \\ \mathfrak{h}_2 - 2\mathfrak{h}_1 & -\mathfrak{h}_4 & \mathfrak{h}_3 + \mathfrak{h}_4 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & 0 & -\mathfrak{h}_3 & -3\mathfrak{h}_2 \\ \sqrt{2}\mathfrak{h}_3 & 2\sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & 0 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_2 - 2\sqrt{2}\mathfrak{h}_1 & \sqrt{2}\mathfrak{h}_3 \\ 3\mathfrak{h}_2 & -\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & -\mathfrak{h}_3 - \mathfrak{h}_4 & -\mathfrak{h}_4 & 2\mathfrak{h}_1 - \mathfrak{h}_2 \\ -3\mathfrak{h}_1 & 0 & \mathfrak{h}_3 & 2\sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_2 & \mathfrak{h}_4 & \mathfrak{h}_4 - \mathfrak{h}_3 & \mathfrak{h}_1 - 2\mathfrak{h}_2 \\ 0 & -3\mathfrak{h}_1 & 3\mathfrak{h}_2 & -\sqrt{2}\mathfrak{h}_3 & \mathfrak{h}_2 - 2\mathfrak{h}_1 & \mathfrak{h}_1 - 2\mathfrak{h}_2 & -2\mathfrak{h}_3 \end{pmatrix} \tag{5.77}$$

Considering $\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3, \mathfrak{h}_4)$ as a Lax operator and calculating its Taub-NUT charge and electromagnetic charges we find:

$$\mathbf{n}_{TN} = -6\mathfrak{h}_1 \quad ; \quad \mathcal{Q} = \left\{ 2\sqrt{3}(\mathfrak{h}_2 - 2\mathfrak{h}_1), -2\mathfrak{h}_4, -2\sqrt{3}\mathfrak{h}_3, -6\mathfrak{h}_2 \right\} \tag{5.78}$$

This implies that constructing the multi-centre solution with harmonic functions the condition $\mathfrak{h}_1 = 0$ might be sufficient to annihilate the Taub-NUT current. We shall demonstrate that also in this case the condition is slightly more complicated. This emphasizes the difference between the Lax operator one-dimensional approach and the multicenter construction based on harmonic functions.

For later convenience let us change the normalization in the basis of harmonic functions as follows:

$$\mathfrak{h}_1^{(0)} = \frac{1}{\sqrt{12}}\mathcal{H}_1 \quad ; \quad \mathfrak{h}_2^{(0)} = \frac{1}{\sqrt{12}}\mathcal{H}_2 \quad ; \quad \mathfrak{h}_3^{(0)} = \frac{1}{2}(\mathcal{H}_3 - 1) \quad ; \quad \mathfrak{h}_4^{(0)} = \frac{1}{2}(\mathcal{H}_4 + 1) \tag{5.79}$$

Implementing the symmetric coset construction with:

$$\mathcal{Y}(\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3, \mathcal{H}_4) \equiv \exp \left[\mathfrak{H} \left(\frac{1}{\sqrt{12}}\mathcal{H}_1, \frac{1}{\sqrt{12}}\mathcal{H}_2, \frac{1}{2}(\mathcal{H}_3 - 1), \frac{1}{2}(\mathcal{H}_4 + 1) \right) \right] \tag{5.80}$$

calculating the upper triangular coset representative $\mathbb{L}(\mathcal{Y})$ according to equations (3.26) and extracting the σ -model scalar fields we obtain an explicit but rather messy answer which we present in the appendix in eq.s (A.1, A.5, A.7). In particular we obtain the Taub-NUT current in the following form:

$$j^{TN} = \sum_{i=1}^4 \mathfrak{R}_i(\mathcal{H}) \nabla \mathcal{H}_i \quad (5.81)$$

where $\mathfrak{R}_i(\mathcal{H})$ are rational functions of the four harmonic functions, the maximal degree of involved polynomials being 16. A priori, imposing the vanishing of the Taub-NUT current is a problem without guaranteed solutions. In the 4-dimensional linear space of the harmonic functions we can introduce r -linear relations of the form:

$$0 = V_\alpha^i \mathcal{H}_i \quad ; \quad \alpha = 1, \dots, r \quad (5.82)$$

Let U_a^i be a set of $4 - r$ linear independent 4-vectors orthogonal to the vectors V_α^i . Then it must happen that on the locus defined by eq.s (5.82), the following rational functions should also vanish

$$0 = \mathfrak{P}_a(\mathcal{H}) \equiv U_a^i \mathfrak{R}_i(\mathcal{H}) \quad ; \quad (a = 1, \dots, r - 4) \quad (5.83)$$

For generic rational functions this will never happen, yet we know that for our system such solutions should exist and in want of a clear cut algorithm it is a matter of ingenuity to find them. We do not find any solution with $r = 1$ but we find two nice solutions with $r = 2$. They are the following ones:

a) $\mathcal{H}_1 = \mathcal{H}_2 = 0$. The complete form of the supergravity solution corresponding to this choice is:

$$\exp[-U] = \sqrt{-\mathcal{H}_3^3 \mathcal{H}_4} \quad (5.84)$$

$$z = i \frac{\sqrt{-\mathcal{H}_3^3 \mathcal{H}_4}}{\mathcal{H}_3^2} \quad (5.85)$$

$$j^{TN} = 0 \quad (5.86)$$

$$j^{EM} = \star \nabla \begin{pmatrix} 0 \\ \frac{\mathcal{H}_4}{\sqrt{2}} \\ \sqrt{\frac{3}{2}} \mathcal{H}_3 \\ 0 \end{pmatrix} \quad (5.87)$$

b) $\mathcal{H}_1 = 0, \mathcal{H}_3 = -\mathcal{H}_4$. The complete form of the supergravity solution corresponding to this choice is:

$$\exp[-U] = \sqrt{-\frac{\mathcal{H}_2^4}{3} - 2\mathcal{H}_4^2 \mathcal{H}_2^2 + \mathcal{H}_4^4} \quad (5.88)$$

$$z = \frac{2\mathcal{H}_2 \mathcal{H}_4 - i \sqrt{-\mathcal{H}_2^4 - 6\mathcal{H}_4^2 \mathcal{H}_2^2 + 3\mathcal{H}_4^4}}{\sqrt{3} (\mathcal{H}_2^2 - \mathcal{H}_4^2)} \quad (5.89)$$

$$j^{TN} = 0 \quad (5.90)$$

$$j^{EM} = \star \nabla \begin{pmatrix} -\frac{\mathcal{H}_2}{\sqrt{2}} \\ \frac{\mathcal{H}_4}{\sqrt{2}} \\ -\sqrt{\frac{3}{2}} \mathcal{H}_4 \\ \sqrt{\frac{3}{2}} \mathcal{H}_2 \end{pmatrix} \quad (5.91)$$

We can now make some comments about the two solutions. First of all both in case a) and in case b) we have to fix the asymptotic value of the harmonic functions at spatial infinity $r = \infty$, in such a way as to obtain asymptotic flatness. This is quite easy and we do not dwell on it. Secondly we have to fix the parameters of the harmonic functions in such a way that the warp factor is always real on the whole physical range. These conditions are also easily spelled out:

$$\begin{aligned} a) \quad & -\mathcal{H}_3\mathcal{H}_4 > 0 \\ b) \quad & -\frac{\mathcal{H}_4^4}{3} - 2\mathcal{H}_4^2\mathcal{H}_2^2 + \mathcal{H}_4^4 > 0 \end{aligned} \quad (5.92)$$

and in a multicenter solution can be easily arranged adjusting the coefficients of each pole. Thirdly we can comment about the structure of the charge vector that we obtain at each pole:

$$\mathcal{H}_i \sim a_i + \frac{Q_i}{|x - x_\alpha|} \quad (5.93)$$

In case a) and b) we respectively obtain:

$$\mathcal{Q}_\alpha = \begin{pmatrix} 0 \\ \frac{Q_4}{\sqrt{2}} \\ \sqrt{\frac{3}{2}}Q_3 \\ 0 \end{pmatrix} \quad (5.94)$$

$$\mathcal{Q}_\alpha = \begin{pmatrix} -\frac{Q_2}{\sqrt{2}} \\ \frac{Q_4}{\sqrt{2}} \\ -\sqrt{\frac{3}{2}}Q_4 \\ \sqrt{\frac{3}{2}}Q_2 \end{pmatrix} \quad (5.95)$$

Comparing with eq.s (5.70,5.71) we see that in both cases the structure of these charges is that imposed by the \mathbb{Z}_3 invariance which characterizes BPS black holes. The necessary choice of signs in the case a)

$$\frac{Q_4}{Q_3} < 0 \quad (5.96)$$

is the same which is required by the reality of the warp factor. Hence in case b) all the black holes encompassed by the solution at each pole are finite area BPS black holes. In case a) the same is true for all the poles common to the harmonic function \mathcal{H}_3 and \mathcal{H}_4 : they are finite area BPS black holes. Yet we can envisage the situation where some poles of \mathcal{H}_3 are not shared by \mathcal{H}_4 and viceversa. In this case the pole of \mathcal{H}_4 defines a very small black hole, while the pole of \mathcal{H}_3 defines a small black hole. This is confirmed by the fact that a charge vector of type $\{0, p, 0, 0\}$ is mapped into $\{0, 0, 0, p\}$ by $\Lambda \left[\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right]$ and as such admits a parabolic subgroup of stability $\Lambda \left[\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \right]$.

Summary For a multicenter solution associated with the \mathcal{O}_{11}^3 orbit there are two possibilities either all the black holes included in one solution are regular, finite area, BPS black holes, either we have a mixture

of very small and small black holes. A finite area BPS black hole emerges when the center of a very small black hole coincides with the center of a small one. This provides the challenging suggestion that a BPS black hole can be considered quantum mechanically as a composite object where the "quarks" are small and very small black holes.

5.4 BPS Kerr-Newman solution

Next we want to show how this orbit encompasses also the BPS Kerr-Newman solution that was found by Luest et al in [45].

To this effect we go back to the general formulae (A.1-A.7) for the scalar fields in this orbit and we make the following reduction from four to two independent harmonic functions:

$$\mathcal{H}_2 = 0 \quad ; \quad \mathcal{H}_4 = -\frac{1}{3} \mathcal{H}_3 \quad (5.97)$$

With such a choice the expressions for all the scalar fields dramatically simplify and we obtain:

$$\mathfrak{W} = \frac{\sqrt{3}}{\mathcal{H}_1^2 + \mathcal{H}_3^2} \quad (5.98)$$

$$z = i \frac{1}{\sqrt{3}} \quad (5.99)$$

$$Z = \begin{pmatrix} -\frac{3\mathcal{H}_1}{\sqrt{2}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \\ \frac{\mathcal{H}_1^2 + (\mathcal{H}_3 - 3)\mathcal{H}_3}{\sqrt{2}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \\ -\frac{\sqrt{\frac{3}{2}}(\mathcal{H}_1^2 + (\mathcal{H}_3 - 1)\mathcal{H}_3)}{\mathcal{H}_1^2 + \mathcal{H}_3^2} \\ -\frac{\mathcal{H}_1}{\sqrt{6}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \end{pmatrix} \quad (5.100)$$

$$a = \frac{5\mathcal{H}_1}{\sqrt{3}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \quad (5.101)$$

Utilizing the above expressions in the final oxidation formulae we obtain the following result for the Taub-Nut current and for the electromagnetic currents:

$$j^{TN} = \frac{2(\star \nabla \mathcal{H}_1 \mathcal{H}_3 - \star \nabla \mathcal{H}_3 \mathcal{H}_1)}{\sqrt{3}} \quad (5.102)$$

$$j^{EM} = \begin{pmatrix} \frac{2\star \nabla \mathcal{H}_3 \mathcal{H}_1(\mathcal{H}_1^2 + (\mathcal{H}_3 - 2)\mathcal{H}_3) - \star \nabla \mathcal{H}_1((2\mathcal{H}_3 + 1)\mathcal{H}_1^2 + \mathcal{H}_3^2(2\mathcal{H}_3 - 3))}{\sqrt{2}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \\ \frac{\star \nabla \mathcal{H}_3(3\mathcal{H}_1^2 - \mathcal{H}_3^2) - 4\star \nabla \mathcal{H}_1 \mathcal{H}_1 \mathcal{H}_3}{3\sqrt{2}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \\ \frac{\sqrt{\frac{3}{2}}(4\star \nabla \mathcal{H}_1 \mathcal{H}_1 \mathcal{H}_3 + \star \nabla \mathcal{H}_3(\mathcal{H}_3^2 - 3\mathcal{H}_1^2))}{\mathcal{H}_1^2 + \mathcal{H}_3^2} \\ \frac{2\star \nabla \mathcal{H}_3 \mathcal{H}_1(\mathcal{H}_1^2 + (\mathcal{H}_3 - 6)\mathcal{H}_3) - \star \nabla \mathcal{H}_1((2\mathcal{H}_3 + 3)\mathcal{H}_1^2 + \mathcal{H}_3^2(2\mathcal{H}_3 - 9))}{\sqrt{6}(\mathcal{H}_1^2 + \mathcal{H}_3^2)} \end{pmatrix} \quad (5.103)$$

Next identifying the two harmonic functions with those introduced in eq.s(3.53-3.56), according to:

$$\mathcal{H}_1 = 3^{\frac{1}{4}} (1 + m \mathcal{P}) \quad ; \quad \mathcal{H}_3 = 3^{\frac{1}{4}} m \mathcal{R} \quad (5.104)$$

we obtain the following result for the warp-factor:

$$\exp[U] = \frac{(m+r)^2 + \alpha^2 \cos^2(\theta)}{r^2 + \alpha^2 \cos^2(\theta)} \quad (5.105)$$

and for the Kaluza-Klein vector:

$$\mathbf{A}^{[KK]} = \omega \equiv \frac{m(m+2r)\alpha \sin^2(\theta)}{r^2 + \alpha^2 \cos^2(\theta)} d\phi \quad (5.106)$$

Indeed one can easily check that, in the spheroidal coordinates (3.51) with flat metric 3.52 we have:

$$2m (\star \nabla \mathcal{P} \mathcal{R} - \mathcal{P} \star \nabla \mathcal{R}) = d\omega \quad (5.107)$$

where $\star \nabla$ denotes the Hodge dual of the exterior derivative d . Writing the corresponding final form of the metric::

$$ds_{BPSKN}^2 = -\exp[U] (dt + \omega)^2 + \exp[-U] d\Omega_{spheroidal}^2 \quad (5.108)$$

we can easily check that it is just the Kerr-Newman metric (2.54) with $q = m$. The only necessary step, in order to verify such an identity is a redefinition of the coordinate r . If in the metric (2.54) one replaces $r \rightarrow r + m$, then (2.54) becomes identical to (5.108).

It is interesting to consider the expressions for the vector field strengths that solve the Maxwell-Einstein system together with the BPS Kerr-Newmann metric. For the first two field strengths (magnetic), from eq.(5.103) we find:

$$\begin{aligned} F^1 = & -\frac{1}{\sqrt{2}(r^2 + \alpha^2 \cos^2 \theta)^2 ((m+r)^2 + \alpha^2 \cos^2 \theta)} \left(\sqrt[4]{3} m \alpha \sin \theta \left(\left((-3 + 2\sqrt[4]{3}) \alpha^4 \cos^4 \theta \right. \right. \right. \\ & + m \left(2\sqrt[4]{3} m + m + 2 \left(1 + \sqrt[4]{3} \right) r \right) \alpha^2 \cos^2 \theta - r(m+r)^2 \left(2\sqrt[4]{3} m + (-3 + 2\sqrt[4]{3}) r \right) \Big) \sin \theta dr \wedge d\phi \\ & + 2(r^2 + \alpha^2) \cos \theta \left(\left((-2 + \sqrt[4]{3}) m + (-3 + 2\sqrt[4]{3}) r \right) \alpha^2 \cos^2 \theta + (m+r) \left(\sqrt[4]{3} m^2 \right. \right. \\ & \left. \left. + (-1 + 3\sqrt[4]{3}) rm + (-3 + 2\sqrt[4]{3}) r^2 \right) \right) d\theta \wedge d\phi \Big) \end{aligned} \quad (5.109)$$

$$\begin{aligned} F^2 = & \frac{1}{\sqrt{2} 3^{3/4} (r^2 + \alpha^2 \cos^2 \theta)^2 ((m+r)^2 + \alpha^2 \cos^2 \theta)} \left(m \sin \theta \left(\alpha^2 (-2 \cos \theta \sin \theta r^3 \right. \right. \\ & + m^2 \sin 2\theta r - 2(2m+r) \alpha^2 \cos^3 \theta \sin \theta) dr \wedge d\phi \\ & - \frac{1}{8} (r^2 + \alpha^2) (8r^4 + 16mr^3 + 8m^2 r^2 + \alpha^4 \\ & \left. - 8\alpha^2 (-3m^2 - 6rm + \alpha^2) \cos^2 \theta - \alpha^4 \cos(4\theta)) d\theta \wedge d\phi \right) \end{aligned} \quad (5.110)$$

while for the second two we get:

$$\begin{aligned} G^3 = & \frac{1}{\sqrt{2} (r^2 + \alpha^2 \cos^2 \theta)^2 ((m+r)^2 + \alpha^2 \cos^2 \theta)} \left(3^{3/4} m \sin \theta \left((\sin 2\theta r^3 - 2m^2 \cos \theta \sin \theta r \right. \right. \\ & + 2(2m+r) \alpha^2 \cos^3 \theta \sin \theta) dr \wedge d\phi \alpha^2 \\ & \left. + \frac{1}{8} (r^2 + \alpha^2) (8r^4 + 16mr^3 + 8m^2 r^2 + \alpha^4 - 8\alpha^2 (-3m^2 - 6rm + \alpha^2) \cos^2 \theta - \alpha^4 \cos(4\theta)) d\theta \wedge d\phi \right) \end{aligned} \quad (5.111)$$

$$\begin{aligned}
G^4 = & -\frac{1}{\sqrt{2}(r^2 + \alpha^2 \cos^2 \theta)^2 ((m+r)^2 + \alpha^2 \cos^2 \theta)} \left(m\alpha \sin \theta \left(\left(-2 + 3^{3/4} \right) \alpha^4 \cos^4 \theta \right. \right. \\
& + m \left(\left(2 + 3^{3/4} \right) m + 2 \left(1 + 3^{3/4} \right) r \right) \alpha^2 \cos^2 \theta + r(m+r)^2 \left(\left(-2 + 3^{3/4} \right) r - 2m \right) \left. \right) \sin \theta dr \wedge d\phi \\
& - 2(r^2 + \alpha^2) \cos \theta \left(-m^3 + \left(-4 + 3^{3/4} \right) rm^2 + \left(-5 + 4^{3/4} \right) r^2 m + \left(-2 + 3^{3/4} \right) r^3 \right. \\
& \left. \left. + \left(\left(-1 + 2^{3/4} \right) m + \left(-2 + 3^{3/4} \right) r \right) \alpha^2 \cos^2 \theta \right) d\theta \wedge d\phi \right) \quad (5.112)
\end{aligned}$$

The above expressions are rather formidable, yet considering them in some limit their meaning can be decoded. First of all we recall that in the limit $\alpha \rightarrow 0$ the metric (5.108) becomes the Reissner-Nordstrom metric. Correspondingly in the same limit the above four-vector of field strengths degenerates into:

$$\begin{pmatrix} F^1 \\ F^2 \\ G^3 \\ G^4 \end{pmatrix} \xrightarrow{\alpha \rightarrow 0} \begin{pmatrix} 0 \\ -\frac{m \sin(\theta) d\theta \wedge d\phi}{\sqrt{2} 3^{3/4}} \\ \frac{3^{3/4} m \sin(\theta) d\theta \wedge d\phi}{\sqrt{2}} \\ 0 \end{pmatrix} \quad (5.113)$$

showing that the black hole charges $\left(0, -\frac{m}{\sqrt{2} 3^{1/4}}, \frac{m 3^{1/4}}{\sqrt{2}}, 0\right)$ have the correct form for a BPS black hole and are endowed with the characteristic \mathbb{Z}_3 symmetry.

Also in the $\alpha \neq 0$ we can easily determine the black hole charges by integrating the field strengths on a two-sphere of very large radius $r \rightarrow \infty$. For this purpose it is important to evaluate the asymptotic expansion of the field strengths for large radius. We find:

$$\begin{pmatrix} F^1 \\ F^2 \\ G^3 \\ G^4 \end{pmatrix} \underset{r \rightarrow \infty}{\simeq} \begin{pmatrix} -\frac{\sqrt{2} 4^{3/4} (-3 + 2 \sqrt{3}) m \alpha \cos \theta \sin \theta d\theta \wedge d\phi}{r} + \mathcal{O}\left(\frac{1}{r^2}\right) \\ -\frac{m \sin \theta d\theta \wedge d\phi}{\sqrt{2} 3^{3/4}} + \mathcal{O}\left(\frac{1}{r^2}\right) \\ \frac{3^{3/4} m \sin \theta d\theta \wedge d\phi}{\sqrt{2}} + \mathcal{O}\left(\frac{1}{r^2}\right) \\ \frac{\sqrt{2} (-2 + 3^{3/4}) m \alpha \cos \theta \sin \theta d\theta \wedge d\phi}{r} + \mathcal{O}\left(\frac{1}{r^2}\right) \end{pmatrix} \quad (5.114)$$

and the integration on the angular variables produces the same result as for the corresponding Reissner-Nordstrom black hole:

$$\mathcal{Q}_{BPSKN} = \left(0, -\frac{m}{\sqrt{2} 3^{1/4}}, \frac{m 3^{1/4}}{\sqrt{2}}, 0\right) \quad (5.115)$$

In conclusion the BPS Kerr-Newman solution is a deformation of the Reissner-Nordstrom BPS black hole. It is extremal in the σ -model sense and for this reason could be retrieved from the nilpotent orbit construction. However it is not extremal in the sense of General Relativity since the mass is less than $\sqrt{q^2 + \alpha^2}$ being equal to m . For this reason we are below the limit of the cosmic censorship, there is no horizon and we have instead a naked singularity.

The important message is that, notwithstanding the deformation and the presence of a Kaluza-Klein vector, the structure of the charges is that pertaining to the orbit where the solution has been constructed, namely the BPS orbit \mathcal{O}_{11}^3 .

5.5 The Large non BPS Black Holes of \mathcal{O}_{22}^3

Next let us consider the orbit \mathcal{O}_{22}^3 , which in the spherical symmetric case leads to non BPS Black holes with a finite horizon area.

W-representation As in the previous case, in order to better appreciate the structure of these solutions, let us slightly generalize our orbit representative, writing the following nilpotent matrix that depends on two parameters (p, q)

$$X_{3|22}(p, q) = \begin{pmatrix} q & 0 & 0 & \frac{q}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & \frac{p+q}{2} & -\frac{p}{2} & 0 & -\frac{q}{2} & 0 & 0 \\ 0 & \frac{p}{2} & \frac{q-p}{2} & 0 & 0 & \frac{q}{2} & 0 \\ -\frac{q}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & -\frac{q}{\sqrt{2}} \\ 0 & \frac{q}{2} & 0 & 0 & \frac{p-q}{2} & \frac{p}{2} & 0 \\ 0 & 0 & -\frac{q}{2} & 0 & -\frac{p}{2} & \frac{1}{2}(-p-q) & 0 \\ 0 & 0 & 0 & \frac{q}{\sqrt{2}} & 0 & 0 & -q \end{pmatrix} \quad (5.116)$$

The standard triple representative mentioned in eq.(4.6) is just the particular case $X_{3|22}(1, 1)$. Applying the usual strategy from the general formula we obtain

$$\mathcal{Q}_{3|22}^{\mathbf{w}} = \text{Tr}(X_{3|22}(p, q)\mathcal{T}^{\mathbf{w}}) = (0, p, \sqrt{3}q, 0) \quad (5.117)$$

Substituting such a result in the expression for the quartic symplectic invariant (see eq.(5.3) we find:

$$\mathcal{I}_4 = -9pq^3 < 0 \quad \text{if } p \text{ and } q \text{ have the same sign} \quad (5.118)$$

This result is meaningful since, by calculating the trace $\text{Tr}(X_{3|22}L_+^E) = 0$, we find that the Taub-NUT charge vanishes. Furthermore we note that the condition that p and q have the same sign was singled out in [28] as the defining condition of the orbit O_{22}^3 which, in the spherical symmetry approach leads to regular non BPS solutions. The choice of opposite signs was proved in [28] to correspond to a different H^* orbit, the non diagonal O_{12}^3 which instead contains only singular solutions.

Addressing the question of stability subgroups of the original duality group in four-dimensions $\text{SL}(2, \mathbb{R})$, we realize that for the charge vector (5.117) this subgroup is just trivial:

$$\text{SL}(2, \mathbb{R}) \supset \mathcal{S}_{3|22} = \mathbf{1} \quad (5.119)$$

H^* -stability subgroup Considering next the stability subgroup of the nilpotent element $X_{3|22}(1, 1)$ in $H^* = \widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$ we obtain:

$$\mathfrak{S}_{3|22} = \mathbb{R} \quad (5.120)$$

the group being generated by a matrix $\mathbb{A}_{3|22}$ of nilpotency degree 2:

$$\mathbb{A}_{3|22}^3 = \mathbf{0} \quad (5.121)$$

We do not give its explicit form which we do not use in the sequel.

Nilpotent algebra $\mathbb{N}_{3|22}$ Considering next the adjoint action of the central element $h_{3|22}$ on the subspace \mathbb{K}^* we find that its eigenvalues are the following ones:

$$\text{Eigenvalues}_{3|22}^{\mathbb{K}^*} = \{-4, 4, -2, -2, 2, 2, 0, 0\} \quad (5.122)$$

Therefore the three eigenoperators A_1, A_2, A_3 corresponding to the three positive eigenvalues 4, 2, 2, respectively, form the restriction to \mathbb{K}^* of a nilpotent algebra $\mathbb{N}_{3|22}$. In this case the A_i do all commute among themselves so that we have $\mathbb{N}_{3|22} = \mathbb{N}_{3|22} \cap \mathbb{K}^*$ and it is abelian. The abelian structure of the nilpotent algebra implies that for the orbit \mathcal{O}_{22}^3 we have only three functions \mathfrak{h}_i^0 which will be harmonic and independent. This is so because $\mathcal{D}\mathbb{N}_{3|22} = 0$

Explicitly we set:

$$\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3) = \sum_{i=1}^3 \mathfrak{h}_i A_i =$$

$$\begin{pmatrix} 2\mathfrak{h}_3 & \mathfrak{h}_1 - 2\mathfrak{h}_2 & 2\mathfrak{h}_1 - \mathfrak{h}_2 & -\sqrt{2}\mathfrak{h}_3 & -3\mathfrak{h}_2 & -3\mathfrak{h}_1 & 0 \\ \mathfrak{h}_1 - 2\mathfrak{h}_2 & \mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_2 - 2\sqrt{2}\mathfrak{h}_1 & \mathfrak{h}_3 & 0 & -3\mathfrak{h}_1 \\ \mathfrak{h}_2 - 2\mathfrak{h}_1 & 0 & \mathfrak{h}_3 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & 0 & -\mathfrak{h}_3 & -3\mathfrak{h}_2 \\ \sqrt{2}\mathfrak{h}_3 & 2\sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & 0 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & \sqrt{2}\mathfrak{h}_2 - 2\sqrt{2}\mathfrak{h}_1 & \sqrt{2}\mathfrak{h}_3 \\ 3\mathfrak{h}_2 & -\mathfrak{h}_3 & 0 & \sqrt{2}\mathfrak{h}_1 - 2\sqrt{2}\mathfrak{h}_2 & -\mathfrak{h}_3 & 0 & 2\mathfrak{h}_1 - \mathfrak{h}_2 \\ -3\mathfrak{h}_1 & 0 & \mathfrak{h}_3 & 2\sqrt{2}\mathfrak{h}_1 - \sqrt{2}\mathfrak{h}_2 & 0 & -\mathfrak{h}_3 & \mathfrak{h}_1 - 2\mathfrak{h}_2 \\ 0 & -3\mathfrak{h}_1 & 3\mathfrak{h}_2 & -\sqrt{2}\mathfrak{h}_3 & \mathfrak{h}_2 - 2\mathfrak{h}_1 & \mathfrak{h}_1 - 2\mathfrak{h}_2 & -2\mathfrak{h}_3 \end{pmatrix} \quad (5.123)$$

Considering $\mathfrak{H}(\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_3)$ as a Lax operator and calculating its Taub-NUT charge and electromagnetic charges we find:

$$\mathbf{n}_{TN} = -6\mathfrak{h}_1 \quad ; \quad \mathcal{Q} = \left\{ 2\sqrt{3}(\mathfrak{h}_2 - 2\mathfrak{h}_1), 0, -2\sqrt{3}\mathfrak{h}_3, -6\mathfrak{h}_2 \right\} \quad (5.124)$$

This implies that constructing the multi-centre solution with harmonic functions the condition $\mathfrak{h}_1 = 0$ might be sufficient to annihilate the Taub-NUT current. In this case we will be lucky and such a condition suffices.

For later convenience let us change the normalization in the basis of harmonic functions as follows:

$$\mathfrak{h}_1^{(0)} = \mathcal{H}_1 \quad ; \quad \mathfrak{h}_2^{(0)} = \frac{1}{2}(1 - \mathcal{H}_2) \quad ; \quad \mathfrak{h}_3^{(0)} = \frac{1}{2}(1 - \mathcal{H}_3) \quad (5.125)$$

Implementing the symmetric coset construction with:

$$\mathcal{Y}(\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3) \equiv \exp \left[\mathfrak{H} \left(\mathcal{H}_1, \frac{1}{2}(1 - \mathcal{H}_2), \frac{1}{2}(1 - \mathcal{H}_3) \right) \right] \quad (5.126)$$

calculating the upper triangular coset representative $\mathbb{L}(\mathcal{Y})$ according to equations (3.26) and extracting the σ -model scalar fields we obtain an explicit expression which is sufficiently simple to be displayed:

$$\exp[-U] = \sqrt{\mathcal{H}_2 \mathcal{H}_3^3 - 4\mathcal{H}_1^2} \quad (5.127)$$

$$\text{Im } z = \frac{\sqrt{\mathcal{H}_2 \mathcal{H}_3^3 - 4\mathcal{H}_1^2}}{\mathcal{H}_3^2} \quad (5.128)$$

$$\text{Re } z = -\frac{2\mathcal{H}_1}{\mathcal{H}_3^2} \quad (5.129)$$

$$Z^M = \begin{pmatrix} -\frac{\sqrt{6}\mathcal{H}_1\mathcal{H}_3}{4\mathcal{H}_1^2-\mathcal{H}_2\mathcal{H}_3^3} \\ \frac{4\mathcal{H}_1^2-(\mathcal{H}_2-1)\mathcal{H}_3^3}{\sqrt{2}(4\mathcal{H}_1^2-\mathcal{H}_2\mathcal{H}_3^3)} \\ \frac{\sqrt{\frac{3}{2}}(4\mathcal{H}_1^2-\mathcal{H}_2(\mathcal{H}_3-1)\mathcal{H}_3^2)}{4\mathcal{H}_1^2-\mathcal{H}_2\mathcal{H}_3^3} \\ \frac{\sqrt{2}\mathcal{H}_1\mathcal{H}_2}{4\mathcal{H}_1^2-\mathcal{H}_2\mathcal{H}_3^3} \end{pmatrix} \quad (5.130)$$

$$a = -\frac{\mathcal{H}_1(\mathcal{H}_2+3\mathcal{H}_3-2)}{4\mathcal{H}_1^2-\mathcal{H}_2\mathcal{H}_3^3} \quad (5.131)$$

Using these results we easily obtain the Taub-NUT current in the following form:

$$j^{TN} = 2 \star \nabla \mathcal{H}_1 \quad (5.132)$$

In this case the predicted condition $\mathcal{H}_1 = 0$ is sufficient to annihilate the Taub-NUT current and we obtain an extremely simple result⁶. The complete form of the supergravity solution corresponding to this choice is:

$$\exp[-U] = \sqrt{\mathcal{H}_3^3\mathcal{H}_2} \quad (5.133)$$

$$z = i \frac{\sqrt{\mathcal{H}_3^3\mathcal{H}_2}}{\mathcal{H}_3^2} \quad (5.134)$$

$$j^{TN} = 0 \quad (5.135)$$

$$j^{EM} = \star \nabla \begin{pmatrix} 0 \\ -\frac{\mathcal{H}_2}{\sqrt{2}} \\ -\sqrt{\frac{3}{2}}\mathcal{H}_3 \\ 0 \end{pmatrix} \quad (5.136)$$

Comparing with the case of the large BPS orbit we see that the only difference is the relative sign of the harmonic functions in the electromagnetic current. What we said for the BPS black holes extends to the non BPS ones in the same way.

Summary For a multicenter solution associated with the \mathcal{O}_{22}^3 orbit we have a mixture of very small and small black holes as in the case of the orbit \mathcal{O}_{22}^3 . Also here a finite area non BPS black hole emerges when the center of a very small black comes to coincides with the center of a small one. The only difference is the relative sign of the two charges. With equal signs we construct a non BPS state, while with opposite charges we construct a BPS one. This reinforces the conjecture that at the quantum level finite black holes can be interpreted as composite states.

This conjecture is also supported by an angular momentum analysis. Looking at the representations in table 1, we see that the representation $2(j=1) + (j=0)$ that corresponds to BPS and non BPS large black holes can be obtained by summing the representation $(j=1) + 2(j=\frac{1}{2})$ that corresponds

⁶Actually even the condition $\mathcal{H}_1 = \text{const}$ suffices to annihilate the Taub-NUT charge allowing for a non trivial real part of the z -field. However in this section we analyze the case $\mathcal{H}_1 = 0$ for its remarkable simplicity

to small black holes with the representation $3(j=0) + 2(j=\frac{1}{2})$ that corresponds to very small black holes. Consider the following table:

1	$\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{1}{2}$	-1
0	$\frac{1}{2}$	$-\frac{1}{2}$	0	$\frac{1}{2}$	$-\frac{1}{2}$	0
1	1	0	0	0	-1	-1

the numbers in the first line are the eigenvalues of the central element h in the triplet (h, X, Y) characterizing the orbit \mathcal{O}_{11}^4 . The second line contains the eigenvalues for the central element of the triplet of the orbit \mathcal{O}_{11}^4 . In the last line we have the eigenvalues for the h in the triplet characterizing the orbit $\mathcal{O}_{i,j}^3$. We realize that the coincidence of centres correspond to the identification of a new $SL(2, \mathbb{R})$ subgroup which is the direct sum of the original two associated with the two small black holes.

5.6 The Largest orbit \mathcal{O}_{11}^1

Next let us consider the orbit \mathcal{O}_{11}^1 , which in the spherical symmetric case leads only to singular solutions.

W-representation Applying the usual strategy from the general formula we obtain a charge vector

$$\mathcal{Q}_{1|11}^{\mathbf{w}} = \text{Tr}(X_{1|11}(p, q)\mathcal{T}^{\mathbf{w}}) \quad (5.137)$$

which has no invariance:

$$SL(2, \mathbb{R}) \supset \mathcal{S}_{1|11} = \mathbf{1} \quad (5.138)$$

and yields a quartic invariant generically different from zero:

$$\mathfrak{I}_4 \neq 0 \quad (5.139)$$

Because of our simplified choice of the representative the Taub-NUT charge is not zero and only later we will enforce the vanishing of the Taub-NUT current on the harmonic function parameterized solution.

H^{*}-stability subgroup Considering next the stability subgroup of the nilpotent element $X_{1|11}$ in $H^* = \widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$ we obtain that it is trivial:

$$\mathfrak{S}_{1|11} = \mathbf{1} \quad (5.140)$$

Nilpotent algebra $\mathbb{N}_{1|11}$ Considering next the adjoint action of the central element $h_{1|11}$ on the subspace \mathbb{K}^* we find that its eigenvalues are the following ones:

$$\text{Eigenvalues}_{3|22}^{\mathbb{K}^*} = \{-5, 5, -3, 3, -1, -1, 1, 1\} \quad (5.141)$$

Therefore the four eigenoperators A_1, A_2, A_3, A_4 corresponding to the four positive eigenvalues 5, 3, 1, 1, respectively, form the restriction to \mathbb{K}^* of a nilpotent algebra $\mathbb{N}_{1|11}$. In this case the A_i do not all

commute among themselves so that we have $\mathbb{N}_{1|11} \neq \mathbb{N}_{1|11} \cap \mathbb{K}^*$. The full algebra involves also two operators $B_1, B_2 \in \mathbb{H}^*$ and the full set of commutation relations is the following one:

$$\begin{aligned}
0 &= [A_1, A_2] = [A_1, A_3] = [A_1, A_4] \\
0 &= [A_2, A_3] \\
0 &= [B_1, B_2] = [B_1, A_1] = [B_1, A_2] \\
0 &= [B_1, A_4] = [B_2, A_1] = [B_2, A_3] \\
B_1 &= [A_2, A_4] \\
B_2 &= [A_3, A_4] \\
-16 A_1 &= [B_1, A_3] \\
-16 A_1 &= [B_2, A_1] \\
24 A_2 &= [B_2, A_4]
\end{aligned} \tag{5.142}$$

By inspection of eq.s(5.142) we easily see that:

$$\mathcal{D}\mathbb{N}_{1|11} = \text{span}\{B_1, B_2, A_1, A_2\} \quad ; \quad \mathcal{D}\mathbb{N}_{1|11} \cap \mathbb{K}^* = \text{span}\{A_1, A_2\} \tag{5.143}$$

$$\mathcal{D}^2\mathbb{N}_{1|11} = \text{span}\{A_1\} = \mathcal{D}^2\mathbb{N}_{1|11} \cap \mathbb{K}^* \tag{5.144}$$

This structure of the nilpotent algebra implies that for the orbit \mathcal{O}_{11}^1 we have only two functions $\mathfrak{h}_3^0, \mathfrak{h}_4^0$ which are harmonic and independent. The other two functions $\mathfrak{h}_1^2, \mathfrak{h}_2^1$, obey instead equations in which the previous two play the role of sources. Not surprisingly $\mathfrak{h}_1^2, \mathfrak{h}_2^1$ correspond to the higher gradings 5 and 3, while $\mathfrak{h}_3^0, \mathfrak{h}_4^0$ correspond to the gradings 1, 1. More precisely \mathfrak{h}_2^1 receives source contributions only from $\mathfrak{h}_3^0, \mathfrak{h}_4^0$, while \mathfrak{h}_1^2 receives source contributions from $\mathfrak{h}_2^1, \mathfrak{h}_3^0, \mathfrak{h}_4^0$

Explicitly we set:

$$\mathfrak{H}(\mathfrak{h}_1, \dots, \mathfrak{h}_4) = \sum_{i=1}^4 \mathfrak{h}_i A_i = \begin{pmatrix} \mathfrak{h}_1 + \mathfrak{h}_4 & \frac{\mathfrak{h}_2}{3} - \mathfrak{h}_3 & \mathfrak{h}_4 & \frac{\sqrt{2}\mathfrak{h}_2}{3} - \sqrt{2}\mathfrak{h}_3 & -\mathfrak{h}_1 & -\mathfrak{h}_2 - \mathfrak{h}_3 & 0 \\ \frac{\mathfrak{h}_2}{3} - \mathfrak{h}_3 & 2\mathfrak{h}_4 & \mathfrak{h}_2 + \mathfrak{h}_3 & -\sqrt{2}\mathfrak{h}_4 & \mathfrak{h}_3 - \frac{\mathfrak{h}_2}{3} & 0 & -\mathfrak{h}_2 - \mathfrak{h}_3 \\ -\mathfrak{h}_4 & -\mathfrak{h}_2 - \mathfrak{h}_3 & \mathfrak{h}_1 - \mathfrak{h}_4 & \frac{\sqrt{2}\mathfrak{h}_2}{3} - \sqrt{2}\mathfrak{h}_3 & 0 & \frac{\mathfrak{h}_2}{3} - \mathfrak{h}_3 & -\mathfrak{h}_1 \\ \sqrt{2}\mathfrak{h}_3 - \frac{\sqrt{2}\mathfrak{h}_2}{3} & \sqrt{2}\mathfrak{h}_4 & \frac{\sqrt{2}\mathfrak{h}_2}{3} - \sqrt{2}\mathfrak{h}_3 & 0 & \frac{\sqrt{2}\mathfrak{h}_2}{3} - \sqrt{2}\mathfrak{h}_3 & -\sqrt{2}\mathfrak{h}_4 & \sqrt{2}\mathfrak{h}_3 - \frac{\sqrt{2}\mathfrak{h}_2}{3} \\ \mathfrak{h}_1 & \frac{\mathfrak{h}_2}{3} - \mathfrak{h}_3 & 0 & \frac{\sqrt{2}\mathfrak{h}_2}{3} - \sqrt{2}\mathfrak{h}_3 & \mathfrak{h}_4 - \mathfrak{h}_1 & -\mathfrak{h}_2 - \mathfrak{h}_3 & \mathfrak{h}_4 \\ -\mathfrak{h}_2 - \mathfrak{h}_3 & 0 & \mathfrak{h}_3 - \frac{\mathfrak{h}_2}{3} & \sqrt{2}\mathfrak{h}_4 & \mathfrak{h}_2 + \mathfrak{h}_3 & -2\mathfrak{h}_4 & \frac{\mathfrak{h}_2}{3} - \mathfrak{h}_3 \\ 0 & -\mathfrak{h}_2 - \mathfrak{h}_3 & \mathfrak{h}_1 & \frac{\sqrt{2}\mathfrak{h}_2}{3} - \sqrt{2}\mathfrak{h}_3 & -\mathfrak{h}_4 & \frac{\mathfrak{h}_2}{3} - \mathfrak{h}_3 & -\mathfrak{h}_1 - \mathfrak{h}_4 \end{pmatrix} \tag{5.145}$$

Considering $\mathfrak{H}(\mathfrak{h}_1, \dots, \mathfrak{h}_4)$ as a Lax operator and calculating its Taub-NUT charge and electromagnetic charges we find:

$$\mathbf{n}_{TN} = -2(\mathfrak{h}_2 + \mathfrak{h}_3) \quad ; \quad \mathcal{Q} = \left\{ -2\sqrt{3}\mathfrak{h}_4, -2(\mathfrak{h}_2 + \mathfrak{h}_3), \frac{2(\mathfrak{h}_2 - 3\mathfrak{h}_3)}{\sqrt{3}}, -2\mathfrak{h}_1 \right\} \tag{5.146}$$

This implies that constructing the multi-centre solution with harmonic functions the condition $\mathfrak{h}_2 = -\mathfrak{h}_3$ might be sufficient to annihilate the Taub-NUT current.

Implementing the symmetric coset construction with:

$$\mathcal{Y}(\mathfrak{h}_1, \dots, \mathfrak{h}_4) \equiv \exp[\mathfrak{H}(\mathfrak{h}_1, \dots, \mathfrak{h}_4)] \quad (5.147)$$

and imposing the field equations (3.14) we obtain the following conditions:

$$\begin{aligned} 0 &= \frac{224}{5} \nabla \mathfrak{h}_3 \circ \nabla \mathfrak{h}_3 \mathfrak{h}_4^3 - \frac{16}{5} \mathfrak{h}_3 \Delta \mathfrak{h}_3 \mathfrak{h}_4^3 - \frac{416}{5} \nabla \mathfrak{h}_3 \circ \nabla \mathfrak{h}_4 \mathfrak{h}_3 \mathfrak{h}_4^2 + \frac{16}{5} \mathfrak{h}_3^2 \Delta \mathfrak{h}_4 \mathfrak{h}_4^2 \\ &\quad + \frac{192}{5} \nabla \mathfrak{h}_4 \circ \nabla \mathfrak{h}_4 \mathfrak{h}_3^2 \mathfrak{h}_4 + \frac{32}{3} \nabla \mathfrak{h}_2 \circ \nabla \mathfrak{h}_3 \mathfrak{h}_4 - \frac{8}{3} \mathfrak{h}_3 \Delta \mathfrak{h}_2 \mathfrak{h}_4 - \frac{8}{3} \mathfrak{h}_2 \Delta \mathfrak{h}_3 \mathfrak{h}_4 \\ &\quad - \frac{16}{3} \nabla \mathfrak{h}_3 \circ \nabla \mathfrak{h}_4 \mathfrak{h}_2 - \frac{16}{3} \nabla \mathfrak{h}_2 \circ \nabla \mathfrak{h}_4 \mathfrak{h}_3 + \Delta \mathfrak{h}_1 + \frac{16}{3} \mathfrak{h}_2 \mathfrak{h}_3 \Delta \mathfrak{h}_4 \\ 0 &= 4 \Delta \mathfrak{h}_3 \mathfrak{h}_4^2 - 8 \nabla \mathfrak{h}_3 \circ \nabla \mathfrak{h}_4 \mathfrak{h}_4 - 4 \mathfrak{h}_3 \Delta \mathfrak{h}_4 \mathfrak{h}_4 + 8 \nabla \mathfrak{h}_4 \circ \nabla \mathfrak{h}_4 \mathfrak{h}_3 + \Delta \mathfrak{h}_2 \\ 0 &= \Delta \mathfrak{h}_3 \\ 0 &= \Delta \mathfrak{h}_4 \end{aligned} \quad (5.148)$$

Solutions of the above system can be quite complicated and can encompass many different types of behaviors, yet what is generically true is that the contributions from the source term introduces in \mathfrak{h}_1 and \mathfrak{h}_2 poles $1/r^p$ stronger than $p = 1$, while \mathfrak{h}_3 and \mathfrak{h}_4 have only simple poles. Hence if the structure of the polynomials in the functions $\mathfrak{h}_{1,2,3,4}$ is such that at simple poles the divergence of the inverse warp factor is already too strong or the coefficient already becomes imaginary, introducing stronger poles can only make the situation worse. For this reason we confine ourselves to analyze solutions encompassed in this orbit in which the source terms vanish identically upon the implementation of some identifications.

There are few different reductions with such a property and we choose just one that has also the additional feature of annihilating the Taub-NUT current. It is the following one:

$$\mathfrak{h}_3 = \mathfrak{h}_4 = -\mathfrak{h}_2 \equiv \mathfrak{h} \quad (5.149)$$

The reader can easily check that with the choice (5.149) the system of equations (5.148) reduces to:

$$\Delta \mathfrak{h} = \Delta \mathfrak{h}_1 = 0 \quad (5.150)$$

For later convenience let us change the normalization in the basis of harmonic functions as follows:

$$\mathfrak{h}_4 = \frac{1}{4} \mathcal{H} \quad ; \quad \mathfrak{h}_3 = \frac{1}{4} \mathcal{H} \quad ; \quad \mathfrak{h}_2 = -\frac{1}{4} \mathcal{H} \quad ; \quad \mathfrak{h}_1 = -\frac{1}{4} + \mathcal{W} \quad (5.151)$$

calculating the upper triangular coset representative $\mathbb{L}(\mathcal{Y})$ according to equations (3.26) and extracting the σ -model scalar fields we obtain explicit expressions which are sufficiently simple to be displayed:

$$\exp[U] = \frac{8\sqrt{15}}{\sqrt{-(\mathcal{H}+2)^3(\mathcal{H}^5+10\mathcal{H}^4+40\mathcal{H}^3+80\mathcal{H}^2-60(4\mathcal{W}+1))}} \quad (5.152)$$

$$\text{Im } z = \frac{3\sqrt{15}(\mathcal{H}+2)}{\sqrt{-\frac{\mathcal{H}+2}{\mathcal{H}^2(\mathcal{H}(\mathcal{H}+10)+40)+80}-60(4\mathcal{W}+1)}((\mathcal{H}(\mathcal{H}(\mathcal{H}+10)+20)-40)\mathcal{H}^2+90(4\mathcal{W}+1))} \quad (5.153)$$

$$\text{Re } z = \frac{15\mathcal{H}(\mathcal{H}+2)(\mathcal{H}+4)}{\mathcal{H}^5+10\mathcal{H}^4+20\mathcal{H}^3-40\mathcal{H}^2+360\mathcal{W}+90} \quad (5.154)$$

We skip the form of the Z fields and of a but we mention their consequences, namely the Taub-NUT current

$$j^{TN} = 0 \quad (5.155)$$

and the electromagnetic currents

$$j^{EM} = \star \nabla \left\{ \frac{1}{2} \sqrt{\frac{3}{2}} \mathcal{H}, 0, \frac{7\mathcal{H}}{6}, \sqrt{2}\mathcal{W} \right\} \quad (5.156)$$

This shows that a black hole belonging to this orbit has a charge vector $\mathcal{Q} = \left\{ \frac{1}{2} \sqrt{\frac{3}{2}} p, 0, \frac{7p}{6}, \sqrt{2}q \right\}$, whose quartic invariant is:

$$\mathfrak{I}_4 = \frac{1}{128} p^3 (49p + 72q) \quad (5.157)$$

This latter can be positive or negative depending on the choices for p and q . The problem, however, is that this solution is always singular around all poles of \mathcal{H} . Indeed setting:

$$\mathcal{H} \sim \frac{p}{r} \quad ; \quad \mathcal{W} \sim \frac{q}{r} \quad (5.158)$$

we find that for $r \rightarrow 0$ the inverse warp factor behaves as follows:

$$\exp[-U] \sim \frac{\sqrt{-p^8}}{8\sqrt{15}r^4} + \frac{\sqrt{-p^8}}{\sqrt{15}pr^3} + \frac{\sqrt{\frac{3}{5}}\sqrt{-p^8}}{p^2r^2} + \frac{4\sqrt{-p^8}}{\sqrt{15}p^3r} + \frac{\sqrt{\frac{3}{5}}p^3(p+5q)}{\sqrt{-p^8}} + \mathcal{O}(r) \quad (5.159)$$

The coefficient $\sqrt{-p^8}$ indicates that approaching the pole the warp factor becomes imaginary at a finite distance from it and the would be horizon $r = 0$ is never reached. If it were reached, the divergence $\frac{1}{r^4}$ would imply an infinite area of the horizon. As we know from our general discussion the Riemann tensor diverges if the warp factor goes to zero faster than r^2 so that the would be horizon would actually be a singularity. Yet since the warp factor becomes imaginary at a finite distance from the pole it remains open the question if solutions of this type can be prolonged by suitably changing the coordinate system. In that case they might acquire a physical meaning. So far such a question has not been tackled but it deserves to be.

6 Classification of the sugra-relevant symmetric spaces and discussion of their general properties

As we highlighted in the introduction there is a general group-theoretical framework underlying the construction of supergravity black holes which allows both for

- 1) a classification of the relevant symmetric spaces,
- 2) a general description of their structures which are relevant to the black hole solutions.

The presentation of both items in the above list is the goal of the present section. To achieve such a goal we need to emphasize a few general aspects of the decomposition (2.20) that relate to the underlying root systems and Dynkin diagrams. In the following we heavily rely on results presented several years

ago in [44]. Indeed from the algebraic view-point a crucial property of the general decomposition in eq.(2.20) is encoded into the following statements which are true for all the cases ⁷:

1. The A_1 root-system associated with the $\mathfrak{sl}(2, \mathbb{R})_E$ algebra in the decomposition (2.20) is made of $\pm \psi$ where ψ is the highest root of $\mathbb{U}_{D=3}$.
2. Out of the r simple roots α_i of $\mathbb{U}_{D=3}$ there are $r - 1$ that have grading zero with respect to ψ and just one α_W that has grading 1:

$$\begin{aligned} (\psi, \alpha_i) &= 0 & i \neq W \\ (\psi, \alpha_W) &= 1 \end{aligned} \quad (6.1)$$

3. The only simple root α_W that has non vanishing grading with respect ψ is just the highest weight of the symplectic representation \mathbf{W} of $\mathbb{U}_{D=4}$ to which the vector fields are assigned.
4. The Dynkin diagram of $\mathbb{U}_{D=4}$ is obtained from that of $\mathbb{U}_{D=3}$ by removing the dot corresponding to the special root α_W .
5. Hence we can arrange a basis for the simple roots of the rank r algebra $\mathbb{U}_{D=3}$ such that:

$$\begin{aligned} \alpha_i &= \{\bar{\alpha}_i, 0\} & i \neq W \\ \alpha_W &= \left\{ \bar{\mathbf{w}}_h, \frac{1}{\sqrt{2}} \right\} \\ \psi &= \{\mathbf{0}, \sqrt{2}\} \end{aligned} \quad (6.2)$$

where $\bar{\alpha}_i$ are $(r - 1)$ -component vectors representing a basis of simple roots for the Lie algebra $\mathbb{U}_{D=4}$, $\bar{\mathbf{w}}_h$ is also an $(r - 1)$ -vector representing the *highest weight* of the representation \mathbf{W} .

This means that the entire root system and the Cartan subalgebra of the $\mathbb{U}_{D=3}$ Lie algebra can be organized as follows:

$$\begin{array}{llll} \pm \psi & = & \pm (\mathbf{0}, \sqrt{2}) & ; & 2 \\ \pm \hat{\alpha} & = & \pm (\alpha, \sqrt{2}) & ; & 2 \times \# \text{ of roots} = 2n_r \\ \pm \hat{w} & = & \pm \left(w, \frac{\sqrt{2}}{2} \right) & ; & 2 \times \# \text{ of weights} = 2 \times \dim \mathbf{W} \\ \mathcal{H}^i \in \text{CSA} \subset \mathbb{U}_{D=4} & & & ; & \text{rank} \mathbb{U}_{D=4} = r \\ \mathcal{H}^\psi & & & & 1 \end{array} \quad (6.3)$$

$$\dim \mathbb{U}_{D=4} = 3 + \dim \mathbb{U}_{D=3} + 2 \times \dim \mathbf{W}$$

This organization of the Lie algebra is very important, as it was thoroughly discussed in [44], for the systematics of the Kač Moody extension which occurs when stepping down from D=3 to D=2 dimensions, but it is equally important in the present context to analyze the structure of the H^* -subalgebra and the Tits Satake projection.

⁷An apparent exception is given by the case of $\mathcal{N} = 3$ supergravity. The extra complicacy, there, is that the duality algebra in $D = 3$, namely $\mathbb{U}_{D=3}$ has rank $r + 2$, rather than $r + 1$ with respect to the rank of the algebra $\mathbb{U}_{D=4}$. Actually in this case there is an extra $U(1)_Z$ factor that is active on the vectors, but not on the scalars and which is responsible for the additional complications. It happens in this case that there are two vector roots, one for the complex representation to which the vectors are assigned and one for its conjugate. They have opposite charges under $U(1)_Z$. This case together with that of $\mathcal{N} = 5$ supergravity and with one of the series of $\mathcal{N} = 2$ theories completes the list of three *exotic models* which are anomalous also from the point of view of the Tits Satake projection (see below).

6.1 Tits Satake projection

In most cases of lower supersymmetry, neither the algebra $\mathbb{U}_{D=4}$ nor the algebra $\mathbb{U}_{D=3}$ are **maximally split**. In short this means that the non-compact rank $r_{nc} < r$ is less than the rank of \mathbb{U} , namely not all the Cartan generators are non-compact. Rigorously r_{nc} is defined as follows:

$$r_{nc} = \text{rank}(\mathbb{U}/\mathbb{H}) \equiv \dim \mathcal{H}^{n.c.} \quad ; \quad \mathcal{H}^{n.c.} \equiv \text{CSA}_{\mathbb{U}(\mathbb{C})} \cap \mathbb{K} \quad (6.4)$$

When this happens it means that, just as the billiard dynamics, also the structure of black hole solutions is effectively determined by a *maximally split subalgebra* $\mathbb{U}^{TS} \subset \mathbb{U}$ named the *Tits Satake* subalgebra of \mathbb{U} , whose rank is equal to r_{nc} . Effectively determined does not mean that solutions of the big system coincide with those of the smaller system rather it means that the former can be obtained from the latter by means of rotations of a compact subgroup of the big algebra $\mathbb{G}_{\text{paint}} \subset \mathbb{U}$ which we name the *paint group*, for whose precise definition we refer the reader to [36]. Here we just emphasize few important facts, relevant for our goals. To this effect we recall that the Tits Satake algebra is obtained from the original algebra via a projection of the root system of \mathbb{U} onto the subspace orthogonal to the compact part of the Cartan subalgebra of \mathbb{U}^{TS} :

$$\Pi^{TS} \quad ; \quad \Delta_{\mathbb{U}} \mapsto \overline{\Delta}_{\mathbb{U}^{TS}} \quad (6.5)$$

In euclidian geometry $\overline{\Delta}_{\mathbb{U}^{TS}}$ is just a collection of vectors in r_{nc} dimensions; a priori there is no reason why it should be the root system of another Lie algebra. Yet in almost all cases, $\overline{\Delta}_{\mathbb{U}^{TS}}$ turns out to be a Lie algebra root system and the maximal split Lie algebra corresponding to it, \mathbb{U}^{TS} , is, by definition, the Tits Satake subalgebra of the original non maximally split Lie algebra: $\mathbb{U}^{TS} \subset \mathbb{U}$. Such algebras \mathbb{U} are called *non-exotic*. The *exotic* non compact algebras are those for which the system $\overline{\Delta}_{\mathbb{U}^{TS}}$ is not an admissible root system. In such cases there is no Tits Satake subalgebra \mathbb{U}^{TS} . Exotic algebras are very few and in supergravity they appear only in three instances that display additional pathologies relevant also for the black hole solutions. For the non exotic models we have that the decomposition (2.20) commutes with the projection, namely:

$$\begin{aligned} \text{adj}(\mathbb{U}_{D=3}) &= \text{adj}(\mathbb{U}_{D=4}) \oplus \text{adj}(\mathfrak{sl}(2, \mathbb{R})_E) \oplus W_{(2,W)} \\ &\Downarrow \\ \text{adj}(\mathbb{U}_{D=3}^{TS}) &= \text{adj}(\mathbb{U}_{D=4}^{TS}) \oplus \text{adj}(\mathfrak{sl}(2, \mathbb{R})_E) \oplus W_{(2,W^{TS})} \end{aligned} \quad (6.6)$$

In other words the projection leaves the A_1 Ehlers subalgebra untouched and has a non trivial effect only on the duality algebra $\mathbb{U}_{D=4}$. Furthermore the image under the projection of the highest root of \mathbb{U} is the highest root of \mathbb{U}^{TS} :

$$\Pi^{TS} \quad : \quad \psi \rightarrow \psi^{TS} \quad (6.7)$$

The reason why the Tits Satake projection is relevant to us was pointed out in [29] where we advocated that the classification of nilpotent orbits and hence of extremal black hole solutions depends only on the Tits Satake subalgebra and therefore is universal for all members of the same Tits Satake universality class. By this name we mean all algebras who share the same Tits Satake projection.

Having clarified these points we can proceed with the classification of homogeneous symmetric spaces relevant to supergravity models and to black hole solutions.

#	TS D=4	TS D=3	coset D=4	coset D=3	Paint Group	subP Group	susy
1	$\frac{E_{7(7)}}{SU(8)}$	$\frac{E_{8(8)}}{SO^*(16)}$	$\frac{E_{7(7)}}{SU(8)}$	$\frac{E_{8(8)}}{SO^*(16)}$	1	1	$\mathcal{N} = 8$
2	$\frac{SU(1,1)}{U(1)}$	$\frac{G_{2(2)}}{SL(2,R) \times SL(2,R)}$	$\frac{SU(1,1)}{U(1)}$	$\frac{G_{2(2)}}{SL(2,R) \times SL(2,R)}$	1	1	$\mathcal{N} = 2$ n=1
3	$\frac{Sp(6,R)}{SU(3) \times U(1)}$	$\frac{F_{4(4)}}{Sp(6,R) \times SL(2,R)}$	$\frac{Sp(6,R)}{SU(3) \times U(1)}$	$\frac{F_{4(4)}}{Sp(6,R) \times SL(2,R)}$	1	1	$\mathcal{N} = 2$ n=6
4			$\frac{SU(3,3)}{SU(3) \times SU(3) \times U(1)}$	$\frac{E_{6(2)}}{SU(3,3) \times SL(2,R)}$	$SO(2) \times SO(2)$	1	$\mathcal{N} = 2$ n=9
5			$\frac{SO^*(12)}{SU(6) \times U(1)}$	$\frac{E_{7(-5)}}{SO^*(12) \times SL(2,R)}$	$SO(3) \times SO(3)$ $\times SO(3)$	$SO(3)_d$	$\mathcal{N} = 6$ $\mathcal{N} = 2$ n=16
6			$\frac{E_{7(-25)}}{E_{6(-78)} \times U(1)}$	$\frac{E_{8(-24)}}{E_{7(-25)} \times SL(2,R)}$	$SO(8)$	$G_{2(-14)}$	$\mathcal{N} = 2$ n=27
7	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,1)}{SO(2)}$	$\frac{SO(4,3)}{SO(2,2) \times SO(2,1)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,1)}{SO(6)}$	$\frac{SO(8,3)}{SO(6,2) \times SO(2,1)}$	$SO(5)$	$SO(4)$	$\mathcal{N} = 4$ n=1
8	$\frac{SL(2,R)}{O(2)} \times \frac{SO(3,2)}{SO(3) \times SO(2)}$	$\frac{SO(5,4)}{SO(3,2) \times SO(2,2)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,2)}{SO(6) \times SO(2)}$	$\frac{SO(8,4)}{SO(6,2) \times SO(2,2)}$	$SO(4)$	$SO(3)$	$\mathcal{N} = 4$ n=2
9	$\frac{SL(2,R)}{O(2)} \times \frac{SO(4,3)}{SO(4) \times SO(3)}$	$\frac{SO(6,5)}{SO(4,2) \times SO(2,3)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,3)}{SO(6) \times SO(3)}$	$\frac{SO(8,5)}{SO(6,2) \times SO(2,3)}$	$SO(3)$	$SO(2)$	$\mathcal{N} = 4$ n=3
10	$\frac{SL(2,R)}{O(2)} \times \frac{SO(5,4)}{SO(5) \times SO(4)}$	$\frac{SO(7,6)}{SO(5,2) \times SO(2,4)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,4)}{SO(6) \times SO(4)}$	$\frac{SO(8,6)}{SO(6,2) \times SO(2,4)}$	$SO(2)$	1	$\mathcal{N} = 4$ n=4
11	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,5)}{SO(6) \times SO(5)}$	$\frac{SO(8,7)}{SO(6,2) \times SO(2,5)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,5)}{SO(6) \times SO(5)}$	$\frac{SO(8,7)}{SO(6,2) \times SO(2,5)}$	1	1	$\mathcal{N} = 4$ n=5
12	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,6)}{SO(6) \times SO(6)}$	$\frac{SO(8,8)}{SO(6,2) \times SO(2,6)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,6)}{SO(6) \times SO(6)}$	$\frac{SO(8,8)}{SO(6,2) \times SO(2,6)}$	1	1	$\mathcal{N} = 4$ n=6
13	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,7)}{SO(6) \times SO(7)}$	$\frac{SO(8,9)}{SO(6,2) \times SO(2,7)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(6,6+p)}{SO(6) \times SO(6+p)}$	$\frac{SO(8,8+p)}{SO(6,2) \times SO(2,6+p)}$	$SO(p)$	$SO(p-1)$	$\mathcal{N} = 4$ n=6+p
14	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,1)}{SO(2)}$	$\frac{SO(4,3)}{SO(2,2) \times SO(2,1)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,1)}{SO(2)}$	$\frac{SO(4,3)}{SO(2,2) \times SO(2,1)}$	1	1	$\mathcal{N} = 2$ n=2
15	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,2)}{SO(2) \times SO(2)}$	$\frac{SO(4,4)}{SO(2,2) \times SO(2,2)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,2)}{SO(2) \times SO(2)}$	$\frac{SO(4,4)}{SO(2,2) \times SO(2,2)}$	1	1	$\mathcal{N} = 2$ n=3
16	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,3)}{SO(2) \times SO(3)}$	$\frac{SO(4,5)}{SO(2,2) \times SO(2,3)}$	$\frac{SL(2,R)}{O(2)} \times \frac{SO(2,2+p)}{SO(2) \times SO(2+p)}$	$\frac{SO(4,4+p)}{SO(2,2) \times SO(2,2+p)}$	$SO(p)$	$SO(p-1)$	$\mathcal{N} = 2$ n=3+p

Table 3: The 16 instances of *non-exotic* homogeneous symmetric scalar manifolds appearing in $D = 4$ supergravity. Non exotic means that the Tits Satake projection of the root system is a standard Lie Algebra root system. The 16 models are grouped according to their Tits Satake Universality classes. The time-like dimensional reduction is listed side by side. Within each class the models are distinguished by the different structure of the Paint Group and of its subPaint subgroup. The Paint group is the same in D=4 and in D=3

6.2 Classification of the sugra-relevant symmetric spaces

The classification of the symmetric coset based supergravity models is exhaustive and it is presented in tables 3 and 4. There are 16 universality classes of non-exotic models and 3 exceptional instances of exotic models which appear in the second table.

In the tables we have also listed the Paint groups and the subpaint groups. These latter are always compact and their different structures is what distinguishes the different elements belonging to the same class. As it was shown in [36], these groups are dimensional reduction invariant, namely they are the same in $D = 4$ and in $D = 3$. Hence the representation \mathbf{W} , which in particular contains the electromagnetic charges of the hole, can be decomposed with respect to the Tits Satake subalgebra and the Paint group revealing a regularity structure inside each Tits Satake universality class which is at the heart of the classification of *charge orbits*. The same decomposition can be given also for the \mathbb{K}^* representation and this is at the heart of the classification of black holes according to nilpotent orbits.

Focusing on the non-exotic models, we note that the 16 classes have a quite different type of population. There are six one element classes whose single member is maximally split. They are the following ones and all have a distinguished standpoint within the panorama of supergravity theories:

1. The $\mathcal{N} = 8$ supergravity theory, which is the maximal one in $D = 4$, (model 1).
2. The $\mathcal{N} = 2$ supergravity theory with a single vector multiplet and non-vanishing Yukawa (model 2).
3. The $\mathcal{N} = 4$ supergravity theory with 5 vector multiplets (model 11).
4. The $\mathcal{N} = 4$ supergravity theory with 6 vector multiplets which is obtained compactifying a type II theory on a T^6/\mathbb{Z}_2 orbifold (model 12).
5. The $\mathcal{N} = 2$ theory with two vector multiplets and non vanishing Yukawa couplings, usually called the *st*-model (model 14).
6. The $\mathcal{N} = 2$ theory with three vector multiplets and non vanishing Yukawa couplings, usually called the *stu*-model (model 15).

Next we have two universality classes, each containing an infinite number of elements. They are

1. The $\mathcal{N} = 4$ supergravity theory with $n = 6 + p$ vector multiplets ($p \geq 1$), (model 13).
2. The $\mathcal{N} = 2$ supergravity theory with $n = 3 + p$ vector multiplets ($p \geq 1$) and non vanishing Yukawa couplings (model 16).

We still have the very interesting 4-element universality class whose maximally split representative corresponds to the maximally split special Kähler manifold $\frac{\mathrm{Sp}(4, \mathbb{R})}{\mathrm{SU}(3) \times \mathrm{U}(1)}$. This class contains the models 3, 4, 5, 6 distinguished by quite peculiar Paint groups. We will thoroughly analyze the structure of this class.

Finally we have the three exotic models whose common feature is that their group and subgroup all belong to the pseudo-unitary series $\mathrm{SU}(p, q)$. The general decomposition (2.20) still holds true, but the Tits Satake projection loses its significance.

#	TS D=4	TS D=3	coset D=4	coset D=3	Paint Group	subP Group	susy
1_e	bc_1	bc_2	$\frac{SU(p+1,1)}{SU(p+1) \times U(1)}$	$\frac{SU(p+2,2)}{SU(p+1,1) \times SL(2, \mathbb{R})_{h^*}}$	$U(1) \times U(1) \times U(p)$	$U(p-1)$	$\mathcal{N} = 2$ $n=p+1$
2_e	bc_3	bc_4	$\frac{SU(p+1,3)}{SU(p+1) \times SU(3) \times U(1)}$	$\frac{SU(p+2,4)}{SU(p+1,2) \times SU(1,2) \times U(1)}$	$U(1) \times U(1) \times U(p)$	$U(p-1)$	$\mathcal{N} = 3$ $n=p+1$
3_e	bc_1	bc_2	$\frac{SU(5,1)}{SU(5) \times U(1)}$	$\frac{E_{6(-14)}}{SO^*(10) \times SO(2)}$	$U(1) \times U(1) \times U(4)$	$U(3)$	$\mathcal{N} = 5$

Table 4: The 3 instances of *exotic* homogenous symmetric scalar manifolds appearing in $D = 4$ supergravity. Exotic means that the Tits Satake projection of the root system is not a standard Lie Algebra root system. Notwithstanding this anomaly the concept of Paint Group, according to its definition as group of external automorphisms of the solvable Lie algebra generating the non compact coset manifold still exists. The Paint group is the same in D=4 and in D=3

6.3 Dynkin diagram analysis of the principal models

Next we analyze the form of the root systems of the $\mathbb{U}_{D=3}$ algebras in relation with the decomposition (2.20).

$\mathcal{N}=8$ This is the case of maximal supersymmetry and it is illustrated by fig. 1.


In this case all the involved Lie algebras are maximally split and we have

$$\text{adj } E_{8(8)} = \text{adj } E_{7(7)} \oplus \text{adj } SL(2, \mathbb{R})_E \oplus (\mathbf{2}, \mathbf{56}) \quad (6.8)$$

The highest root of $E_{8(8)}$ is

$$\psi = 3\alpha_1 + 4\alpha_2 + 5\alpha_3 + 6\alpha_4 + 3\alpha_5 + 4\alpha_6 + 2\alpha_7 + 2\alpha_8 \quad (6.9)$$

and the unique simple root not orthogonal to ψ is $\alpha_8 = \alpha_W$, according to the labeling of roots as in fig. 1. This root is the highest weight of the fundamental $\mathbf{56}$ -representation of $E_{7(7)}$.



$$\begin{aligned}\psi &= 3\alpha_1 + 4\alpha_2 + 5\alpha_3 + 6\alpha_4 + 3\alpha_5 + 4\alpha_6 + 2\alpha_7 + 2\alpha_8 \\ (\psi, \alpha_8) &= 1 \quad ; \quad (\psi, \alpha_i) = 0 \quad i \neq 8\end{aligned}$$

Figure 1: The Dynkin diagram of $E_{8(8)}$. The only simple root which has grading one with respect to the highest root ψ is α_8 (painted black). With respect to the algebra $\mathbb{U}_{D=4} = E_{7(7)}$ whose Dynkin diagram is obtained by removal of the black circle, α_8 is the highest weight of the symplectic representation of the vector fields, namely $\mathbf{W} = \mathbf{56}$.

The well adapted basis of simple E_8 roots is constructed as follows:

$$\begin{aligned}\alpha_1 &= \{1, -1, 0, 0, 0, 0, 0, 0\} &= \{\bar{\alpha}_1, 0\} \\ \alpha_2 &= \{0, 1, -1, 0, 0, 0, 0, 0\} &= \{\bar{\alpha}_2, 0\} \\ \alpha_3 &= \{0, 0, 1, -1, 0, 0, 0, 0\} &= \{\bar{\alpha}_3, 0\} \\ \alpha_4 &= \{0, 0, 0, 1, -1, 0, 0, 0\} &= \{\bar{\alpha}_4, 0\} \\ \alpha_5 &= \{0, 0, 0, 0, 1, -1, 0, 0\} &= \{\bar{\alpha}_5, 0\} \\ \alpha_6 &= \{0, 0, 0, 0, 1, 1, 0, 0\} &= \{\bar{\alpha}_6, 0\} \\ \alpha_7 &= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{\sqrt{2}}, 0\} &= \{\bar{\alpha}_7, 0\} \\ \alpha_8 &= \{-1, 0, 0, 0, 0, 0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\} &= \{\mathbf{w}_h, \frac{1}{\sqrt{2}}\}\end{aligned} \tag{6.10}$$

In this basis we recognize that the seven 7-vectors $\bar{\alpha}_i$ constitute a simple root basis for the E_7 root system, while:


$$\mathbf{w}_h = \{-1, 0, 0, 0, 0, 0, -\frac{1}{\sqrt{2}}\} \tag{6.11}$$

is the highest weight of the fundamental $\mathbf{56}$ dimensional representation. Finally in this basis the highest root ψ defined by eq.(6.9) takes the expected form:

$$\psi = \{0, 0, 0, 0, 0, 0, 0, \sqrt{2}\} \tag{6.12}$$

N=6 In this case the $D = 4$ duality algebra is $\mathbb{U}_{D=4} = \text{SO}^*(12)$, whose maximal compact subgroup is $H = \text{SU}(6) \times \text{U}(1)$. The scalar manifold:

$$\mathcal{SK}_{N=6} \equiv \frac{\text{SO}^*(12)}{\text{SU}(6) \times \text{U}(1)} \tag{6.13}$$



$$\begin{aligned}\psi &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 2\alpha_5 + 3\alpha_6 + 2\alpha_7 \\ (\psi, \alpha_7) &= 1 \quad ; \quad (\psi, \alpha_i) = 0 \quad i \neq 7\end{aligned}$$

Figure 2: The Dynkin diagram of $E_{7(-5)}$. The only simple root which has grading one with respect to the highest root ψ is α_4 (painted black). With respect to the algebra $\mathbb{U}_{D=4} = \text{SO}^*(12)$ whose Dynkin diagram is obtained by removal of the black circle, α_7 is the highest weight of the symplectic representation of the vector fields, namely the $\mathbf{W} = \mathbf{32}_s$.

is an instance of special Kähler manifold which can also be utilized in an $\mathcal{N} = 2$ supergravity context. The $D = 3$ algebra is $\mathbb{U}_{D=3} = E_{7(-5)}$. The 16 vector fields of $D = 4$ $\mathcal{N} = 6$ supergravity with their electric and magnetic field strengths fill the spinor representation $\mathbf{32}_s$ of $\text{SO}^*(12)$, so that the decomposition (2.20), in this case becomes:

$$\text{adj } E_{7(-5)} = \text{adj } \text{SO}^*(12) \oplus \text{adj } \text{SL}(2, \mathbb{R})_{\text{E}} \oplus (\mathbf{2}, \mathbf{32}_s) \quad (6.14)$$

The simple root α_W is α_7 and the highest root is:


$$\psi = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 2\alpha_5 + 3\alpha_6 + 2\alpha_7 \quad (6.15)$$

A well adapted basis of simple E_7 roots can be written as follows:

$$\begin{aligned}\alpha_1 &= \{1, -1, 0, 0, 0, 0, 0\} &= \{\bar{\alpha}_1, 0\} \\ \alpha_2 &= \{0, 1, -1, 0, 0, 0, 0\} &= \{\bar{\alpha}_2, 0\} \\ \alpha_3 &= \{0, 0, 1, -1, 0, 0, 0\} &= \{\bar{\alpha}_3, 0\} \\ \alpha_4 &= \{0, 0, 0, 1, -1, 0, 0\} &= \{\bar{\alpha}_4, 0\} \\ \alpha_5 &= \{0, 0, 0, 0, 1, -1, 0\} &= \{\bar{\alpha}_5, 0\} \\ \alpha_6 &= \{0, 0, 0, 0, 1, 1, 0\} &= \{\bar{\alpha}_6, 0\} \\ \alpha_7 &= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{\sqrt{2}}\} &= \{\bar{\mathbf{w}}_h, \frac{1}{\sqrt{2}}\}\end{aligned} \quad (6.16)$$

In this basis we recognize that the six 6-vectors $\bar{\alpha}_i$ ($i = 1, \dots, 6$) constitute a simple root basis for the $D_6 \simeq \text{SO}^*(12)$ root system, while:

$$\mathbf{w}_h = \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\} \quad (6.17)$$



$$\begin{aligned}\psi &= 2\varpi_1 + 3\varpi_2 + 4\varpi_3 + 2\varpi_4 \\ (\psi, \varpi_1) &= 2 \quad ; \quad (\psi, \varpi_i) = 0 \quad i \neq 1\end{aligned}$$

Figure 3: The Dynkin diagram of $F_{4(4)}$. The only root which is not orthogonal to the highest root is $\varpi_V = \varpi_1$. In the Tits Satake projection Π^{TS} the highest root ψ of $F_{4(4)}$ is the image of the highest root of $E_{7(-5)}$ and the root $\varpi_V = \varpi_1 = \Pi^{TS}(\alpha_7)$ is the image of the root associated with the vector fields.

is the highest weight of the spinor **32**-dimensional representation of $\text{SO}^*(12)$. Finally in this basis the highest root ψ defined by eq.(6.15) takes the expected form:

$$\psi = \{0, 0, 0, 0, 0, 0, \sqrt{2}\} \quad (6.18)$$

In this case, as in most cases of lower supersymmetry, neither the algebra $\mathbb{U}_{D=4}$ nor the algebra $\mathbb{U}_{D=3}$ are **maximally split**. The Tits Satake projection of $E_{7(-5)}$ is $F_{4(4)}$ and the explicit form of eq.(6.6) is the following one:

$$\begin{aligned}\text{adj}(E_{7(-5)}) &= \text{adj}(\text{SO}^*(12)) \oplus \text{adj}(\text{SL}(2, \mathbb{R})_E) \oplus (\mathbf{2}, \mathbf{32}_s) \\ &\Downarrow \\ \text{adj}(F_{4(4)}) &= \text{adj}(\text{Sp}(6, \mathbb{R}) \oplus \text{adj}(\text{SL}(2, \mathbb{R})_E) \oplus (\mathbf{2}, \mathbf{14}')\end{aligned} \quad (6.19)$$

The representation $\mathbf{14}'$ of $\text{Sp}(6, \mathbb{R})$ is that of an antisymmetric symplectic traceless tensor:

$$\dim_{\text{Sp}(6, \mathbb{R})} \begin{array}{c} \widetilde{\square} \\ \square \\ \square \end{array} = \mathbf{14}' \quad (6.20)$$


The Dynkin diagram of the Tits Satake subalgebra $\mathfrak{f}_{4(4)}$ is discussed in fig.3.

N=5 The case of $\mathcal{N} = 5$ supergravity is described by fig.4 and it is one of the three exotic models whose Tits - Satake projection does not produce a Lie algebra root system.

In the $\mathcal{N} = 5$ theory the scalar manifold is a complex coset of rank $r = 1$,

$$\mathcal{M}_{N=5, D=4} = \frac{\text{SU}(1, 5)}{\text{SU}(5) \times \text{U}(1)} \quad (6.21)$$

$E_{6(-14)}$



$$\begin{aligned}\psi &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 \\ (\psi, \alpha_4) &= 1 \quad ; \quad (\psi, \alpha_i) = 0 \quad i \neq 4\end{aligned}$$

Figure 4: The Dynkin diagram of $E_{6(-14)}$. The only simple root which has grading one with respect to the highest root ψ is α_7 (painted black). With respect to the algebra $\mathbb{U}_{D=4} = \text{SU}(5, 1)$ whose Dynkin diagram is obtained by removal of the black circle, α_4 is the highest weight of the symplectic representation of the vector fields, namely the $\mathbf{W} = \mathbf{20}$.

and there are $\mathbf{10}$ vector fields whose electric and magnetic field strengths are assigned to the $\mathbf{20}$ -dimensional representation of $\text{SU}(1, 5)$, which is that of an antisymmetric three-index tensor

$$\dim_{\text{SU}(1,5)} \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} = 20 \quad (6.22)$$

The decomposition (2.20) takes the explicit form:

$$\text{adj}(E_{6(-14)}) = \text{adj}(\text{SU}(1, 5) \oplus \text{adj}(\text{SL}(2, \mathbb{R})_E) \oplus (\mathbf{2}, \mathbf{20}) \quad (6.23)$$

and we have that the highest root of E_6 , namely

$$\psi = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6 \quad (6.24)$$

has non vanishing scalar product only with the root α_4 in the form depicted in fig.4.

Writing a well adapted basis of E_6 roots is a little bit more laborious but it can be done. We find:

$$\begin{aligned}\alpha_1 &= \left\{ 0, 0, -\frac{\sqrt{3}}{2}, \frac{1}{2\sqrt{5}}, \sqrt{\frac{6}{5}}, 0 \right\} &= \{ \bar{\alpha}_1, 0 \} \\ \alpha_2 &= \left\{ -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{6}}, \frac{2}{\sqrt{3}}, 0, 0, 0 \right\} &= \{ \bar{\alpha}_2, 0 \} \\ \alpha_3 &= \{ \sqrt{2}, 0, 0, 0, 0, 0 \} &= \{ \bar{\alpha}_3, 0 \} \\ \alpha_4 &= \left\{ -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{6}}, -\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{5}}, -\sqrt{\frac{3}{10}}, \frac{1}{\sqrt{2}} \right\} &= \left\{ \bar{\mathbf{w}}_h, \frac{1}{\sqrt{2}} \right\} \\ \alpha_5 &= \left\{ -\frac{1}{\sqrt{2}}, -\sqrt{\frac{3}{2}}, 0, 0, 0, 0 \right\} &= \{ \bar{\alpha}_4, 0 \} \\ \alpha_6 &= \left\{ 0, \sqrt{\frac{2}{3}}, -\frac{1}{2\sqrt{3}}, -\frac{\sqrt{5}}{2}, 0, 0 \right\} &= \{ \bar{\alpha}_5, 0 \}\end{aligned} \quad (6.25)$$

In this basis we can check that the five 5-vectors $\bar{\alpha}_i$ ($i = 1, \dots, 5$) constitute a simple root basis for the

$A_5 \simeq \text{SU}(1, 5)$ root system, namely:

$$\langle \bar{\alpha}_i, \bar{\alpha}_j \rangle = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix} = \text{Cartan matrix of } A_5 \quad (6.26)$$

while:

$$\mathbf{w}_h = \left\{ -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{6}}, -\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{5}}, -\sqrt{\frac{3}{10}} \right\} \quad (6.27)$$

is the highest weight of the **20**-dimensional representation of $\text{SU}(1, 5)$. Finally in this basis the highest root ψ defined by eq.(6.24) takes the expected form:

$$\psi = \{0, 0, 0, 0, 0, \sqrt{2}\} \quad (6.28)$$

N=4 The case of $\mathcal{N} = 4$ supergravity is the first where the scalar manifold is not completely fixed, since we can choose the number n_m of vector multiplets that we can couple to the graviton multiplet. In any case, once n_m is fixed the scalar manifold is also fixed and we have:

$$\mathcal{M}_{N=4, D=4} = \frac{\text{SL}(2, \mathbb{R})_0}{\text{O}(2)} \otimes \frac{\text{SO}(6, \mathbf{n}_m)}{\text{SO}(6) \times \text{SO}(\mathbf{n}_m)} \quad (6.29)$$

The total number of vectors $n_v = 6 + n_m$ is also fixed and the symplectic representation \mathbf{W} of the duality algebra

$$\mathbb{U}_{D=4} = \text{SL}(2, \mathbb{R})_0 \times \text{SO}(6, \mathbf{n}_m) \quad (6.30)$$

to which the vectors are assigned and which determines the embedding:

$$\text{SL}(2, \mathbb{R})_0 \times \text{SO}(6) \times \text{SO}(\mathbf{n}_m) \mapsto \text{Sp}(12 + 2\mathbf{n}_m, \mathbb{R}) \quad (6.31)$$

is also fixed, namely $\mathbf{W} = (\mathbf{2}_0, \mathbf{6} + \mathbf{n}_m)$, $\mathbf{2}_0$ being the fundamental representation of $\text{SL}(2, \mathbb{R})_0$ and $\mathbf{6} + \mathbf{n}_m$ the fundamental vector representation of $\text{SO}(6, \mathbf{n}_m)$.

The $D = 3$ algebra is, $\mathbb{U}_{D=3} = \text{SO}(8, \mathbf{n}_m + 2)$. Correspondingly the form taken by the general decomposition (2.20) is the following one:


$$\text{adj}(\text{SO}(8, \mathbf{n}_m + 2)) = \text{adj}(\text{SL}(2, \mathbb{R})_0) \oplus \text{adj}(\text{SO}(6, \mathbf{n}_m)) \oplus \text{adj}(\text{SL}(2, \mathbb{R})_E) \oplus (\mathbf{2}_E, \mathbf{2}_0, \mathbf{6} + \mathbf{n}_m) \quad (6.32)$$

where $\mathbf{2}_{E,0}$ are the fundamental representations respectively of $\text{SL}(2, \mathbb{R})_E$ and of $\text{SL}(2, \mathbb{R})_0$.

In order to give a Dynkin Weyl description of these algebras, we are forced to distinguish the case of an odd and even number of vector multiplets. In the first case both $\mathbb{U}_{D=3}$ and $\mathbb{U}_{D=4}$ are non simply laced algebras of the B -type, while in the second case they are both simply laced algebras of the D -type

$$n_m = \begin{cases} 2k & \rightarrow \mathbb{U}_{D=4} \simeq D_{k+3} \\ 2k+1 & \rightarrow \mathbb{U}_{D=4} \simeq B_{k+3} \end{cases} \quad (6.33)$$

Just for simplicity and for shortness we choose to discuss only the even case $n_m = 2k$ which is described by fig.5.



$$\begin{aligned}\psi &= \alpha_1 + 2\alpha_2 + 2\alpha_3 + \dots + 2\alpha_{\ell-2} + \alpha_{\ell-1} + \alpha_{\ell} \\ (\psi, \alpha_2) &= 1 \quad ; \quad (\psi, \alpha_i) = 0 \quad i \neq 2\end{aligned}$$

Figure 5: The Dynkin diagram of D_{4+k+1} . The algebra D_{4+k+1} is that of the group $\text{SO}(8, 2k+2)$ corresponding to the σ -model reduction of $\mathcal{N} = 4$ supergravity coupled to $n_m = 2k$ vector multiplets. The only simple root which has non vanishing grading with respect to the highest one ψ is α_2 . Removing it (black circle) we are left with the algebra $D_{4+k-1} \oplus A_1$ which is indeed the duality algebra in $D = 4$, namely $\text{SO}(6, 2k) \oplus \text{SL}(2, \mathbb{R})_0$. The black root α_2 is the highest weight of the symplectic representation of the vector fields, namely the $\mathbf{W} = (2_0, \mathbf{6} + 2\mathbf{k})$.

In this case we consider the $\mathbb{U}_{D=3} = \text{SO}(8, 2k+2)$ Lie algebra whose Dynkin diagram is that of D_{5+k} . Naming ϵ_i the unit vectors in an Euclidean ℓ -dimensional space where $\ell = 5 + k$, a well adapted basis of simple roots for the considered algebra is the following one:

$$\begin{aligned}\alpha_1 &= \sqrt{2} \epsilon_1 \\ \alpha_2 &= -\frac{1}{\sqrt{2}} \epsilon_1 - \epsilon_2 + \frac{1}{\sqrt{2}} \epsilon_{\ell} \\ \alpha_3 &= \epsilon_2 - \epsilon_3 \\ \alpha_4 &= \epsilon_3 - \epsilon_4 \\ \dots &= \dots \\ \alpha_{\ell-1} &= \epsilon_{\ell-2} - \epsilon_{\ell-1} \\ \alpha_{\ell} &= \epsilon_{\ell-2} + \epsilon_{\ell-1}\end{aligned}\tag{6.34}$$

which is quite different from the usual presentation but yields the correct Cartan matrix. In this basis the highest root of the algebra:

$$\psi = \alpha_1 + 2\alpha_2 + 2\alpha_3 + \dots + 2\alpha_{\ell-2} + \alpha_{\ell-1} + \alpha_{\ell}\tag{6.35}$$

takes the desired form:

$$\psi = \sqrt{2} \epsilon_{\ell}\tag{6.36}$$

In the same basis the $\alpha_W = \alpha_2$ root has also the expect form:

$$\alpha_W = \left(\mathbf{w}, \frac{1}{\sqrt{2}} \right)\tag{6.37}$$

where:

$$\mathbf{w} = -\frac{1}{\sqrt{2}} \epsilon_1 - \epsilon_2\tag{6.38}$$

is the weight of the symplectic representation $\mathbf{W} = (\mathbf{2}_0, \mathbf{6} + \mathbf{2k})$. Indeed $-\frac{1}{\sqrt{2}}\epsilon_1$ is the fundamental weight for the Lie algebra $\mathrm{SL}(2, \mathbb{R})_0$, whose root is $\alpha_1 = \sqrt{2}\epsilon_1$, while $-\epsilon_2$ is the highest weight for the vector representation of the algebra $\mathrm{SO}(6, 2k)$, whose roots are $\alpha_3, \alpha_4, \dots, \alpha_\ell$.

Next we briefly comment on the Tits Satake projection. The algebra $\mathrm{SO}(8, n_m + 2)$ is maximally split only for $n_m = 5, 6, 7$. The case $n_m = 6$, from the superstring view point, corresponds to the case of Neveu-Schwarz vector multiplets in a toroidal compactification. For a different number of vector multiplets, in particular for $n_m > 7$ the study of extremal black holes involves considering the Tits Satake projection, which just yields the universal algebra

$$\mathbb{U}_{N=4, D=3}^{TS} = \mathfrak{so}(8, 9) \quad (6.39)$$

7 Tits Satake decompositions of the \mathbf{W} representations

As we stressed in the introduction, one of our main goals is to compare the classification of extremal black holes by means of *charge orbits* with their classification by means of H^* *orbits*. Charge orbits means orbits of the $\mathbb{U}_{D=4}$ group in the \mathbf{W} -representation.

For this reason, in the present section we consider the decomposition of the \mathbf{W} -representations with respect to Tits-Satake subalgebras and Paint groups for all the non-exotic models. The relevant \mathbf{W} -representations are listed in table 6. In table 7 we listed the \mathbf{W} -representations for the exotic models.

In [36] the *paint algebra* was defined as the algebra of external automorphism of the solvable Lie algebra $\mathrm{Solv}_{\mathcal{M}}$ generating the non-compact symmetric space: $\mathcal{M} = \mathbb{U}/H$, namely

$$\mathbb{G}_{\mathrm{paint}} = \mathrm{Aut}_{\mathrm{Ext}} [\mathrm{Solv}_{\mathcal{M}}]. \quad (7.1)$$

where:

$$\mathrm{Aut}_{\mathrm{Ext}} [\mathrm{Solv}_{\mathcal{M}}] \equiv \frac{\mathrm{Aut} [\mathrm{Solv}_{\mathcal{M}}]}{\mathrm{Solv}_{\mathcal{M}}}, \quad (7.2)$$

Given the paint algebra $\mathbb{G}_{\mathrm{paint}} \subset \mathbb{U}$ and the Tits Satake subalgebra $\mathbb{G}_{\mathrm{TS}} \subset \mathbb{U}$, whose construction we have briefly recalled above, following [36] one introduces the *sub Tits Satake* and *sub paint* algebras as the centralizers of the paint algebra and of the Tits Satake algebra, respectively. In other words we have:

$$\mathfrak{s} \in \mathbb{G}_{\mathrm{subTS}} \subset \mathbb{G}_{\mathrm{TS}} \subset \mathbb{U} \quad \Leftrightarrow \quad [\mathfrak{s}, \mathbb{G}_{\mathrm{paint}}] = 0 \quad (7.3)$$

and

$$\mathfrak{t} \in \mathbb{G}_{\mathrm{subpaint}} \subset \mathbb{G}_{\mathrm{paint}} \subset \mathbb{U} \quad \Leftrightarrow \quad [\mathfrak{t}, \mathbb{G}_{\mathrm{TS}}] = 0 \quad (7.4)$$

A very important property of the paint and subpaint algebras is that they are conserved in the dimensional reduction, namely they are the same for $\mathbb{U}_{D=4}$ and $\mathbb{U}_{D=3}$.

In the next lines we analyze the decomposition of the \mathbf{W} -representations with respect to these subalgebras for each Tits Satake universality class of non maximally split models. In the case of maximally split models there is no paint algebra and there is nothing with respect to which to decompose.

7.1 Universality class $\mathfrak{sp}(6, \mathbb{R}) \Rightarrow \mathfrak{f}_{4(4)}$

In this case the sub Tits Satake Lie algebra is

$$\mathbb{G}_{\mathrm{subTS}} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R}) \subset \mathfrak{sp}(6, \mathbb{R}) = \mathbb{G}_{\mathrm{TS}} \quad (7.5)$$

and the **W**-representation of the maximally split model decomposes as follows:

$$\mathbf{14}' \xrightarrow{\mathbb{G}_{\text{subTS}}} (\mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{2}, \mathbf{2}) \quad (7.6)$$

This decomposition combines in the following way with the paint group representations in the various models belonging to the same universality class.

7.1.1 $\mathfrak{su}(3, 3)$ model

For this case the paint algebra is

$$\mathbb{G}_{\text{paint}} = \mathfrak{so}(2) \oplus \mathfrak{so}(2) \quad (7.7)$$

and the **W**-representation is the **20** dimensional of $\mathfrak{su}(3, 3)$ corresponding to an antisymmetric tensor with a reality condition of the form:

$$t_{\alpha\beta\gamma}^* = \frac{1}{3!} \epsilon_{\alpha\beta\gamma\delta\eta\theta} t_{\delta\eta\theta} \quad (7.8)$$

The decomposition of this representation with respect to the Lie algebra $\mathbb{G}_{\text{paint}} \oplus \mathbb{G}_{\text{subTS}}$ is the following one:

$$\mathbf{20} \xrightarrow{\mathbb{G}_{\text{paint}} \oplus \mathbb{G}_{\text{subTS}}} (2, q_1 | \mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (2, q_2 | \mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (2, q_3 | \mathbf{1}, \mathbf{1}, \mathbf{2}) \oplus (1, 0 | \mathbf{2}, \mathbf{2}, \mathbf{2}) \quad (7.9)$$

where $(2, q)$ means a doublet of $\mathfrak{so}(2) \oplus \mathfrak{so}(2)$ with a certain grading q with respect to the generators, while $(1, 0)$ means the singlet that has 0 grading with respect to both generators. The subpaint algebra in this case is $\mathbb{G}_{\text{subpaint}} = 0$ and the decomposition of the same **W**-representation with respect to $\mathbb{G}_{\text{subpaint}} \oplus \mathbb{G}_{\text{TS}}$ is:

$$\mathbf{20} \xrightarrow{\mathbb{G}_{\text{subpaint}} \oplus \mathbb{G}_{\text{TS}}} \mathbf{6} \oplus \mathbf{14} \quad (7.10)$$

This follows from the decomposition of the **6** of $\mathfrak{sp}(6, \mathbf{R})$ with respect to the sub Tits Satake algebra (7.5):

$$\mathbf{6} \xrightarrow{\mathbb{G}_{\text{subTS}}} (\mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{2}) \quad (7.11)$$

7.1.2 $\mathfrak{so}^*(12)$ model

For this case the paint algebra is

$$\mathbb{G}_{\text{paint}} = \mathfrak{so}(3) \oplus \mathfrak{so}(3) \oplus \mathfrak{so}(3) \quad (7.12)$$

and the **W**-representation is the $\mathbf{32}_s$ dimensional spinorial representation of $\mathfrak{so}^*(12)$. The decomposition of this representation with respect to the Lie algebra $\mathbb{G}_{\text{paint}} \oplus \mathbb{G}_{\text{subTS}}$ is the following one:

$$\mathbf{32}_s \xrightarrow{\mathbb{G}_{\text{paint}} \oplus \mathbb{G}_{\text{subTS}}} (\underline{2}, \underline{2}, \underline{1} | \mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\underline{2}, \underline{1}, \underline{2} | \mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (\underline{1}, \underline{1}, \underline{2} | \mathbf{1}, \mathbf{1}, \mathbf{2}) \oplus (\underline{1}, \underline{1}, \underline{1} | \mathbf{2}, \mathbf{2}, \mathbf{2}) \quad (7.13)$$

where $\underline{2}$ means the doublet spinor representation of $\mathfrak{so}(3)$. The subpaint algebra in this case is $\mathbb{G}_{\text{paint}} = \mathfrak{so}(3)_{\text{diag}}$ and the decomposition of the same **W**-representation with respect to $\mathbb{G}_{\text{subpaint}} \oplus \mathbb{G}_{\text{TS}}$ is:

$$\mathbf{32}_s \xrightarrow{\mathbb{G}_{\text{TS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{6} | \underline{3}) \oplus (\mathbf{14}' | \underline{1}) \quad (7.14)$$

This follows from the decomposition of the product $\underline{2} \times \underline{2}$ of $\mathfrak{so}(3)_{\text{diag}}$ times the Tits Satake algebra (7.5):

$$\underline{2} \times \underline{2} = \underline{3} \oplus \underline{1} \quad (7.15)$$

7.1.3 $\mathfrak{e}_{7(-25)}$ model

For this case the paint algebra is

$$\mathbb{G}_{\text{paint}} = \mathfrak{so}(8) \quad (7.16)$$

and the \mathbf{W} -representation is the fundamental **56** dimensional representation of $\mathfrak{e}_{7(-25)}$. The decomposition of this representation with respect to the Lie algebra $\mathbb{G}_{\text{paint}} \oplus \mathbb{G}_{\text{subTS}}$ is the following one:

$$\mathbf{56} \xrightarrow{\mathbb{G}_{\text{paint}} \oplus \mathbb{G}_{\text{subTS}}} (\mathbf{8}_v | \mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{8}_s | \mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (\mathbf{8}_c | \mathbf{1}, \mathbf{1}, \mathbf{2}) \oplus (\mathbf{1} | \mathbf{2}, \mathbf{2}, \mathbf{2}) \quad (7.17)$$

where $\mathbf{8}_{v,s,c}$ are the three inequivalent eight-dimensional representations of $\mathfrak{so}(8)$, the vector, the spinor and the conjugate spinor. The subpaint algebra in this case is $\mathbb{G}_{\text{paint}} = \mathfrak{g}_{2(-14)}$ with respect to which all three 8-dimensional representations of $\mathfrak{so}(8)$ branch as follows:

$$\mathbf{8}_{v,s,c} \xrightarrow{\mathfrak{g}_{2(-14)}} \mathbf{7} \oplus \mathbf{1} \quad (7.18)$$

In view of this the decomposition of the same \mathbf{W} -representation with respect to $\mathbb{G}_{\text{subpaint}} \oplus \mathbb{G}_{\text{TS}}$ is:

$$\mathbf{56} \xrightarrow{\mathbb{G}_{\text{TS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{6} | \mathbf{7}) \oplus (\mathbf{14}' | \mathbf{1}) \quad (7.19)$$

7.2 Universality class $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 3) \Rightarrow \mathfrak{so}(4, 5)$

This case corresponds to one of the possible infinite families of $\mathcal{N} = 2$ theories with a symmetric homogeneous special Kähler manifold and a number of vector multiplets larger than three ($n = 3 + p$). The other infinite family corresponds instead to one of the three exotic models.

The generic element of this infinite class corresponds to the following algebras:

$$\begin{aligned} \mathbb{U}_{D=4} &= \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 2 + p) \\ \mathbb{U}_{D=3} &= \mathfrak{so}(4, 4 + p) \end{aligned} \quad (7.20)$$

In this case the sub Tits Satake algebra is:

$$\mathbb{G}_{\text{subTS}} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R}) \simeq \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 2) \subset \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 3) = \mathbb{G}_{\text{TS}} \quad (7.21)$$

an the paint and subpaint algebras are as follows:

$$\begin{aligned} \mathbb{G}_{\text{paint}} &= \mathfrak{so}(p) \\ \mathbb{G}_{\text{subpaint}} &= \mathfrak{so}(p - 1) \end{aligned} \quad (7.22)$$

The symplectic \mathbf{W} representation of $\mathbb{U}_{D=4}$ is the tensor product of the fundamental representation of $\mathfrak{sl}(2)$ with the fundamental vector representation of $\mathfrak{so}(2, 2 + p)$, namely

$$\mathbf{W} = (\mathbf{2} | \mathbf{4} + p) \quad ; \quad \dim \mathbf{W} = 8 + 2p \quad (7.23)$$

The decomposition of this representation with respect to $\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{subpaint}}$ is the following one:

$$\mathbf{W} \xrightarrow{\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{2}, \mathbf{2}, \mathbf{2} | \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}, \mathbf{1} | \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}, \mathbf{1} | p - 1) \quad (7.24)$$

where $\mathbf{2}, \mathbf{2}, \mathbf{2}$ denotes the tensor product of the three fundamental representations of $\mathfrak{sl}(2, \mathbb{R})^3$. Similarly $\mathbf{2}, \mathbf{1}, \mathbf{1}$ denotes the doublet of the first $\mathfrak{sl}(2, \mathbb{R})$ tensored with the singlets of the following two $\mathfrak{sl}(2, \mathbb{R})$ algebras. The representations appearing in (7.24) can be grouped in order to reconstruct full representations either of the complete Tits Satake or of the complete paint algebras. In this way one obtains:

$$\begin{aligned} \mathbf{W} &\xrightarrow{\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{paint}}} (\mathbf{2}, \mathbf{2}, \mathbf{2}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}, \mathbf{1}|p+1) \\ \mathbf{W} &\xrightarrow{\mathbb{G}_{\text{TS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{2}, \mathbf{5}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|p-1) \end{aligned} \quad (7.25)$$

7.3 Universality class $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, 7) \Rightarrow \mathfrak{so}(8, 9)$

This case, which corresponds to an $\mathcal{N} = 4$ theory with a number of vector multiplets larger than six ($n = 6 + p$) presents a very strong similarity with the previous $\mathcal{N} = 2$ case.

The generic element of this infinite class corresponds to the following algebras:

$$\begin{aligned} \mathbb{U}_{D=4} &= \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, 6+p) \\ \mathbb{U}_{D=3} &= \mathfrak{so}(8, 8+p) \end{aligned} \quad (7.26)$$

In this case the sub Tits Satake algebra is:

$$\mathbb{G}_{\text{subTS}} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, 6) \subset \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, 7) = \mathbb{G}_{\text{TS}} \quad (7.27)$$

an the paint and subpaint algebras are the same as in the previous $\mathcal{N} = 2$ case, namely:

$$\begin{aligned} \mathbb{G}_{\text{paint}} &= \mathfrak{so}(p) \\ \mathbb{G}_{\text{subpaint}} &= \mathfrak{so}(p-1) \end{aligned} \quad (7.28)$$

The symplectic \mathbf{W} representation of $\mathbb{U}_{D=4}$ is the tensor product of the fundamental representation of $\mathfrak{sl}(2)$ with the fundamental vector representation of $\mathfrak{so}(6, 6+p)$, namely

$$\mathbf{W} = (\mathbf{2}|\mathbf{12}+p) \quad ; \quad \dim \mathbf{W} = 24 + 2p \quad (7.29)$$

The decomposition of this representation with respect to $\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{subpaint}}$ is the following one:

$$\mathbf{W} \xrightarrow{\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{2}, \mathbf{12}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|p) \quad (7.30)$$

Just as above the three representations appearing in (7.30) can be grouped in order to obtain either representation of the complete Tits Satake or of the complete paint algebras. This yields

$$\begin{aligned} \mathbf{W} &\xrightarrow{\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{paint}}} (\mathbf{2}, \mathbf{12}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|p+1) \\ \mathbf{W} &\xrightarrow{\mathbb{G}_{\text{TS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{2}, \mathbf{13}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|p) \end{aligned} \quad (7.31)$$

7.4 The universality classes $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, n) \Rightarrow \mathfrak{so}(8, n+2)$ with $n \leq 5$

These classes correspond to the $\mathcal{N} = 4$ theories with a number $n = 1, 2, 3, 4, 5$ of vector multiplets. In each case we have the following algebras:

$$\begin{aligned} \mathbb{U}_{D=4} &= \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(6, n) \\ \mathbb{U}_{D=3} &= \mathfrak{so}(8, n+2) \end{aligned} \quad (7.32)$$

In all these cases the Tits Satake and sub Tits Satake algebras are:

$$\begin{aligned}\mathbb{G}_{\text{TS}} &= \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(n+1, n) \\ \mathbb{G}_{\text{subTS}} &= \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(n, n)\end{aligned}\tag{7.33}$$

and the paint and subpaint algebras are:

$$\begin{aligned}\mathbb{G}_{\text{paint}} &= \mathfrak{so}(6-n) \\ \mathbb{G}_{\text{subpaint}} &= \mathfrak{so}(5-n)\end{aligned}\tag{7.34}$$

The symplectic \mathbb{W} representation is the tensor product of the doublet representation of $\mathfrak{sl}(2)$ with the fundamental representation of $\mathfrak{so}(6, n)$, namely

$$\mathbf{W} = (\mathbf{2}, \mathbf{6} + \mathbf{n})\tag{7.35}$$

and its decomposition with respect to the $\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{subpaint}}$ algebra is as follows

$$\mathbf{W} \xrightarrow{\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{2}, \mathbf{2n}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|\mathbf{5} - \mathbf{n})\tag{7.36}$$

which, with the same procedure as above leads to:

$$\begin{aligned}\mathbf{W} &\xrightarrow{\mathbb{G}_{\text{subTS}} \oplus \mathbb{G}_{\text{paint}}} (\mathbf{2}, \mathbf{2n}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|\mathbf{6} - \mathbf{n}) \\ \mathbf{W} &\xrightarrow{\mathbb{G}_{\text{TS}} \oplus \mathbb{G}_{\text{subpaint}}} (\mathbf{2}, \mathbf{2n} + \mathbf{1}|\mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}|\mathbf{5} - \mathbf{n})\end{aligned}\tag{7.37}$$

7.5 \mathbf{W} -representations of the maximally split non exotic models

In the previous subsections we have analysed the Tits-Satake decomposition of the \mathbf{W} -representation for all those models that are non maximally split. The remaining models are the maximally split ones for which there is no paint algebra and the Tits Satake projection is the identity map. For the reader's convenience we have extracted from table 3 the list of such models and presented it in table 5. As we see from the table we have essentially five type of models:

1. The $E_{7(7)}$ model corresponding to $\mathcal{N} = 8$ supergravity where the \mathbf{W} -representation is the fundamental $\mathbf{56}$.
2. The $SU(1, 1)$ non exotic model where the \mathbf{W} -representation is the $j = \frac{3}{2}$ of $\mathfrak{so}(1, 2) \sim \mathfrak{su}(1, 1)$
3. The $Sp(6, \mathbb{R})$ model where the \mathbf{W} -representation is the $\mathbf{14}'$ (antisymmetric symplectic traceless three-tensor).
4. The models $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q)$ where the \mathbf{W} -representation is the $(2, 2q)$, namely the tensor product of the two fundamentals.
5. The models $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q+1)$ where the \mathbf{W} -representation is the $(2, 2q+1)$, namely the tensor product of the two fundamentals.

#	TS D=4	TS D=3	coset D=4	coset D=3	Paint Group	subP Group	susy
1	$\frac{E_7(7)}{SU(8)}$	$\frac{E_8(8)}{SO^*(16)}$	$\frac{E_7(7)}{SU(8)}$	$\frac{E_8(8)}{SO^*(16)}$	1	1	$\mathcal{N} = 8$
2	$\frac{SU(1,1)}{U(1)}$	$\frac{G_2(2)}{SL(2,\mathbb{R}) \times SL(2,\mathbb{R})}$	$\frac{SU(1,1)}{U(1)}$	$\frac{G_2(2)}{SL(2,\mathbb{R}) \times SL(2,\mathbb{R})}$	1	1	$\mathcal{N} = 2$ $n=1$
3	$\frac{Sp(6,\mathbb{R})}{SU(3) \times U(1)}$	$\frac{F_4(4)}{Sp(6,\mathbb{R}) \times SL(2,\mathbb{R})}$	$\frac{Sp(6,\mathbb{R})}{SU(3) \times U(1)}$	$\frac{F_4(4)}{Sp(6,\mathbb{R}) \times SL(2,\mathbb{R})}$	1	1	$\mathcal{N} = 2$ $n=6$
11	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(6,5)}{SO(6) \times SO(5)}$	$\frac{SO(8,7)}{SO(6,2) \times SO(2,5)}$	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(6,5)}{SO(6) \times SO(5)}$	$\frac{SO(8,7)}{SO(6,2) \times SO(2,5)}$	1	1	$\mathcal{N} = 4$ $n=5$
12	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(6,6)}{SO(6) \times SO(6)}$	$\frac{SO(8,8)}{SO(6,2) \times SO(2,6)}$	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(6,6)}{SO(6) \times SO(6)}$	$\frac{SO(8,8)}{SO(6,2) \times SO(2,6)}$	1	1	$\mathcal{N} = 4$ $n=6$
13	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(6,7)}{SO(6) \times SO(7)}$	$\frac{SO(8,9)}{SO(6,2) \times SO(2,7)}$	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(6,7)}{SO(6) \times SO(7)}$	$\frac{SO(8,9)}{SO(6,2) \times SO(2,7)}$	1	1	$\mathcal{N} = 4$ $n=7$
14	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(2,1)}{SO(2)}$	$\frac{SO(4,3)}{SO(2,2) \times SO(2,1)}$	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(2,1)}{SO(2)}$	$\frac{SO(4,3)}{SO(2,2) \times SO(2,1)}$	1	1	$\mathcal{N} = 2$ $n=2$
15	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(2,2)}{SO(2) \times SO(2)}$	$\frac{SO(4,4)}{SO(2,2) \times SO(2,2)}$	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(2,2)}{SO(2) \times SO(2)}$	$\frac{SO(4,4)}{SO(2,2) \times SO(2,2)}$	1	1	$\mathcal{N} = 2$ $n=3$
16	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(2,3)}{SO(2) \times SO(3)}$	$\frac{SO(4,5)}{SO(2,2) \times SO(2,3)}$	$\frac{SL(2,\mathbb{R})}{O(2)} \times \frac{SO(2,3)}{SO(2) \times SO(3)}$	$\frac{SO(4,5)}{SO(2,2) \times SO(2,3)}$	1	1	$\mathcal{N} = 2$ $n=4$

Table 5: The list of *non-exotic* homogenous symmetric scalar manifolds appearing in $D = 4$ supergravity which are also maximally split. For these models the paint group is the identity group.

Therefore, for the above maximally split models, the *charge classification* of black holes reduces to the classification of $U_{D=4}$ orbits in the mentioned \mathbf{W} -representations. Actually such orbits are sufficient also for the non maximally split models. Indeed each of the above 5-models correspond to one Tits Satake universality class and, within each universality class, the only relevant part of the \mathbf{W} -representation is the subpaint group singlet which is universal for all members of the class. This is precisely what we verified in the previous subsections.

For instance for all members of the universality class of $Sp(6, \mathbb{R})$, the \mathbf{W} -representation splits as follows with respect to the subalgebra $\mathfrak{sp}(6, \mathbb{R}) \oplus \mathbb{G}_{\text{subpaint}}$:

$$\mathbf{W}^{\mathfrak{sp}(6, \mathbb{R}) \oplus \mathbb{G}_{\text{subpaint}}} \xrightarrow{\quad} (\mathbf{6} | \mathcal{D}_{\text{subpaint}}) + (\mathbf{14}' | \mathbf{1}_{\text{subpaint}}) \quad (7.38)$$

where the representation $\mathcal{D}_{\text{subpaint}}$ is the following one for the three non-maximally split members of the class:

$$\mathcal{D}_{\text{subpaint}} = \begin{cases} \mathbf{1} & \text{of } \mathbf{1} & \text{for the } \mathfrak{su}(3, 3) - \text{model} \\ \mathbf{3} & \text{of } \mathfrak{so}(3) & \text{for the } \mathfrak{so}^*(12) - \text{model} \\ \mathbf{7} & \text{of } \mathfrak{g}_{2(-14)} & \text{for the } \mathfrak{e}_{7(-25)} - \text{model} \end{cases} \quad (7.39)$$

Clearly the condition:

$$(\mathbf{6} | \mathcal{D}_{\text{subpaint}}) = 0 \quad (7.40)$$

imposed on a vector in the \mathbf{W} -representation breaks the group $U_{D=4}$ to its Tits Satake subgroup. The key point is that each \mathbf{W} -orbit of the big group $U_{D=4}$ crosses the locus (7.40) so that the classification of $\mathrm{Sp}(6, \mathbb{R})$ orbits in the $\mathbf{14}'$ -representation exhausts the classification of \mathbf{W} -orbits for all members of the universality class.

In order to prove that the gauge (7.40) is always reachable it suffices to show that the representation $(\mathbf{6} | \mathcal{D}_{\text{subpaint}})$ always appears at least once in the decomposition of the Lie algebra $\mathbb{U}_{D=4}$ with respect to the subalgebra $\mathfrak{sp}(6, \mathbb{R}) \oplus \mathbb{G}_{\text{subpaint}}$. The corresponding parameters of the big group can be used to set to zero the projection of the \mathbf{W} -vector onto $(\mathbf{6} | \mathcal{D}_{\text{subpaint}})$.

The required condition is easily verified since we have:

$$\begin{array}{ccc}
\underbrace{\text{adj } \mathfrak{su}(3, 3)}_{\mathbf{35}} & \xrightarrow{\mathfrak{sp}(6, \mathbb{R})} & \underbrace{\text{adj } \mathfrak{sp}(6, \mathbb{R})}_{\mathbf{21}} \oplus \mathbf{6} \oplus \mathbf{6} \oplus \mathbf{1} \oplus \mathbf{1} \\
\underbrace{\text{adj } \mathfrak{so}^*(12)}_{\mathbf{66}} & \xrightarrow{\mathfrak{sp}(6, \mathbb{R}) \oplus \mathfrak{so}(3)} & \underbrace{\text{adj } \mathfrak{sp}(6, \mathbb{R})}_{\mathbf{21}} \oplus \underbrace{\text{adj } \mathfrak{so}(3)}_{\mathbf{3}} \oplus (\mathbf{6}, \mathbf{3}) \oplus (\mathbf{6}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{3}) \\
\underbrace{\text{adj } \mathfrak{e}_{7(-25)}}_{\mathbf{133}} & \xrightarrow{\mathfrak{sp}(6, \mathbb{R}) \oplus \mathfrak{g}_{2(-14)}} & \underbrace{\text{adj } \mathfrak{sp}(6, \mathbb{R})}_{\mathbf{21}} \oplus \underbrace{\text{adj } \mathfrak{g}_{2(-14)}}_{\mathbf{14}} \oplus (\mathbf{6}, \mathbf{7}) \oplus (\mathbf{6}, \mathbf{7}) \oplus (\mathbf{1}, \mathbf{7}) \oplus (\mathbf{1}, \mathbf{7})
\end{array} \tag{7.41}$$

The reader cannot avoid being impressed by the striking similarity of the above decompositions which encode the very essence of Tits Satake universality. Indeed the representations of the common Tits Satake subalgebra appearing in the decomposition of the adjoint are the same for all members of the class. They are simply uniformly assigned to the fundamental representation of the subpaint algebra which is different in the three cases. The representation $(\mathbf{6} | \mathcal{D}_{\text{subpaint}})$ appears twice in these decompositions and can be used to reach the gauge (7.40) as we claimed above.

For the models of type $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q + p)$ having $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q + 1)$ as Tits Satake subalgebra and $\mathfrak{so}(p - 1)$ as subpaint algebra the decomposition of the \mathbf{W} -representation is the following one:

$$\mathbf{W} = (\mathbf{2}, \mathbf{2q} + \mathbf{p}) \xrightarrow{\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q+1) \oplus \mathfrak{so}(p-1)} (\mathbf{2}, \mathbf{2q} + \mathbf{1} | \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1} | \mathbf{p} - \mathbf{1}) \tag{7.42}$$

and the question is whether each $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q + p)$ orbit in the $(\mathbf{2}, \mathbf{2q} + \mathbf{p})$ representation intersects the $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q + 1) \oplus \mathfrak{so}(p - 1)$ -invariant locus:

$$(\mathbf{2}, \mathbf{1} | \mathbf{p} - \mathbf{1}) = 0 \tag{7.43}$$

The answer is yes since we always have enough parameters in the coset

$$\frac{\mathrm{SL}(2, \mathbb{R}) \times \mathrm{SO}(q, q + p)}{\mathrm{SL}(2, \mathbb{R}) \times \mathrm{SO}(q, q + 1) \times \mathrm{SO}(p - 1)} \tag{7.44}$$

to reach the desired gauge (7.43). Indeed let us observe the decomposition:

$$\text{adj } [\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(q, q + p)] = \text{adj } [\mathfrak{sl}(2, \mathbb{R})] \oplus \text{adj } [\mathfrak{so}(q, q + 1)] \oplus \text{adj } [\mathfrak{so}(p - 1)] \oplus (\mathbf{1}, \mathbf{2q} + \mathbf{1} | \mathbf{p} - \mathbf{1}) \tag{7.45}$$

The $2q + 1$ vectors of $\mathfrak{so}(p - 1)$ appearing in (7.45) are certainly sufficient to set to zero the 2 vectors of $\mathfrak{so}(p - 1)$ appearing in \mathbf{W} .

The conclusion therefore is that the classification of charge-orbits for all supergravity models can be performed by restriction to the Tits Satake sub-model. The same we show, in the next section, to be true at the level of the classification based on \mathbb{H}^* orbits of the Lax operators, so that the final comparison of the two classifications can be performed by restriction to the Tits Satake subalgebras.

8 Tits Satake reduction of the \mathbb{H}^* subalgebra and of its representation \mathbb{K}^*

In the σ -model approach to black hole solutions one arrives at the new coset manifold (2.19). The structure of the enlarged group $\mathbb{U}_{D=3}$ and of its Lie algebra $\mathbb{U}_{D=3}$ was discussed in eq.(2.20). The subgroups \mathbb{H}^* are listed in table (6) for the non exotic models and in table (7) for the exotic ones. The coset generators fall into a representation of \mathbb{H}^* that we name \mathbb{K}^* . The Lax operator L_0 which determines the spherically symmetric black hole solution up to boundary conditions of the scalar fields at infinity is just an element of such a representation:

$$L_0 \in \mathbb{K}^* \quad (8.1)$$

so that the classification of spherical black holes is reduced to the classification of \mathbb{H}^* orbits in the \mathbb{K}^* representation. On the other hand, in part one of the paper we already saw how nilpotent orbits can be associated to multicenter solutions.

In this paper we focus on non-exotic models that admit a regular Tits Satake projection and we postpone the analysis of the exotic ones to a future publication.

A first general remark concerns the structure of \mathbb{H}^* in all those models that correspond to $\mathcal{N} = 2$ supersymmetry. In these cases the \mathbb{H}^* subalgebra is isomorphic to $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathbb{U}_{D=4}$ so that we have a decomposition of the $\mathbb{U}_{D=3}$ Lie algebra with respect to \mathbb{H}^* completely analogous to that in equation (2.20), namely:

$$\text{adj}(\mathbb{U}_{D=3}) = \underbrace{\text{adj}(\widehat{\mathbb{U}_{D=4}})}_{\mathbb{H}^*} \oplus \underbrace{\text{adj}(\mathfrak{sl}(2, \mathbb{R})_{\mathbb{H}^*})}_{\mathbb{K}^*} \oplus \underbrace{(2_{\mathbb{H}^*}, \widehat{\mathbf{W}})}_{\mathbb{K}^*} \quad (8.2)$$

Hence the representation \mathbb{K}^* which contains the Lax operators has a structure analogous to the representation which contains the generators of $\mathbb{U}_{D=4}$ that originate from the vector fields, namely: $(2_{\mathbb{H}^*}, \widehat{\mathbf{W}})$. This means that in all these models, by means of exactly the same argument as utilized above, we can always reach the gauge where the \mathbb{K}^* representation is localized on the image of the Tits Satake projection \mathbb{K}_{TS}^* . For instance, for the models in the $\mathfrak{f}_{4(4)}$ universality class we have:

$$\mathbb{H}_{\text{TS}}^* = \mathfrak{sl}(2, \mathbb{R})_{\mathbb{H}^*} \oplus \widehat{\mathfrak{sp}(6, \mathbb{R})} \quad (8.3)$$

and:

$$\begin{aligned} \mathbb{H}^* &\xrightarrow{\mathbb{H}_{\text{TS}}^* \oplus \mathbb{G}_{\text{subpaint}}} \text{adj } \mathfrak{sl}(2, \mathbb{R})_{\mathbb{H}^*} \oplus \text{adj } \widehat{\mathfrak{sp}(6, \mathbb{R})} \\ &\quad \oplus (\mathbf{6} | \mathcal{D}_{\text{subpaint}}) \oplus (\mathbf{6} | \mathcal{D}_{\text{subpaint}}) \\ &\quad \oplus (\mathbf{1} | \mathcal{D}_{\text{subpaint}}) \oplus (\mathbf{1} | \mathcal{D}_{\text{subpaint}}) \\ \mathbb{K}^* &\xrightarrow{\mathbb{H}_{\text{TS}}^* \oplus \mathbb{G}_{\text{subpaint}}} (2_{\mathbb{H}^*}, \mathbf{14}' | \mathbf{1}_{\text{subpaint}}) \oplus (2_{\mathbb{H}^*}, \mathbf{6} | \mathcal{D}_{\text{subpaint}}) \end{aligned} \quad (8.4)$$

and the two representations $(\mathbf{6} | \mathcal{D}_{\text{subpaint}})$ appearing in the adjoint representation of \mathbb{H}^* can be utilized to get rid of $(2_{\mathbb{H}^*}, \mathbf{6} | \mathcal{D}_{\text{subpaint}})$ appearing in the decomposition of \mathbb{K}^* .

#	$\mathbb{U}_{D=3}$	\mathbb{H}^*	\mathbb{K}^*	$\mathbb{U}_{D=4}$	rep. W	\mathbb{H}_c
1	$\mathfrak{e}_{8(8)}$	$\widehat{\mathfrak{so}^*(16)}$	128_s	$\mathfrak{e}_{7(7)}$	56	$\mathfrak{su}(8)$
2	$\mathfrak{g}_{2(2)}$	$\widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(4_{3/2}, 2_{h^*})$	$\mathfrak{sl}(2, \mathbb{R})$	$4_{3/2}$	$\mathfrak{so}(2)$
3	$\mathfrak{f}_{4(4)}$	$\widehat{\mathfrak{sp}(6, \mathbb{R})} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{14}, 2_{h^*})$	$\mathfrak{sp}(6, \mathbb{R})$	14'	$\mathfrak{u}(3)$
4	$\mathfrak{e}_{6(2)}$	$\widehat{\mathfrak{su}(3, 3)} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{20}, 2_{h^*})$	$\mathfrak{su}(3, 3)$	20	$\mathfrak{su}(3) \oplus \mathfrak{su}(3) \oplus \mathfrak{u}(1)$
5	$\mathfrak{e}_{7(-5)}$	$\widehat{\mathfrak{so}^*(12)} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{32_{spin}}, 2_{h^*})$	$\mathfrak{so}^*(12)$	32_{spin}	$\mathfrak{u}(6)$
6	$\mathfrak{e}_{8(-24)}$	$\widehat{\mathfrak{e}_{7(-25)}} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{56}, 2_{h^*})$	$\mathfrak{e}_{7(-25)}$	56	$\mathfrak{u}(6)$
7	$\mathfrak{so}(8, 3)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 1)$	$(8, 3)$	$\mathfrak{so}(6, 1) \oplus \mathfrak{sl}(2, \mathbb{R})$	(7, 2)	$\mathfrak{so}(6) \oplus \mathfrak{u}(1)$
8	$\mathfrak{so}(8, 4)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 2)$	$(8, 4)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{sl}(2, \mathbb{R})$	(8, 2)	$\mathfrak{so}(6) \oplus \mathfrak{so}(2) \oplus \mathfrak{u}(1)$
9	$\mathfrak{so}(8, 5)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 3)$	$(8, 5)$	$\mathfrak{so}(6, 3) \oplus \mathfrak{sl}(2, \mathbb{R})$	(9, 2)	$\mathfrak{so}(6) \oplus \mathfrak{so}(3) \oplus \mathfrak{u}(1)$
10	$\mathfrak{so}(8, 6)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 4)$	$(8, 6)$	$\mathfrak{so}(6, 4) \oplus \mathfrak{sl}(2, \mathbb{R})$	(10, 2)	$\mathfrak{so}(6) \oplus \mathfrak{so}(4) \oplus \mathfrak{u}(1)$
11	$\mathfrak{so}(8, 7)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 5)$	$(8, 7)$	$\mathfrak{so}(6, 5) \oplus \mathfrak{sl}(2, \mathbb{R})$	(11, 2)	$\mathfrak{so}(6) \oplus \mathfrak{so}(5) \oplus \mathfrak{u}(1)$
12	$\mathfrak{so}(8, 8)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 6)$	$(8, 8)$	$\mathfrak{so}(6, 6) \oplus \mathfrak{sl}(2, \mathbb{R})$	(12, 2)	$\mathfrak{so}(6) \oplus \mathfrak{so}(6) \oplus \mathfrak{u}(1)$
13	$\mathfrak{so}(8, 8+p)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{so}(2, 6+p)$	$(8, 8+p)$	$\mathfrak{so}(6, 6+p) \oplus \mathfrak{sl}(2, \mathbb{R})$	(12+p, 2)	$\mathfrak{so}(6) \oplus \mathfrak{so}(6+p) \oplus \mathfrak{u}(1)$
14	$\mathfrak{so}(4, 3)$	$\widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \widehat{\mathfrak{so}(2, 1)} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{2}, \widehat{3}, 2_{h^*})$	$\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 1)$	(2, 3)	$\mathfrak{so}(2) \oplus \mathfrak{u}(1)$
15	$\mathfrak{so}(4, 4)$	$\widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \widehat{\mathfrak{so}(2, 2)} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{2}, \widehat{4}, 2_{h^*})$	$\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 2)$	(2, 4)	$\mathfrak{so}(2) \oplus \mathfrak{so}(2) \oplus \mathfrak{u}(1)$
16	$\mathfrak{so}(4, 4+p)$	$\widehat{\mathfrak{sl}(2, \mathbb{R})} \oplus \widehat{\mathfrak{so}(2, 2+p)} \oplus \mathfrak{sl}(2, \mathbb{R})_{h^*}$	$(\widehat{2}, \widehat{4+p}, 2_{h^*})$	$\mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{so}(2, 2)$	(2, 4+p)	$\mathfrak{so}(2) \oplus \mathfrak{so}(2+p) \oplus \mathfrak{u}(1)$

Table 6: Table of \mathbb{H}^* subalgebras of $\mathbb{U}_{D=3}$, \mathbb{K}^* -representations and \mathbf{W} representations of $\mathbb{U}_{D=4}$ for the supergravity models based on *non-exotic* scalar symmetric spaces

What is important to stress is that, although isomorphic \mathbb{H}^* and $\mathfrak{sl}(2, \mathbb{R}) \oplus \mathbb{U}_{D=4}$ are different subalgebras of $\mathbb{U}_{D=3}$:

$$\mathbb{U}_{D=3} \supset \mathfrak{sl}(2, \mathbb{R})_{h^*} \neq \mathfrak{sl}(2, \mathbb{R})_E \subset \mathbb{U}_{D=3} \quad ; \quad \mathbb{U}_{D=3} \supset \widehat{\mathbb{U}_{D=4}} \neq \mathbb{U}_{D=4} \subset \mathbb{U}_{D=3} \quad (8.5)$$

Moreover, while the decomposition (2.20) is universal and holds true for all supergravity models, the structure (8.3) of the \mathbb{H}^* subalgebra is peculiar to the $\mathcal{N} = 2$ models. In other cases the structure of \mathbb{H}^* is different.

The reduction to the Tits Satake projection however is universal and applies to all non maximally split cases.

Indeed the remaining cases are of the form:

$$\frac{\mathbb{U}_{D=3}}{\mathbb{H}^*} = \frac{\mathrm{SO}(2+q, q+2+p)}{\mathrm{SO}(q, 2) \times \mathrm{SO}(2, q+p)} \quad (8.6)$$

#	$\mathbb{U}_{D=3}$	\mathbb{H}^*	\mathbb{K}^*	$\mathbb{U}_{D=4}$	symp. rep. W	\mathbb{H}_c
1_e	$\mathfrak{su}(\mathbf{p}+2, 2)$	$\widehat{\mathfrak{su}(\mathbf{p}+1, 1) \oplus \mathfrak{u}(1)} \oplus \mathfrak{sl}(2, \mathbb{R})_{\mathbb{H}^*}$	$(\mathbf{p}+2, \mathbf{2}_{h^*})$	$\mathfrak{su}(\mathbf{p}+1, 1) \oplus \mathfrak{u}(1)$	$\mathbf{p}+2$	$\mathfrak{su}(\mathbf{p}+1) \oplus \mathfrak{u}(1)$
2_e	$\mathfrak{su}(\mathbf{p}+2, 4)$	$\mathfrak{su}(\mathbf{p}+1, 2) \oplus \mathfrak{u}(1) \oplus \mathfrak{su}(1, 2)$	$(\mathbf{p}+3, \mathbf{3})$	$\mathfrak{su}(\mathbf{p}+1, 1) \oplus \mathfrak{u}(1)$	$\mathbf{p}+4$	$\mathfrak{su}(\mathbf{p}+1) \oplus \mathfrak{su}(3) \oplus \mathfrak{u}(1)$
3_e	$\mathfrak{e}_{6(-14)}$	$\mathfrak{so}^*(10) \oplus \mathfrak{so}(2)$	$(\mathbf{16}_s, \mathbf{2})$	$\mathfrak{su}(5, 1)$	$\mathbf{10}$	$\mathfrak{u}(5)$

Table 7: Table of \mathbb{H}^* subalgebras of $\mathbb{U}_{D=3}$, \mathbb{K}^* -representations and \mathbf{W} representations of $\mathbb{U}_{D=4}$ for the supergravity models based on *exotic* scalar symmetric spaces

leading to

$$\mathbb{K}^* = (\mathbf{q}+2, \mathbf{q}+\mathbf{p}+2) \xrightarrow{\mathfrak{so}(q,2) \oplus \mathfrak{so}(2,q+1) \oplus \mathfrak{so}(p-1)} (\mathbf{q}+2, \mathbf{q}+1, \mathbf{1}) \oplus (\mathbf{q}+2, \mathbf{1}, \mathbf{p}-1) \quad (8.7)$$

where:

$$\mathfrak{so}(q, 2) \oplus \mathfrak{so}(2, q+1) = \mathbb{H}_{\text{TS}}^* \quad (8.8)$$

$$\mathfrak{so}(p-1) = \mathbb{G}_{\text{subpaint}} \quad (8.9)$$

Considering the coset:

$$\frac{\mathbb{H}^*}{\mathbb{H}_{\text{TS}}^* \times \mathbb{G}_{\text{subpaint}}} = \frac{\text{SO}(2, \mathbf{q}+\mathbf{p})}{\text{SO}(\mathbf{q}+1, 2) \times \text{SO}(\mathbf{p}-1)} \quad (8.10)$$

we see that its $(q+3) \times (p-1)$ parameters are arranged into the

$$(\mathbf{q}+3|\mathbf{p}-1) \quad (8.11)$$

representation of $\mathfrak{so}(q+1, 2) \oplus \mathfrak{so}(p-1)$ and can be used to put to zero the component $(\mathbf{q}+2, \mathbf{1}, \mathbf{p}-1)$ in the decomposition (8.7). Note that the $\mathcal{N} = 4$ cases with more than 6 vector multiplets are covered by the above formulae by setting:

$$q = 6 \quad ; \quad p > 1 \quad (8.12)$$

Similarly the $\mathcal{N} = 2$ cases with more than 3 vector multiplets are covered by the above formulae by setting:

$$q = 2 \quad ; \quad p > 1 \quad (8.13)$$

Finally the $\mathcal{N} = 4$ cases with less than 6 vector multiplets are covered by the above formulae by setting:

$$q = n \quad ; \quad p = 6 - n \quad ; \quad n = 1, 2, 3, 4, 5 \quad (8.14)$$

9 The general structure of the $\mathbb{H}^* \oplus \mathbb{K}^*$ decomposition in the maximally split models

In the previous section we have shown that all \mathbb{H}^* orbits in the \mathbb{K}^* representation cross the locus defined by:

$$\Pi_{\text{TS}}(\mathbb{K}^*) = \mathbb{K}^* \quad (9.1)$$

where Π_{TS} is the Tits-Satake projection. In other words just as for the \mathbf{W} -representation of $\mathbb{U}_{D=4}$, it suffices to classify the orbits \mathbb{H}_{TS}^* in the \mathbb{K}_{TS}^* representation. In view of this result, in the present section we study the general structure of the $\mathbb{H}^* \oplus \mathbb{K}^*$ decomposition for maximally split algebras $\mathbb{U}_{D=3}$.

A key point in our following discussion is provided by the structure of the root system of $\mathbb{U}_{D=3}$ as described in section 6.3. The entire set of positive roots can be written as follows:

$$0 < \mathfrak{a} = \begin{cases} \alpha &= \{\bar{\alpha}, 0\} \\ \mathfrak{w} &= \{\bar{\mathfrak{w}}, \frac{1}{\sqrt{2}}\} \\ \psi &= \{0, \sqrt{2}\} \end{cases} \quad (9.2)$$

where $\bar{\alpha} > 0$ denotes the set of all positive roots of $\mathbb{U}_{D=4}$, while $\bar{\mathfrak{w}}$ denotes the complete set of weights (positive, negative and null) of the \mathbf{W} representation of $\mathbb{U}_{D=4}$. The root ψ is the highest root of the $\mathbb{U}_{D=3}$ root system and is also the root of the Ehlers subalgebra $\mathfrak{sl}(2, \mathbb{R})_E$. Accordingly, a basis of the Cartan subalgebra of $\mathbb{U}_{D=3}$ is constructed as follows:

$$\underbrace{\text{CSA}}_{\text{of } \mathbb{U}_{D=3}} = \text{span of } \left\{ \underbrace{\mathcal{H}_1, \mathcal{H}_2, \dots, \mathcal{H}_r}_{\text{CSA generators of } \mathbb{U}_{D=4}}, \underbrace{\mathcal{H}_\psi}_{\text{CSA generator of } \mathfrak{sl}(2, \mathbb{R})_E} \right\} \quad (9.3)$$

For all maximally split Lie algebras \mathbb{U} of rank $r + 1$, the maximal compact subalgebra $\mathbb{H} \subset \mathbb{U}$ is generated by:

$$T^{\mathfrak{a}} = E^{\mathfrak{a}} - E^{-\mathfrak{a}} \quad (9.4)$$

while the complementary orthogonal space \mathbb{K} is generated by

$$K^{\mathfrak{a}} = E^{\mathfrak{a}} + E^{-\mathfrak{a}} \quad (9.5)$$

$$K^I = \mathcal{H}^I \quad ; \quad I = 1, \dots, r + 1 \quad (9.6)$$

The splitting $\mathbb{H}^* \oplus \mathbb{K}^*$ is obtained by means of just one change of sign which, thanks to the structure (9.2) of the root system is consistent, namely still singles out a subalgebra.

The generators of the \mathbb{H}^* subalgebra are as follows:

$$\begin{aligned} T_{\star}^{\alpha} &= E^{\alpha} - E^{-\alpha} \\ T_{\star}^{\mathfrak{w}} &= E^{\mathfrak{w}} + E^{-\mathfrak{w}} \\ T_{\star}^{\psi} &= E^{\psi} - E^{-\psi} \end{aligned} \quad (9.7)$$

while the generators of the \mathbb{K}^* complementary subspace are as follows:

$$\begin{aligned} K_{\star}^{\alpha} &= E^{\alpha} + E^{-\alpha} \\ K_{\star}^{\mathfrak{w}} &= E^{\mathfrak{w}} - E^{-\mathfrak{w}} \\ K_{\star}^{\psi} &= E^{\psi} + E^{-\psi} \\ K^I &= \mathcal{H}^I \quad ; \quad I = 1, \dots, r + 1 \end{aligned} \quad (9.8)$$

From eq.(9.7) we see that \mathbb{H}^* contains the maximal compact subalgebra of the original $\mathbb{U}_{D=4}$ and the maximal compact subalgebra $\mathfrak{so}(2) \subset \mathfrak{sl}(2, \mathbb{R})_E$ of the Ehlers group. Using this structure we can now compare the classification of \mathbb{K}^* orbits with the classification of \mathbf{W} -orbits.

10 \mathbb{K}^\star orbits versus \mathbf{W} -orbits

In the σ -model approach the complete black hole spherically symmetric supergravity solution is obtained from two data⁸, namely the Lax operator L_0 evaluated at spatial infinity (see eq.(8.1)) and the coset representative \mathbb{L}_0 also evaluated at spatial infinity. In terms of these data one defines the matrix of conserved Noether charges:

$$Q^{Noether} = \mathbb{L}_0 L_0 \mathbb{L}_0^{-1} = \mathbb{L}(\tau) L(\tau) \mathbb{L}^{-1}(\tau) \quad (10.1)$$

from which the electromagnetic charges of the black hole, belonging to the \mathbf{W} -representation of $U_{D=4}$, can be obtained by means of the following trace:

$$\mathcal{Q}^{\mathbf{W}} = \text{Tr} \left(Q^{Noether} \mathcal{T}^{\mathbf{W}} \right) \quad (10.2)$$

where

$$\mathcal{T}^{\mathbf{W}} \propto E^{\mathbf{W}} \quad (10.3)$$

are the generators of the solvable Lie algebra corresponding to the \mathbf{W} -representation.

It is important to stress that, because of physical boundary conditions, the coset representative at spatial infinity \mathbb{L}_0 belongs to the subgroup $U_{D=4} \subset U_{D=3}$. Indeed it simply encodes the boundary values at infinity of the $D = 4$ scalar fields:

$$U_{D=3} \supset U_{D=4} \ni \mathbb{L}_0 = \exp \left[\phi_0^\alpha E^\alpha + \sum_{i=1}^r \phi_0^i \mathcal{H}_i \right] \quad (10.4)$$

Using this information in eq.(10.2) we obtain

$$\mathcal{Q}^{\mathbf{W}} = \text{Tr} \left(L_0 \mathbb{L}_0^{-1}(\phi) \mathcal{T}^{\mathbf{W}} \mathbb{L}_0(\phi) \right) = R(\phi)^{\mathbf{W}}_{\mathbf{W}'} \mathcal{Q}^{\mathbf{W}'} \quad (10.5)$$

where:

$$\mathcal{Q}^{\mathbf{W}'} = \text{Tr} \left(L_0 \mathcal{T}^{\mathbf{W}'} \right) \quad (10.6)$$

are the electromagnetic charges obtained with no scalar field dressing at infinity and

$$R(\phi)^{\mathbf{W}}_{\mathbf{W}'} \in U_{D=4} \quad (10.7)$$

is the matrix representing the group element $\mathbb{L}_0(\phi)$ in the \mathbf{W} -representation.

This result has a very significant consequence. The scalar field dressing at infinity simply rotates the charge vector along the same \mathbf{W} -orbit and is therefore irrelevant.

Hence we conclude that for each Lax operator, the \mathbf{W} -orbit of charges is completely determined and unique. The next question that was already tackled in the introduction is whether the charge-orbit \mathbf{W} is the same for all Lax operators belonging to the same \mathbf{H}^\star -orbit. As already anticipated, the answer is no and it is quite easy to produce counter examples.

Yet if we impose the condition that the Taub-NUT charge should be zero:

$$\text{Tr} \left(L_0 L_-^E \right) = 0 \quad (10.8)$$

⁸See papers [23, 25, 28, 29] for detailed explanations.

then for all Lax operators in the same H^\star , satisfying the additional constraint (10.8), the corresponding charges $Q^w = \text{Tr}(L_0 T^w)$ fall into the same \mathbf{W} -orbit.

We were not able to prove this statement, but we assert it as a *conjecture*, since we analyzed many cases and it was always true, no counter example being ever found.

In the case of multicenter non spherically symmetric solutions our conjecture appears to be true as long as we impose the condition of vanishing of the Taub-NUT current:

$$j^{TN} = 0 \quad (10.9)$$

So doing, at every pole of the involved harmonic functions, we obtain a black hole that always falls into the same \mathbf{W} -orbit.

What happens instead when the Taub-NUT current is turned on cannot be predicted in general terms at the present status of our knowledge and more study is certainly in order.

11 Conclusions and Perspectives

In this paper we examined the systematics of extremal stationary solutions of supergravity models whose scalar manifold is a symmetric coset space.

We provided a comprehensive group theoretical analysis of all such theories, organizing them in universality classes according to their Tits Satake projection. This was instrumental to our double classification of stationary solutions according to orbits of the non-compact isotropy subgroup H^\star in the three-dimensional approach, and to orbits of the symplectic \mathbf{W} -representation of the duality group $U_{D=4}$ in the four-dimensional approach. In both cases we provided full evidence that the solutions can be gauge rotated to the Tits Satake subalgebra so that classifying nilpotent orbits for the finite list of maximally split coset manifolds U_{TS}/H_{TS}^\star suffices to classify all stationary supergravity black holes. In force of this, restricting our attention to the maximally split cases, we provided the general form of the H^\star subalgebra which can be considered one of the new results attained by our paper (sect.9).

Within this general group-theoretical setup we analyzed the construction of multicenter solutions following the strategy laid down in [38, 39, 40, 41, 42, 43] which utilizes the nilpotent subalgebra singled out by every nilpotent orbit. We provided the missing link necessary to transform such a strategy into a complete constructive algorithm: such link is the general procedure described in subsection 3.2 for the transformation of the coset representative from the symmetric to the solvable gauge where the supergravity fields can be read off. We consider this another relevant result of our paper.

Next we performed a complete survey of the stationary solutions associated with the nilpotent orbits of the S^3 -model, considering the structure of charge-orbits for the corresponding black holes. In this case the $D = 4$ duality algebra is $U_{D=4} = \mathfrak{sl}(2, \mathbb{R})$ and the \mathbf{W} -representation encoding the charges \mathcal{Q} , is the $j = \frac{3}{2}$ of $\mathfrak{sl}(2, \mathbb{R})$. We found that there are the following orbits⁹:

- $\mathfrak{J}_4(\mathcal{Q}) = 0$ with stability subgroup $\Gamma \mathcal{Q} = \mathcal{Q}$ given by $\Gamma = \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$, *i.e.* by a parabolic subgroup of $SL(2, \mathbb{R})$. This orbit corresponds to the *very small black holes*.

⁹As we already mentioned the authors express their gratitude to Alessio Marrani who attracted their attention, after the first appearance of the present paper in the arXive, to paper [46] where the existence of a discrete stability subgroup for the charges of the regular BPS orbit in the S^3 model had already been found.

- $\mathfrak{I}_4(\mathcal{Q}) = 0$ with stability subgroup $\Gamma \mathcal{Q} = \mathcal{Q}$ given by $\Gamma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ *i.e.* by the identity of $\text{SL}(2, \mathbb{R})$. This orbit corresponds to the *small black holes*.
- $\mathfrak{I}_4(\mathcal{Q}) > 0$ with stability subgroup $\Gamma \mathcal{Q} = \mathcal{Q}$ given by $\Gamma = \mathbb{Z}_3 \subset \text{SL}(2, \mathbb{R})$. This orbit corresponds to the regular *BPS black holes*.
- $\mathfrak{I}_4(\mathcal{Q}) < 0$ with stability subgroup $\Gamma \mathcal{Q} = \mathcal{Q}$ given by $\Gamma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ *i.e.* by the identity of $\text{SL}(2, \mathbb{R})$. This orbit corresponds to the regular *non BPS black holes*.
- $\mathfrak{I}_4(\mathcal{Q}) > 0$ with stability subgroup $\Gamma \mathcal{Q} = \mathcal{Q}$, given by $\Gamma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ *i.e.* by the identity of $\text{SL}(2, \mathbb{R})$. This orbit corresponds to the generically singular *very large black holes*.

Up to our knowledge, this detailed structure of the S^3 model was not discussed in the literature so far and can be considered still another of our main results. The above pattern is quite inspiring and leads to the conjecture that also in other models the charges of BPS black holes might be characterized by their invariance under some suitable finite subgroup of the duality group.

In our analysis of the multicenter solutions we came to the conclusion that, within each H^* orbit, when the vanishing of the Taub-NUT current is imposed, all the black holes that are located at the various poles of the involved harmonic functions, emerge with charges always assigned to the same \mathbf{W} -orbit. Hence at vanishing Taub-NUT current the main question that motivated our paper has been answered. At least in the S^3 model there is a rigid association between the H^* orbit utilized to construct the solution and the \mathbf{W} -orbit of their charges.

Confirming this rigid association for other Tits Satake universality classes is one of the issue that emerge from our paper and that we live for future publications. Other issues raised by our results that we plan to further investigate are:

1. The appropriate interpretation of the solutions associated with higher nilpotency orbits.
2. Fitting of extremal and non extremal rotating black holes into the scheme.

12 Acknowledgements

The authors would like to express their gratitude to Sergio Ferrara and Mario Trigiante for enlightening discussions during the development of this work. Furthermore it is also a pleasure to acknowledge some remarks from and useful communications with Alessio Marrani.

A.S. Sorin acknowledges partial financial support from the RFBR Grants No. 11-02-01335-a and 11-02-12232-ofi-m-2011.

A The complete form of the fields for the $\mathcal{O}_{3|11}$ orbit

In this appendix we present the complete result for the scalar fields of the three-dimensional σ -model parameterized by four harmonic functions $\mathcal{H}_{1,2,3,4}$ in the case of the Large BPS orbit \mathcal{O}_{11}^3

$$\begin{aligned}
\mathfrak{W} &= \frac{2\sqrt{3}}{\sqrt{4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)}} \\
\text{Im}z &= \frac{\sqrt{4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)}}{2\sqrt{3}(\mathcal{H}_1^2 - \mathcal{H}_2^2 + \mathcal{H}_3^2)} \\
\text{Re}z &= \frac{\mathcal{H}_2(\mathcal{H}_3 - 3\mathcal{H}_4) + \mathcal{H}_1(\mathcal{H}_3 + 3\mathcal{H}_4)}{2\sqrt{3}(\mathcal{H}_1^2 - \mathcal{H}_2^2 + \mathcal{H}_3^2)} \tag{A.1}
\end{aligned}$$

76

$$\begin{aligned}
Z^1 &= -\frac{3\sqrt{2}(2\mathcal{H}_1^3 + 2\mathcal{H}_2\mathcal{H}_1^2 + (\mathcal{H}_3(\mathcal{H}_3 - 3\mathcal{H}_4) - 2\mathcal{H}_2^2)\mathcal{H}_1 + \mathcal{H}_2(\mathcal{H}_3(\mathcal{H}_3 + 3\mathcal{H}_4) - 2\mathcal{H}_2^2))}{4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)} \\
Z^2 &= (4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6(\mathcal{H}_4 + 1)\mathcal{H}_3 - 3\mathcal{H}_4(\mathcal{H}_4 + 2))\mathcal{H}_1^2 + 2\mathcal{H}_2(3(\mathcal{H}_3^2 + 3\mathcal{H}_4(\mathcal{H}_4 + 2)) - 4\mathcal{H}_2^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 \\
&\quad - 12\mathcal{H}_3^3(\mathcal{H}_4 + 1) + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6(\mathcal{H}_4 + 1)\mathcal{H}_3 - 3\mathcal{H}_4(\mathcal{H}_4 + 2))) \times \left(\sqrt{2}(4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 \right. \\
&\quad \left. + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)) \right)^{-1} \tag{A.2}
\end{aligned}$$

$$\begin{aligned}
Z^3 &= -\left(\sqrt{\frac{3}{2}}(4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + (3\mathcal{H}_3^2 - 2(9\mathcal{H}_4 + 1)\mathcal{H}_3 + 3(2 - 3\mathcal{H}_4)\mathcal{H}_4)\mathcal{H}_1^2 \right. \\
&\quad \left. + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2 - 2\mathcal{H}_3)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12(\mathcal{H}_3 - 1)\mathcal{H}_3^2\mathcal{H}_4 + \mathcal{H}_2^2(3\mathcal{H}_3^2 + 2(9\mathcal{H}_4 - 1)\mathcal{H}_3 - 3\mathcal{H}_4(3\mathcal{H}_4 + 2))) \right) \times \\
&\quad \times (4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 \\
&\quad + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2))^{-1} \tag{A.3}
\end{aligned}$$

$$\begin{aligned}
Z^4 &= -\frac{\sqrt{\frac{2}{3}}(2\mathcal{H}_1^3 + 6\mathcal{H}_2\mathcal{H}_1^2 + (6\mathcal{H}_2^2 - 9\mathcal{H}_4(\mathcal{H}_3 + \mathcal{H}_4))\mathcal{H}_1 + \mathcal{H}_2(2\mathcal{H}_2^2 + 9\mathcal{H}_4(\mathcal{H}_4 - \mathcal{H}_3)))}{4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)} \tag{A.4}
\end{aligned}$$

$$a = \frac{\mathcal{N}}{\mathcal{D}}$$

$$\begin{aligned}
\mathcal{N} = & 240\mathcal{H}_1^{11} + 768\mathcal{H}_2\mathcal{H}_1^{10} - 24(2\mathcal{H}_2^2 - 32\mathcal{H}_3^2 + 27\mathcal{H}_4^2 + 63\mathcal{H}_3\mathcal{H}_4)\mathcal{H}_1^9 + 4\mathcal{H}_2(8\mathcal{H}_2^2(3\mathcal{H}_4 - 70) \\
& - 3((6\mathcal{H}_4 - 187)\mathcal{H}_3^2 + 162\mathcal{H}_4\mathcal{H}_3 + 3\mathcal{H}_4^2(6\mathcal{H}_4 + 7)))\mathcal{H}_1^8 + 3(32(\mathcal{H}_4 - 15)\mathcal{H}_2^4 - 8((3\mathcal{H}_4 - 10)\mathcal{H}_3^2 \\
& - 216\mathcal{H}_4\mathcal{H}_3 + 9(\mathcal{H}_4 - 8)\mathcal{H}_4^2)\mathcal{H}_2^2 + 299\mathcal{H}_3^4 + 81\mathcal{H}_4^4 + 486\mathcal{H}_3\mathcal{H}_4^3 + 108\mathcal{H}_3^2\mathcal{H}_4^2 - 1518\mathcal{H}_3^3\mathcal{H}_4)\mathcal{H}_1^7 + \mathcal{H}_2(-32(9\mathcal{H}_4 - 68)\mathcal{H}_2^4 \\
& + 144((2\mathcal{H}_4 - 35)\mathcal{H}_3^2 + (40 - 3\mathcal{H}_4)\mathcal{H}_4\mathcal{H}_3 + \mathcal{H}_4^2(3\mathcal{H}_4 - 5))\mathcal{H}_2^2 - 162\mathcal{H}_3^2\mathcal{H}_4^2 + \mathcal{H}_3^4(2667 - 54\mathcal{H}_4) + 324\mathcal{H}_3^3(\mathcal{H}_4 - 15)\mathcal{H}_4 \\
& + 324\mathcal{H}_3\mathcal{H}_4^3(3\mathcal{H}_4 - 1) + 81\mathcal{H}_4^4(6\mathcal{H}_4 + 7))\mathcal{H}_1^6 - (96(2\mathcal{H}_4 - 27)\mathcal{H}_2^6 - 16(2\mathcal{H}_3^3 + 9(\mathcal{H}_4 - 31)\mathcal{H}_3^2 + 9\mathcal{H}_4(3\mathcal{H}_4 - 41)\mathcal{H}_3 \\
& + 27\mathcal{H}_4^2(2\mathcal{H}_4 - 3))\mathcal{H}_2^4 + 3(8\mathcal{H}_3^5 - 435\mathcal{H}_3^4 + 66\mathcal{H}_4(2\mathcal{H}_4 - 55)\mathcal{H}_3^3 + 36\mathcal{H}_4^2(3\mathcal{H}_4 + 17)\mathcal{H}_3^2 + 54\mathcal{H}_4^3(6\mathcal{H}_4 + 17)\mathcal{H}_3 + 81\mathcal{H}_4^4(4\mathcal{H}_4 + 7))\mathcal{H}_2^2 \\
& - 18\mathcal{H}_3^2(25\mathcal{H}_3^4 - 272\mathcal{H}_4\mathcal{H}_3^3 + 216\mathcal{H}_4^2\mathcal{H}_3^2 + 180\mathcal{H}_4^3\mathcal{H}_3 + 27\mathcal{H}_4^4))\mathcal{H}_1^5 + \mathcal{H}_2(64(3\mathcal{H}_4 - 14)\mathcal{H}_2^6 + 8(8\mathcal{H}_3^3 + (411 - 27\mathcal{H}_4)\mathcal{H}_3^2 \\
& + 18\mathcal{H}_4(6\mathcal{H}_4 - 47)\mathcal{H}_3 - 81(\mathcal{H}_4 - 3)\mathcal{H}_4^2)\mathcal{H}_2^4 + 3(-16\mathcal{H}_3^5 + (18\mathcal{H}_4 - 1271)\mathcal{H}_3^4 - 8\mathcal{H}_4(45\mathcal{H}_4 - 514)\mathcal{H}_3^3 + 18\mathcal{H}_4^2(6\mathcal{H}_4 + 17)\mathcal{H}_3^2 \\
& - 216\mathcal{H}_4^3(3\mathcal{H}_4 - 1)\mathcal{H}_3 + 27\mathcal{H}_4^4(6\mathcal{H}_4 + 23))\mathcal{H}_2^2 + 18\mathcal{H}_3^2(77\mathcal{H}_3^4 + 2\mathcal{H}_4(6\mathcal{H}_4 - 149)\mathcal{H}_3^3 - 108\mathcal{H}_4^2\mathcal{H}_3^2 + 6\mathcal{H}_4^3(6\mathcal{H}_4 - 41)\mathcal{H}_3 - 81\mathcal{H}_4^4))\mathcal{H}_1^4 \\
& + 3(16(4\mathcal{H}_4 - 39)\mathcal{H}_2^8 - 24((3\mathcal{H}_4 - 70)\mathcal{H}_3^2 + 4\mathcal{H}_4(3\mathcal{H}_4 - 10)\mathcal{H}_3 + \mathcal{H}_4^2(9\mathcal{H}_4 - 20))\mathcal{H}_2^6 + (8\mathcal{H}_3^5 - (30\mathcal{H}_4 + 1423)\mathcal{H}_3^4 \\
& + 2\mathcal{H}_4(144\mathcal{H}_4 - 1361)\mathcal{H}_3^3 + 36\mathcal{H}_4^2(3\mathcal{H}_4 + 17)\mathcal{H}_3^2 + 54\mathcal{H}_4^3(12\mathcal{H}_4 - 1)\mathcal{H}_3 + 27\mathcal{H}_4^4(6\mathcal{H}_4 - 7))\mathcal{H}_2^4 - 6\mathcal{H}_3^2(\mathcal{H}_3^5 - 2(3\mathcal{H}_4 + 29)\mathcal{H}_3^4 \\
& + 4\mathcal{H}_4(3\mathcal{H}_4 - 103)\mathcal{H}_3^3 - 18(\mathcal{H}_4 - 24)\mathcal{H}_4^2\mathcal{H}_3^2 + 3\mathcal{H}_4^3(9\mathcal{H}_4 + 28)\mathcal{H}_3 - 54\mathcal{H}_4^4)\mathcal{H}_2^2 + 3\mathcal{H}_3^4(9\mathcal{H}_3^4 - 242\mathcal{H}_4\mathcal{H}_3^3 + 468\mathcal{H}_4^2\mathcal{H}_3^2 \\
& + 234\mathcal{H}_4^3\mathcal{H}_3 + 27\mathcal{H}_4^4))\mathcal{H}_1^3 - \mathcal{H}_2(128\mathcal{H}_2^8 + 16(2\mathcal{H}_3^3 - 9\mathcal{H}_3^2 + 9\mathcal{H}_4(3\mathcal{H}_4 - 28)\mathcal{H}_3 - 27(\mathcal{H}_4 - 3)\mathcal{H}_4^2)\mathcal{H}_2^6 \\
& + (-48\mathcal{H}_3^5 - 441\mathcal{H}_3^4 - 12\mathcal{H}_4(63\mathcal{H}_4 - 905)\mathcal{H}_3^3 + 162\mathcal{H}_4^2(2\mathcal{H}_4 + 3)\mathcal{H}_3^2 - 324\mathcal{H}_4^3(3\mathcal{H}_4 - 1)\mathcal{H}_3 + 81\mathcal{H}_4^4(12\mathcal{H}_4 + 19))\mathcal{H}_2^4 \\
& + 18\mathcal{H}_3^2(\mathcal{H}_3^5 + 38\mathcal{H}_3^4 + 2\mathcal{H}_4(9\mathcal{H}_4 - 256)\mathcal{H}_3^3 - 216\mathcal{H}_4^2\mathcal{H}_3^2 + 3\mathcal{H}_4^3(15\mathcal{H}_4 - 88)\mathcal{H}_3 - 54\mathcal{H}_4^4)\mathcal{H}_2^2 \\
& - 27\mathcal{H}_3^4(9\mathcal{H}_3^4 - 92\mathcal{H}_4\mathcal{H}_3^3 - 78\mathcal{H}_4^2\mathcal{H}_3^2 - 132\mathcal{H}_4^3\mathcal{H}_3 - 27\mathcal{H}_4^4))\mathcal{H}_1^2 + ((528 - 96\mathcal{H}_4)\mathcal{H}_2^{10} - 8(4\mathcal{H}_3^3 - 6(3\mathcal{H}_4 - 37)\mathcal{H}_3^2 \\
& + 27(3 - 2\mathcal{H}_4)\mathcal{H}_4\mathcal{H}_3 + 81\mathcal{H}_4^2)\mathcal{H}_2^8 + 3(16\mathcal{H}_3^5 + (30\mathcal{H}_4 + 737)\mathcal{H}_3^4 + 6(123 - 34\mathcal{H}_4)\mathcal{H}_4\mathcal{H}_3^3 - 108\mathcal{H}_4^2\mathcal{H}_3^2 - 162\mathcal{H}_4^3(2\mathcal{H}_4 - 3)\mathcal{H}_3 \\
& + 81\mathcal{H}_4^4(2\mathcal{H}_4 + 3))\mathcal{H}_2^6 - 6\mathcal{H}_3^2(3\mathcal{H}_3^5 + (34\mathcal{H}_4 + 201)\mathcal{H}_3^4 - 36(\mathcal{H}_4 - 13)\mathcal{H}_4\mathcal{H}_3^3 + 54(\mathcal{H}_4 - 12)\mathcal{H}_4^2\mathcal{H}_3^2 \\
& - 27\mathcal{H}_4^3(5\mathcal{H}_4 - 16)\mathcal{H}_3 + 243\mathcal{H}_4^4)\mathcal{H}_2^4 + 9\mathcal{H}_3^4((8\mathcal{H}_4 + 27)\mathcal{H}_3^4 + 174\mathcal{H}_4\mathcal{H}_3^3 + 12\mathcal{H}_4^2(2\mathcal{H}_4 - 39)\mathcal{H}_3^2 + 90\mathcal{H}_4^3\mathcal{H}_3 + 81\mathcal{H}_4^4)\mathcal{H}_2^2 \\
& - 324\mathcal{H}_3^7\mathcal{H}_4(\mathcal{H}_3^2 - 4\mathcal{H}_4\mathcal{H}_3 - \mathcal{H}_4^2))\mathcal{H}_1 + 3\mathcal{H}_2(\mathcal{H}_2^2 - \mathcal{H}_3^2)^2(64\mathcal{H}_2^6 - 12(7\mathcal{H}_3^2 + 30\mathcal{H}_4\mathcal{H}_3 - 9\mathcal{H}_4^2)\mathcal{H}_2^4 + 3(9\mathcal{H}_3^4 + 136\mathcal{H}_4\mathcal{H}_3^3 \\
& + 90\mathcal{H}_4^2\mathcal{H}_3^2 - 27\mathcal{H}_4^4)\mathcal{H}_2^2 - 108\mathcal{H}_3^3\mathcal{H}_4(\mathcal{H}_3 + \mathcal{H}_4)^2)
\end{aligned} \tag{A.6}$$

$$\begin{aligned}
\mathcal{D} = & 3\sqrt{3}(\mathcal{H}_1^2 - \mathcal{H}_2^2 + \mathcal{H}_3^2)^2(4\mathcal{H}_1^4 + 8\mathcal{H}_2\mathcal{H}_1^3 + 3(\mathcal{H}_3^2 - 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2)\mathcal{H}_1^2 + 2\mathcal{H}_2(-4\mathcal{H}_2^2 + 3\mathcal{H}_3^2 + 9\mathcal{H}_4^2)\mathcal{H}_1 - 4\mathcal{H}_2^4 \\
& - 12\mathcal{H}_3^3\mathcal{H}_4 + 3\mathcal{H}_2^2(\mathcal{H}_3^2 + 6\mathcal{H}_4\mathcal{H}_3 - 3\mathcal{H}_4^2))^2
\end{aligned} \tag{A.7}$$

References

- [1] S. Ferrara, R. Kallosh, *Supersymmetry and attractors*, Phys. Rev. **D54**, 1514-1524 (1996). [hep-th/9602136].
S. Ferrara, R. Kallosh, A. Strominger, *$N=2$ extremal black holes* Phys. Rev. **D52** (1995) 5412-5416. [hep-th/9508072].
- [2] A. Strominger and C. Vafa, *Microscopic Origin of the Bekenstein-Hawking Entropy*, Phys. Lett. B **379** (1996) 99 [arXiv:hep-th/9601029].
- [3] G. W. Gibbons, R. Kallosh, B. Kol, *Moduli, scalar charges, and the first law of black hole thermodynamics*, Phys. Rev. Lett. **77** (1996) 4992-4995. [hep-th/9607108].
- [4] M. Bertolini, P. Fre, R. Iengo and C. A. Scrucca, *Black holes as D3-branes on Calabi-Yau threefolds* Phys. Lett. B **431** (1998) 22 [arXiv:hep-th/9803096].
- [5] K. Goldstein, N. Iizuka, R. P. Jena and S. P. Trivedi, *Non-supersymmetric attractors* Phys. Rev. D **72**, 124021 (2005) [arXiv:hep-th/0507096].
P. K. Tripathy and S. P. Trivedi, *Non-Supersymmetric Attractors in String Theory* JHEP **0603**, 022 (2006) [arXiv:hep-th/0511117].
R. Kallosh, *New Attractors* JHEP **0512**, 022 (2005) [arXiv:hep-th/0510024].
A. Giryavets, *New attractors and area codes* JHEP **0603**, 020 (2006) [arXiv:hep-th/0511215].
R. Kallosh, N. Sivanandam and M. Soroush, *The non-BPS black hole attractor equation* JHEP **0603**, 060 (2006) [arXiv:hep-th/0602005].
S. Bellucci, S. Ferrara and A. Marrani, *On some properties of the attractor equations* Phys. Lett. B **635**, 172 (2006) [arXiv:hep-th/0602161].
S. Bellucci, S. Ferrara, M. Gunaydin, A. Marrani, *Charge orbits of symmetric special geometries and attractors*, Int. J. Mod. Phys. **A21** (2006) 5043-5098. [hep-th/0606209]. L. Andrianopoli, R. D'Auria, S. Ferrara, M. Trigiante, *Extremal black holes in supergravity*, Lect. Notes Phys. **737** (2008) 661-727. [hep-th/0611345].
- [6] M. Gunaydin, A. Neitzke, B. Pioline and A. Waldron, *BPS black holes, quantum attractor flows and automorphic forms*, Phys. Rev. **D73** (2006) 084019; B. Pioline, *Lectures on black holes, topological strings and quantum attractors*, Class. Quant. Grav. **23** (2006) S981. [hep-th/0607227]; M. Gunaydin, A. Neitzke, B. Pioline, A. Waldron, *Quantum Attractor Flows*, JHEP **0709** (2007) 056;
- [7] G. W. Moore, *Arithmetic and attractors*, arXiv:hep-th/9807087. L. Andrianopoli, R. D'Auria, S. Ferrara *et al.*, *$E(7)(7)$ duality, BPS black hole evolution and fixed scalars* Nucl. Phys. **B509** (1998) 463-518. [hep-th/9707087].
L. Andrianopoli, R. D'Auria, S. Ferrara *et al.*, *Solvable Lie algebras in type IIA, type IIB and M theories* Nucl. Phys. **B493** (1997) 249-280. [hep-th/9612202].
L. Andrianopoli, R. D'Auria, S. Ferrara *et al.*, *RR scalars, U duality and solvable Lie algebras*, Nucl. Phys. **B496** (1997) 617-629. [hep-th/9611014].
G. Arcioni, A. Ceresole, F. Cordaro *et al.*, *$N=8$ BPS black holes with $1/2$ or $1/4$ supersymmetry and solvable Lie algebra decompositions* Nucl. Phys. **B542** (1999) 273-307. [hep-th/9807136].
M. J. Duff, J. T. Liu, J. Rahmfeld, *Four Dimensional String/String/String Triality*, Nucl.Phys.

- B459 (1996) 125, hep-th/9508094;
R. Kallosh, M. Shmakova, W. K. Wong, *Freezing of Moduli by $N=2$ Dyons*, Phys.Rev. D54 (1996) 6284, hep-th/9607077;
K. Behrndt, R. Kallosh, J. Rahmfeld, M. Smachova and W. K. Wond, *STU Black holes and string triality*, hep-th/9608059;
G. Lopes Cardoso, D. Lust and T. Mohaupt, *Modular Symmetries of $N=2$ Black Holes*, Phys.Lett. B388 (1996) 266, hep-th/9608099;
- [8] M. Bertolini, M. Trigiante, *Regular BPS black holes: Macroscopic and microscopic description of the generating solution*, Nucl. Phys. **B582** (2000) 393-406. [hep-th/0002191]. M. Bertolini, M. Trigiante, *Regular RR and NS NS BPS black holes*, Int. J. Mod. Phys. **A15** (2000) 5017. [hep-th/9910237].
- [9] Older references on susy black holes:
G. Gibbons, in “*Unified theories of Elementary Particles. Critical Assessment and Prospects*”, Proceedings of the Heisemberg Symposium, München, West Germany, 1981, ed. by P. Breitenlohner and H. P. Dürr, Lecture Notes in Physics Vol. 160 (Springer-Verlag, Berlin, 1982);
G. W. Gibbons and C. M. Hull, Phys. Lett. **109B** (1982) 190;
G. W. Gibbons, in “*Supersymmetry, Supergravity and Related Topics*”, Proceedings of the XVth GIFT International Physics, Girona, Spain, 1984, ed. by F. del Aguila, J. de Azcárraga and L. Ibáñez, (World Scientific, 1995), pag. 147;
R. Kallosh, A. Linde, T. Ortin, A. Peet and A. Van Proeyen, Phys. Rev. **D46** (1992) 5278;
R. Kallosh, T. Ortin and A. Peet, Phys. Rev. **D47** (1993) 5400; R. Kallosh, Phys. Lett. **B282** (1992) 80;
R. Kallosh and A. Peet, Phys. Rev. **D46** (1992) 5223;
A. Sen, Nucl. Phys. **B440** (1995) 421; Phys. Lett. **B303** (1993) 221; Mod. Phys. Lett. **A10** (1995) 2081;
J. Schwarz and A. Sen, Phys. Lett. **B312** (1993) 105;
- [10] R. Kallosh, T. Ortin, “*Charge quantization of Axion Dilaton black-holes*” Phys. rev. **D48** (1993) 742, hep-th 9302109,
E. Bergshoeff, R. Kallosh, T. Ortin “*Stationary Axion Dilaton solutions and supersymmetry* Nuc. Phys. **B478** (1996) 156, hep-th 9605059.
- [11] M. Cvetič, J. Louis, B. A. Ovrut, *A String Calculation of the Kahler Potentials for Moduli of $Z(N)$ Orbifolds*, Phys. Lett. **B206** (1988) 227.
S. Ferrara, P. Fre and P. Soriani, *On the moduli space of the $T^{**6} / Z(3)$ orbifold and its modular group*, Class. Quant. Grav. **9** (1992) 1649 [arXiv:hep-th/9204040].
G. Lopes Cardoso, D. Lust and T. Mohaupt, *Moduli spaces and target space duality symmetries in $(0,2)$ $Z(N)$ orbifold theories with continuous Wilson lines* Nucl. Phys. B **432** (1994) 68 [arXiv:hep-th/9405002].
- [12] *Pioneering papers on vector multiplet coupling to Supergravity:*
K. S. Stelle and P. C. West, Phys. Lett. B **77** (1978) 376; Nucl. Phys. B **145** (1978) 175;
B. de Wit, P.G. Lauwers, R. Philippe, Su S.Q. and A. Van Proeyen, Phys. Lett. 134B (1984) 37.

- S. J. Gates, Nucl. Phys. B238 (1984) 349.
 B. de Wit and A. Van Proeyen, Nucl. Phys. B245 (1984) 89;
 E. Cremmer, C. Kounnas, A. Van Proeyen, J. P. Derendinger, S. Ferrara, B. de Wit and L. Girardello, Nucl. Phys. B250 (1985) 385.
Definition of Special Kähler Geometry:
 A. Strominger, Comm. Math. Phys. 133 (1990) 163.
 L. Castellani, R. D'Auria and S. Ferrara, Phys. Lett. 241B (1990) 57; Class. Quantum Grav. 7 (1990) 1767.
 R. D'Auria, S. Ferrara and P. Fré, Nucl. Phys. B359 (1991) 705.
- [13] L. Andrianopoli, M. Bertolini, A. Ceresole, R. D'Auria, S. Ferrara, P. Fre and T. Magri, *$N = 2$ supergravity and $N = 2$ super Yang-Mills theory on general scalar manifolds: Symplectic covariance, gaugings and the momentum map*, J. Geom. Phys. **23** (1997) 111 [arXiv:hep-th/9605032].
- [14] *Homogeneous Special Kähler Geometry:*
 B. de Wit, A. Van Proeyen, *Special geometry, cubic polynomials and homogeneous quaternionic spaces*, Commun.Math.Phys.149:307-334 (1992), hep-th/9112027;
 B. de Wit, A. Van Proeyen, *Broken sigma model isometries in very special geometry*, Phys.Lett.B293:94-99 (1992), hep-th/9207091; B. de Wit, F. Vanderseypen, A. Van Proeyen, *Symmetry structure of special geometries*, Nucl.Phys. B400 (1993) 463-524, hep-th/9210068;
 B. de Wit, A. Van Proeyen, *Hidden symmetries, special geometry and quaternionic manifolds*, Int.J.Mod.Phys. D3 (1994) 31-48, hep-th/9310067; .
- [15] C. M. Hull and P. K. Townsend, *Unity of superstring dualities*, Nucl. Phys. B **438** (1995) 109 [arXiv:hep-th/9410167]; L. Borsten, D. Dahanayake, M. J. Duff and W. Rubens, *Black holes admitting a Freudenthal dual*, Phys. Rev. D **80** (2009) 026003 [arXiv:0903.5517 [hep-th]].
- [16] A. Strominger, *Black hole entropy from near horizon microstates* JHEP **9802** (1998) 009.[hep-th/9712251].
 G. T. Horowitz, J. M. Maldacena, A. Strominger, *Nonextremal black hole microstates and U duality* Phys. Lett. **B383** (1996) 151-159. [hep-th/9603109].
 A. Dabholkar, *Microstates of nonsupersymmetric black holes* Phys. Lett. **B402** (1997) 53-58. [hep-th/9702050].
 F. Larsen, *A String model of black hole microstates* Phys. Rev. **D56** (1997) 1005-1008.[hep-th/9702153].
 A. Ghosh, P. Mitra, *Counting of black hole microstates* Indian J. Phys. **80** (2006) 867. [gr-qc/0603029]; J. M. Maldacena, A. Strominger, E. Witten, *Black hole entropy in M theory*, JHEP **9712** (1997) 002. [hep-th/9711053]. M. Bertolini, M. Trigiante, *Microscopic entropy of the most general four-dimensional BPS black hole*, JHEP **0010** (2000) 002. [hep-th/0008201].
- [17] G. Lopes Cardoso, A. Ceresole, G. Dall'Agata, J. M. Oberreuter and J. Perz, *First-order flow equations for extremal black holes in very special geometry*, JHEP **0710** (2007) 063 [arXiv:0706.3373 [hep-th]].
- [18] D. Gaiotto, W. W. Li and M. Padi, *Non-Supersymmetric Attractor Flow in Symmetric Spaces*, JHEP **12** (2007) 093,

- [19] A. Ceresole and G. Dall'Agata, *Flow Equations for Non-BPS Extremal Black Holes* JHEP **0703**, 110 (2007) [arXiv:hep-th/0702088].
- [20] A. Ceresole, G. Dall'Agata, S. Ferrara and A. Yeranyan, *First order flows for $N=2$ extremal black holes and duality invariants* Nucl. Phys. B **824**, 239 (2010) [arXiv:0908.1110 [hep-th]].
G. Bossard, Y. Michel and B. Pioline, *Extremal black holes, nilpotent orbits and the true fake superpotential* JHEP **1001**, 038 (2010) [arXiv:0908.1742 [hep-th]].
A. Ceresole, G. Dall'Agata, S. Ferrara and A. Yeranyan, *Universality of the superpotential for $d = 4$ extremal black holes*, Nucl. Phys. B **832**, 358 (2010) [arXiv:0910.2697 [hep-th]].
- [21] L. Andrianopoli, R. D'Auria, E. Orazi and M. Trigiante, *First Order Description of Black Holes in Moduli Space* JHEP **0711**, 032 (2007) [arXiv:0706.0712 [hep-th]];
L. Andrianopoli, R. D'Auria, E. Orazi and M. Trigiante, *First Order Description of $D=4$ static Black Holes and the Hamilton-Jacobi equation* Nucl. Phys. B **833**, 1 (2010) [arXiv:0905.3938 [hep-th]];
L. Andrianopoli, R. D'Auria, S. Ferrara, M. Trigiante, *Fake Superpotential for Large and Small Extremal Black Holes*, JHEP **1008** (2010) 126. [arXiv:1002.4340 [hep-th]].
- [22] E. Bergshoeff, W. Chemissany, A. Ploegh, M. Trigiante and T. Van Riet, *Generating Geodesic Flows and Supergravity Solutions*, Nucl. Phys. B **812** (2009) 343 [arXiv:0806.2310].
- [23] P. Fré and A.S. Sorin, *The Integration Algorithm for Nilpotent Orbits of G/H^* Lax systems: for Extremal Black Holes* [arXiv:0903.3771]. P. Fré and A.S. Sorin, *The arrow of time and the Weyl group: all supergravity billiards are integrable*, Nucl. Phys. B **815** (2009) 430 [arXiv:0710.1059].
P. Fre and A. S. Sorin, "Integrability of supergravity billiards and the generalized Toda lattice equation," Nucl. Phys. B **733** (2006) 334 [hep-th/0510156].
- [24] W. Chemissany, P. Fre and A. S. Sorin, *The Integration Algorithm of Lax equation for both Generic Lax matrices and Generic Initial Conditions*, Nucl. Phys. B **833** (2010) 220 [arXiv:0904.0801 [hep-th]].
- [25] W. Chemissany, P. Fre, J. Rosseel, A. S. Sorin, M. Trigiante and T. Van Riet, *Black holes in supergravity and integrability*, JHEP **1009** (2010) 080.
- [26] P. Fré, A.S. Sorin, *Supergravity Black Holes and Billiards and Liouville integrable structure of dual Borel algebras*, JHEP **03** (2010) 066 [arXiv:0903.2559].
- [27] W. Chemissany, J. Rosseel, M. Trigiante, T. Van Riet, *The full integration of black hole solutions to symmetric supergravity theories*, Nucl. Phys. B **830** (2010) 391 [arXiv:0903.2777].
- [28] P. Frè, A. S. Sorin and M. Trigiante, *Integrability of Supergravity Black Holes and New Tensor Classifiers of Regular and Nilpotent Orbits*, JHEP **1204** (2012) 015, [arXiv:1103.0848 [hep-th]].
- [29] P. Fre, A. S. Sorin and M. Trigiante, *Black Hole Nilpotent Orbits and Tits Satake Universality Classes*, arXiv:1107.5986 [hep-th].
- [30] A. Ceresole, S. Ferrara, A. Marrani, *Small $N=2$ Extremal Black Holes in Special Geometry*, [arXiv:1006.2007v1]

- [31] A. Ceresole, G. Dall’Agata, S. Ferrara, A. Yeranyan, *First order flows for $N=2$ extremal black holes and duality invariants*, [arXiv:0908.1110v2]
- [32] P. Breitenlohner, D. Maison and G. W. Gibbons, *Four-dimensional black holes from Kaluza-Klein theories*, Commun. Math. Phys. **120** (1988) 295;
- [33] S. Ferrara, S. Sabharwal, *Quaternionic manifolds for type II superstring vacua of Calabi-Yau spaces*, Nucl.Phys.B332:317 (1990)
S. Cecotti, *Homogeneous Kahler Manifolds And T Algebras In $N=2$ Supergravity And Superstrings,* Commun. Math. Phys. **124** (1989) 23-55.
R. D’Auria, S. Ferrara, M. Trigiante, *C - map, very special quaternionic geometry and dual Kahler spaces* Phys. Lett. **B587** (2004) 138-142. [hep-th/0401161].
- [34] The main reference on this issue is D. H. Collingwood and W.M. McGovern, *Nilpotent Orbits in Semisimple Lie Algebras*, Van Nostrand Reinhold 1993.
- [35] S. Kim, J. Lindman Hörnlund, J. Palmkvist and A. Virmani, *Extremal solutions of the S^3 model and nilpotent $G_{2(2)}$ orbits* [arXiv:1004.5242v2].
- [36] P. Frè , F. Gargiulo, J. Rosseel *et al.*, *Tits-Satake projections of homogeneous special geometries,* Class. Quant. Grav. **24** (2007) 27-78. [hep-th/0606173].
- [37] W. Chemissany, P. Giaccone, D. Ruggeri and M. Trigiante, *Black hole solutions to the F_4 -model and their orbits (I)*, arXiv:1203.6338 [hep-th].
- [38] G. Bossard and H. Nicolai, *Multi-black holes from nilpotent Lie algebra orbits*, Gen. Rel. Grav. **42** (2010) 509 [arXiv:0906.1987 [hep-th]].
- [39] G. Bossard, H. Nicolai, K. S. Stelle, *Universal BPS structure of stationary supergravity solutions*, JHEP **0907** (2009) 003.
- [40] G. Bossard, *Extremal black holes and nilpotent orbits*, arXiv:0910.0689 [hep-th].
- [41] G. Bossard, *The Extremal black holes of $N=4$ supergravity from $so(8,2+n)$ nilpotent orbits*, Gen. Rel. Grav. **42** (2010) 539
- [42] G. Bossard and C. Ruef, *Interacting non-BPS black holes*, Gen. Rel. Grav. **44** (2012) 21 [arXiv:1106.5806 [hep-th]].
- [43] G. Bossard, *Octonionic black holes* arXiv:1203.0530 [hep-th].
- [44] P. Fre’, F. Gargiulo, K. Rulik and M. Trigiante, *The general pattern of Kac Moody extensions in supergravity and the issue of cosmic billiards*, Nucl. Phys. B **741** (2006) 42 [arXiv:hep-th/0507249].
- [45] K. Behrndt, D. Lust, W. A. Sabra, “Stationary solutions of $N=2$ supergravity,” Nucl. Phys. **B510** (1998) 264-288. [hep-th/9705169];
- [46] L. Borsten, M. J. Duff, S. Ferrara, A. Marrani and W. Rubens, *Small Orbits*, Phys. Rev. D **85** (2012) 086002 [arXiv:1108.0424 [hep-th]].